

Melbourne Veterinary School

Structure and Function of the Kidney – Autoregulation, Glomerular Filtration and Urine Composition

Jenni Bauquier

jbauquier@unimelb.edu.au












Intended Learning Outcomes


- Describe how glomerular filtration occurs, why it is important and the factors that regulate it
- Explain the processes of autoregulation and the factors and hormones that regulate it
- Explain how GFR can be measured and used as an assessment of renal function
- Describe how urine is formed and modified through glomerular filtration, tubular reabsorption and secretion
- Describe the composition of normal urine
- Describe transport mechanisms responsible for sodium reabsorption by the nephron
- Explain coupling of sodium and water reabsorption in the proximal tubule
- Describe how organic ions are secreted and why this is important

Glomerular filtration

- High renal blood flow → large volumes filtered through glomeruli
- Most reabsorbed
- Filtrate/ultrafiltrate very similar composition to plasma
 - Minus protein and protein-bound substances
- Flow of fluid from plasma into Bowman's space over time = glomerular filtration rate




Glomerular filtration rate



- Mainly determined by:
 - Hydrostatic and colloid osmotic forces across glomerular membrane
 - = net filtration pressure
 - Capillary filtration coefficient (K_f)
 - = permeability & surface area of glomerular capillaries
 - GFR = K_f x net filtration pressure

Measuring GFR



- Can't easily measure GFR, therefore we estimate it by measuring other things that are affected by GFR
- Typically these substances are filtered at the glomerulus but don't undergo any reabsorption or secretion in the tubules

Measuring/estimating GFR


- Endogenous substances produced at a relatively constant rate
 - Creatinine
 - Blood urea nitrogen (dogs and cats)
- These are indirect estimations of GFR. Insensitive ~75% of kidney function lost before they increase
- Creatinine measurement in plasma and urine to calculate eGFR
 - Creatinine clearance
- More complex testing involving administration of a substance that is not reabsorbed or secreted, measuring plasma and urine concentrations to calculate eGFR

Glomerular filtration rate


- Kept relatively constant despite changes in BP, intravascular volume, etc.
- How?
 - Can't change permeability and surface area of capillaries (in health)
 - PRESSURE

Glomerular filtration rate

- Bowman's capsule pressure and glomerular colloid osmotic pressure
 OPPOSE filtration
- Glomerular hydrostatic pressure
 DRIVES filtration
 - This is the easiest factor to change
 - Controlled by:
 - Arterial pressure
 - Afferent arteriolar resistance
 - Efferent arteriolar resistance




What could possibly go wrong?



Plasma creatinine concentration = 926 μ mol/L; reference range 40-80 μ mol/L

Physiological control of GFR


- Mostly changes hydrostatic pressure in the glomerular capillaries
 - Influenced by:
 - Arterial blood pressure
 - Afferent arteriolar resistance
 - Efferent arteriolar resistance
- Mechanisms:
 - Sympathetic nervous system
 - Autacoid/hormonal control
 - Tubuloglomerular feedback (autoregulation)



How do the arterioles know what to do?

- \uparrow BP \rightarrow afferent arteriolar constriction $\rightarrow \downarrow$ glomerular hydrostatic pressure and \downarrow filtration at the glomerulus
- ↓ BP → efferent arteriolar constriction → ↑
 glomerular hydrostatic pressure and ↑ filtration at
 the glomerulus
- However, if efferent arterioles are severely constricted, glomerular filtration actually decreases:
 - Glomerular colloid osmotic pressure increases
 - Overall renal blood flow decreases





Sympathetic nervous system

- All renal blood vessels innervated by sympathetic nerves
- Only strong sympathetic activation will cause constriction of renal arterioles
- Most important in severe, acute disturbances, e.g. severe haemorrhage, defense reaction

Autacoid/hormonal control

- Adrenaline/epinephrine and noradrenaline/norepinephrine
 - Same effect as SNS activation
 - Constrict renal arterioles; ↓ GFR
- (Endothelin; ↓ GFR)
- Angiotensin II
 - Produced when BP decreases
 - Preferentially constricts efferent arteriole → maintenance of normal GFR
- Nitric oxide
 - Vasodilatory,

 GFR
- Prostaglandins
 - PGE₂ & PGI₂
 - Vasodilatory, oppose vasoconstriction, maintenance of normal GFR


Autacoid/hormonal control — what could possibly go wrong?









Tubuloglomerular feedback



- Juxtaglomerular apparatus/complex
 - Juxtaglomerular cells
 - Macula densa
- Macula densa senses Na⁺ & Cl⁻ concentration in distal tubule





Tubular reabsorption and secretion

- Large quantities, highly selective
- Small changes in GFR or tubular reabsorption/secretion can mean huge changes in overall urinary excretion
- MOST of what is filtered at the glomerulus is reabsorbed
- Tubules fine-tune reabsorption and secretion


Tubular reabsorption and secretion

• % filtered load reabsorbed of different substances:

Substance	% filtered load reabsorbed	
Glucose	100	
Bicarbonate	>99.9	
Na ⁺	99.4	
K ⁺	87.8	
Cl ⁻	99.1	
Urea	50	
Creatinine	0	


Tubular reabsorption and secretion

- Substances have to move between tubule and capillaries
- Several different mechanisms as to how tubules reabsorb and secrete substances
 - Active or passive
 - Paracellular or transcellular
- Active transport
 - Primary or secondary



Coupling Na⁺ and water reabsorption

- Active transport of sodium out of tubule creates osmotic gradient
- Drives water reabsorption
- Na⁺ main driver as greatest quantity
- Parts of tubule that are highly permeable to water
- Water moves through aquaporins and tight junctions



Passive diffusion of solutes

- Negative ions diffuse along electrical potential gradients
 - Active transport of Na⁺ \rightarrow paracellular diffusion of Cl⁻, (urea)
- Urea permeates tubule less readily than water
 - 50% reabsorbed, 50% excreted in urine
- Creatinine
 - Large molecule, does not cross tubule
 - No reabsorption


Proximal tubule


Reabsorption

- ≈ 65% filtered Na⁺, Cl⁻ and water
- Co-transport of Na⁺ with glucose and amino acids, Cl⁻
- Counter-transport of Na⁺ with H⁺ secretion
- HCO₃-, K⁺, water also reabsorbed

Secretion


- H+
- Organic acids and bases
- Catecholamines






Klein; Cunningham 6th Edition, 2020

Proximal Tubule Cell






Gattineni & Baum, 2015

Proximal tubule

Glucose

- Transport maximum filtered load exceeds ability to reabsorb
- Glucose will appear in urine before all nephrons at transport maximum
- Plasma glucose concentration 9.0-10.0 mmol/L
- Glucose in tubular fluid keeps osmotic gradient – more water stays in tubule











Thin descending loop of Henle

- ≈20% filtered water reabsorbed
- Simple diffusion only
- Highly permeable to water
- Moderately permeable to most solutes
 - Urea
 - Na⁺
- Main effect is movement of water out of the tubule











Thin ascending loop of Henle

- Impermeable to water
- Very small amount of solute reabsorption






Bachmann & Kriz, 1999

Thick ascending loop of Henle


Reabsorption

- ≈25% filtered Na⁺, K⁺ and Cl⁻ reabsorbed
- Na⁺, Cl⁻, K⁺ co-transport (NKCC2 transporter)
- Secretion
 - Also Na⁺-H⁺ countertransport; H⁺ secretion
- Impermeable to water




Bachmann & Kriz, 1999

Thick Ascending Limb Cell




Distal tubule – first half


- Macula densa
 - No role in reabsorption/secretion
 - Detection of Na⁺ and Cl⁻ concentrations



Distal tubule – first half


- Reabsorption
 - ≈5% filtered Na⁺ and Cl⁻ reabsorbed
 - Similar functions to thick ascending loop of Henle
- Also impermeable to water






Bachmann & Kriz, 1999

Distal Convoluted Tubule Cell



Distal tubule – second half & cortical collecting tubules

- Principal cells
 - Reabsorption Na⁺ (aldosterone)
 - Secretion K⁺
- Intercalated cells acid-base
- Intercalated cells type A
 - Secretion H⁺
 - Reabsorption HCO₃⁻
- Intercalated cells type B
 - Opposite function to type A
 - Reabsorption H⁺
 - Secretion HCO₃




Intercalated cell

Type B

- HCO-3

Distal tubule – second half & cortical collecting tubules

- Impermeable to urea
- Water reabsorption
 - ADH (vasopressin)



Medullary collecting duct

- <5% water and Na⁺ reabsorbed here
- Final site urine processing
- Water permeability controlled by ADH
- Permeable to urea (some reabsorption)




Regulation of tubular reabsorption

- Glomerulotubular balance
 - Proximal tubules increase their absorption rate when flow through the tubules is increased
 - ↑ GFR → proportional ↑ proximal tubular reabsorption
 - Keeps % reabsorption constant
 - Also occurs in other segments (mainly loop of Henle)
 - Prevents overloading of distal tubule
 - Works alongside other autoregulatory mechanisms (tubuloglomerular feedback)
 - Independent of hormonal control

Regulation of tubular reabsorption

- Fluid physical forces
 - Hydrostatic and colloid osmotic pressures in the tubules, interstitial fluid and peritubular capillaries



Calcium reabsorption

- Proximal tubule
 - 60-70% filtered Ca²⁺ reabsorbed here
 - Mostly passive, paracellular, along electrical and chemical gradients
- Thick ascending loop of Henle
 - 20% filtered Ca²⁺ reabsorbed here also passive, paracellular diffusion
- Distal tubule
 - 10-20% filtered Ca²⁺ reabsorbed here
 - Transcellular
- Reabsorption in thick ascending loop of Henle and distal tubule under control of parathyroid hormone

Phosphate reabsorption

- Majority reabsorbed in proximal tubule
- Influenced by PTH

Wastes

- Creatinine
 - All filtered, none reabsorbed
- Urea
 - Some reabsorbed
- Ammonia
 - Produced in kidney

Measuring the kidney's handling of solutes: fractional excretion

- Plasma vs. urine concentration of solutes, measured in comparison to a substance that is filtered but not reabsorbed (usually creatinine).
- Tells us about whole body concentration of solutes (mainly electrolytes) based on how much the kidney is reabsorbing or secreting them
- Eg. for FEx of sodium:

 $FEx = [(urinary\ Na^+/plasma\ Na^+)\ x\ (plasma\ creatinine/urine\ creatinine)]\ x\ 100$

Normal values – fractional excretion

Substance	% filtered load reabsorbed	
Glucose	100	
Bicarbonate	>99.9	
Na⁺	99.4	
K ⁺	87.8	
Cl ⁻	99.1	
Urea	50	
Creatinine	0	

Normal values – fractional excretion

Substance	% filtered load reabsorbed	Fractional excretion (%)
Glucose	100	0
Bicarbonate	>99.9	<0.1
Na ⁺	99.4	<1
K ⁺	87.8	12.2
Cl ⁻	99.1	0.9
Urea	50	50
Creatinine	0	100

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING

This material has been reproduced and communicated to you by or on behalf of the University of Melbourne pursuant to Part VB of the *Copyright Act 1968 (the Act)*.

The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.