

Machine Learning

Linear Algebra review (optional)

Matrices and vectors

Matrix: Rectangular array of numbers:



4 x 2 matrix

$\mathbb{R}^{4\times 2}$



2 x 3 matrix

$\mathbb{R}^{2\times 3}$

Dimension of matrix: number of rows x number of columns

Matrix Elements (entries of matrix)

$$A = \begin{bmatrix} 1402 & 191 \\ 1371 & 821 \\ 949 & 1437 \\ 147 & 1448 \end{bmatrix}$$

A_{ij} = "i, j entry" in the i^{th} row, j^{th} column.

$A_{11} = 1402$

$A_{12} = 191$

$A_{32} = 1437$

$A_{41} = 147$

A_{43} = Undefined (error)

Vector: An n x 1 matrix.

$$y = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$$

$n = 4$

← 4-dimensional vector. $\mathbb{R}^4$

$\mathbb{R}^{3\times 2}$

$y_i = i^{th}$ element

$y_1 = 460$

$y_2 = 232$

$y_3 = 315$

A, B, C, X

a, b, x, y

1-indexed vs 0-indexed:

$$y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}$$

1-indexed

y[1]

$$y = \begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

0-indexed

y[0]



Machine Learning

Linear Algebra review (optional)

Addition and scalar multiplication

Matrix Addition





Scalar Multiplication

real number

$$3 \times \begin{bmatrix} 1 & 0 \\ 2 & 5 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 6 & 15 \\ 9 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 5 \\ 3 & 1 \end{bmatrix} \times 3$$

3×2 $\qquad$ 3×2

$$\begin{bmatrix} 4 & 0 \\ 6 & 3 \end{bmatrix} /4 = \frac{1}{4} \begin{bmatrix} 4 & 0 \\ 6 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \frac{3}{2} & \frac{3}{4} \end{bmatrix}$$

Combination of Operands





Machine Learning

Linear Algebra review (optional)

Matrix-vector multiplication

Example


$\begin{bmatrix} 1 & 3 \\ 4 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 16 \\ 4 \\ 7 \end{bmatrix}$
3×2
2×1
3×1 matrix
1×1 + 3×5 = 16
4×1 + 0×5 = 4
2×1 + 1×5 = 7

Details:



m x n matrix (m rows, n columns)

n x 1 matrix (n-dimensional vector)

m-dimensional vector

To get y_i, multiply A's i^{th} row with elements of vector x, and add them up.

Example



$1 \times 1 + 2 \times 3 + 1 \times 2 + 5 \times 1 = 14$

$0 \times 1 + 3 \times 3 + 0 \times 2 + 4 \times 1 = 13$

$-1 \times 1 + (-2) \times 3 + 0 \times 2 + 0 \times 1 = -7$

House sizes:

- 2104
- 1416
- 1534
- 852

$$h_\theta(x) = -40 + 0.25x$$





Machine Learning

Linear Algebra review (optional)

Matrix-matrix multiplication

Example

$$\begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 0 & 1 \\ 5 & 2 \end{bmatrix} = \begin{bmatrix} 11 & 10 \\ 9 & 14 \end{bmatrix}$$

$2 \times 3 \quad 3 \times 2 \quad 2 \times 2$

$$\begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 11 \\ 9 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 3 & 2 \\ 4 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 10 \\ 14 \end{bmatrix}$$

Details:

$A \times B = C$



m x n matrix
(m rows,
n columns)

n x o matrix
(n rows,
o columns)

m x o
matrix

The i^{th} column of the matrix C is obtained by multiplying A with the i^{th} column of B. (for i = 1,2,…,o)

Example

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 9 & 7 \\ 15 & 12 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 0 \\ 3 \end{bmatrix} = \begin{bmatrix} 1\times 0 + 3\times 3 \\ 2\times 0 + 5\times 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 15 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1\times 1 + 3\times 2 \\ 2\times 1 + 5\times 2 \end{bmatrix} = \begin{bmatrix} 7 \\ 12 \end{bmatrix}$$

House sizes:

$$\begin{Bmatrix} 2104 \\ 1416 \\ 1534 \\ 852 \end{Bmatrix}$$

Have 3 competing hypotheses:

1. $h_\theta(x) = -40 + 0.25x$
2. $h_\theta(x) = 200 + 0.1x$
3. $h_\theta(x) = -150 + 0.4x$

Matrix

$$\begin{bmatrix} 1 & 2104 \\ 1 & 1416 \\ 1 & 1534 \\ 1 & 852 \end{bmatrix}$$

$\times$

Matrix

$$\begin{bmatrix} -40 & 200 & -150 \\ 0.25 & 0.1 & 0.4 \end{bmatrix}$$

$=$

$$\begin{bmatrix} 486 & 410 & 692 \\ 314 & 342 & 416 \\ 344 & 353 & 464 \\ 173 & 285 & 191 \end{bmatrix}$$

Prediction of first ho

Predictions of 2nd ho



Machine Learning

Linear Algebra review (optional)

Matrix multiplication properties

$3 \times 5 = 5 \times 3$ "Commutative"

Let A and B be matrices. Then in general,

$A \times B \neq B \times A$. (not commutative.)

E.g.

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}$$

$\neq$

$$\begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 2 & 2 \end{bmatrix}$$

$A \times B$

$m \times n$ $\quad$ $n \times m$

$A \times B$ is $m \times m$

$B \times A$ is $n \times n$

$3 \times 5 \times 2$

$3 \times 10 = 30 = 15 \times 2$

$3 \times (5 \times 2) = (3 \times 5) \times 2$

"Associative"

$A \times (B \times C)$

$(A \times B) \times C$

$A \times B \times C.$

Let $D = B \times C.$ Compute $A \times D.$

Let $E = A \times B.$ Compute $E \times C.$

$A \times (B \times C)$

$(A \times B) \times C$

Same answer.

Identity Matrix

1 is identity

$1 \times z = z \times 1 = z$ for any z

Denoted I (or $I_{n \times n}$).

Examples of identity matrices:

$[1]$ 1 x 1

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$ 2 x 2

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$ 3 x 3

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$ 4 x 4

Informally:



For any matrix A,

$$A \cdot I = I \cdot A = A$$

$I_{n \times n}$

A: m x n, I: n x n, I: m x m, A: m x n, result: m x n

Note:

$AB \neq BA$ in general

$AI = IA$ ✓



Machine Learning

Linear Algebra review (optional)

Inverse and transpose

1 = "identity"

$3 \times (3^{-1}) = 1$, $3^{-1} = \frac{1}{3}$

$12 \times (12^{-1}) = 1$, $12^{-1} = \frac{1}{12}$

$0 \times (0^{-1})$ undefined

Not all numbers have an inverse.

Matrix inverse:

If A is an m x m matrix, and if it has an inverse,

(square matrix (# rows = # columns))

A^{-1}

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$A(A^{-1}) = A^{-1}A = I.$$

E.g.

$$\begin{bmatrix} 3 & 4 \\ 2 & 16 \end{bmatrix} \begin{bmatrix} 0.4 & -0.1 \\ -0.05 & 0.075 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_{2\times 2}$$

2x2

A $\quad$ A^{-1} $\quad$ $A^{-1}A$

Matrices that don't have an inverse are "singular" or "degenerate"

Matrix Transpose

Example: $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & 5 & 9 \end{bmatrix}$ $B = A^T = \begin{bmatrix} 1 & 3 \\ 2 & 5 \\ 0 & 9 \end{bmatrix}$

Let A be an m x n matrix, and let $B = A^T$.

Then B is an n x m matrix, and

$$B_{ij} = A_{ji}.$$

$B_{12} = A_{21} = 2$

$B_{32} = 9$ $A_{23} = 9$.