

Assignment 2

Due date: Thursday, 3 October 2024, 23:59

Exercise 2.3, Simplifying a Formula (★)

(8 Points)

Consider the propositional formula

$$F = ((B \vee C) \rightarrow ((A \vee \neg B) \wedge C)) \vee (A \wedge \neg C)$$

Give a formula G that is equivalent to F , but in which each atomic formula A , B , and C appears at most once. Prove that $F \equiv G$ by providing a sequence of equivalence transformations with *at most* 12 steps.

Expectation. Your proof should be in the form of a sequence of steps, where each step consists of applying the definition of \rightarrow (that is $F \rightarrow G \equiv \neg F \vee G$), one of the rules given in Lemma 2.1 of the lecture notes ¹, or one of the following rules: $F \wedge \neg F \equiv \perp$, $F \wedge \perp \equiv \perp$, $F \vee \perp \equiv F$, $F \vee \neg F \equiv \top$, $F \wedge \top \equiv F$, and $F \vee \top \equiv \top$. For this exercise, associativity is to be applied as in Lemma 2.1.3. Each step of your proof should apply a *single* rule *once* and state *which* rule was applied.

The following 12 steps show the equivalence between F and G , where $G := \neg B \vee A$.

$F = ((B \vee C) \rightarrow ((A \vee \neg B) \wedge C)) \vee (A \wedge \neg C)$	
$F = (\neg(B \vee C) \vee ((A \vee \neg B) \wedge C)) \vee (A \wedge \neg C)$	def of \rightarrow
$F = (\neg B \wedge \neg C \vee ((A \vee \neg B) \wedge C)) \vee (A \wedge \neg C)$	def De Morgan
$F = (\neg B \wedge \neg C \vee (C \wedge (A \vee \neg B))) \vee (A \wedge \neg C)$	def commutative
$F = (\neg B \wedge \neg C \vee (C \wedge A) \vee (C \wedge \neg B)) \vee (A \wedge \neg C)$	def 1st distributive law
$F = (\neg B \wedge \neg C \vee (C \wedge \neg B) \vee (C \wedge A)) \vee (A \wedge \neg C)$	def commutative
$F = (\neg B \wedge (C \vee \neg C)) \vee (C \wedge A) \vee (A \wedge \neg C)$	def 1st distributive law
$F = (\neg B \wedge \top) \vee (C \wedge A) \vee (A \wedge \neg C)$	$F \vee \neg F \equiv \top$
$F = (\neg B \vee (C \wedge A)) \vee (A \wedge \neg C)$	$F \wedge \top \equiv F$
$F = \neg B \vee ((C \wedge A) \vee (A \wedge \neg C))$	def associativity
$F = \neg B \vee (A \wedge (C \vee \neg C))$	def 1st distributive law
$F = \neg B \vee (A \wedge \top)$	$F \vee \neg F \equiv \top$
$F = \neg B \vee A$	$F \wedge \top \equiv F$

¹Lemma 2.1 states rules involving propositional symbols, but you may apply those rules at the level of formulas (see Section 2.3.5 of the lecture notes).

Finally, for the formula G defined as $G := \neg B \vee A$, we have shown that $F \equiv G$ by applying a 12-step sequence of equivalence transformations.