

## International application of the Incidence Rate/Window Period model

Estimating risks of transfusion-transmitted infections (TTIs) is essential for monitoring blood safety and for helping physicians and patients make informed decisions when discussing allogeneic transfusion versus other therapeutic options. Since the 1980s and the HIV pandemic, proposed interventions to decrease TTI risks have mostly centered on reducing viral-transmission risks, with consequent implementation of increasingly stringent donor behavioral and highly sensitive laboratory screening procedures in many countries. As these measures have driven risks down, direct methods to estimate risks, such as prospective studies of transfusion recipients or testing of donor samples with sensitive research tests, have become impractical and/or prohibitively expensive due to the very large sample size required to conduct such studies.<sup>1</sup> This has led to the use of mathematical approaches that can be applied to a limited data set and allow for accurate estimation of risks if based on reasonable assumptions.

Prevalence and incidence rate (IR) have commonly been used to measure TTI risks in the blood supply. Prevalence measures the percentage of donations that test positive on a screening and confirmatory test (including old and new infections), whereas incidence measures the rate at which new infections develop in a population at risk. Since donations that have positive screening tests are discarded, the greatest threat to the safety of the blood supply is the donation of blood in the infectious window period (WP), the time between infection and detectability by screening tests. Additional sources of transfusion risk attributable to variant viral strains, immunosilent carriers, and testing error<sup>2-7</sup> are thought to only minimally contribute to overall risk, especially when dual testing systems exist (i.e., serology and NAT).

### RESIDUAL RISK ESTIMATION: THE IR/WP MODEL

The IR/WP model was developed in parallel by investigators from the National Heart, Lung, and Blood Insti-

tute-sponsored Retrovirus Epidemiology Donor Study (REDS),<sup>8,9</sup> the CDC,<sup>10</sup> and the American National Red Cross<sup>10</sup> and has been used in the US to estimate the probability of a WP donation for each viral TTI since the early 1990s. The model estimates the probability of having a potentially infectious donation being released in the blood supply, or residual risk, by multiplying the IR by the number of days during which an infection may be present but not detectable by current screening assays (i.e., the length of the WP). This modeling approach can be implemented without the need to fund special research projects and can be used to generate risk estimates for multiple agents using the same data set. Although the approach requires the presence of an appropriate infrastructure to compile and analyze blood donor data from a proportion of the donor population in any given country, this model has now been applied successfully worldwide to provide country-specific viral TTI estimates, as exemplified by several reports<sup>11-14</sup> in this issue of *Transfusion*.

The residual risk estimates obtained for Australia<sup>15</sup> and countries represented in this issue of *Transfusion*,<sup>11-14</sup> albeit showing some variation, are uniformly low and suggest that the donor selection process (self-selection, behavioral screening, and test screening) adopted by these countries is very effective (Tables 1-3). More importantly, these reports clearly demonstrate that HIV and HCV residual risk estimates are or will be further reduced to extremely low levels after mini-pool NAT implementation. For HIV, the estimated probability of having a potentially infectious donation being released in the blood supply varied from about 1 in 450,000 (Italy) to 1 in 3 million (Australia) donations before NAT, whereas residual risk estimates after NAT varied between 1 in 900,000 (Italy) and 1 in 5 million (Australia) (Table 1). Even more marked was the reduction for HCV, with risk estimates decreasing from about 1 in 120,000 (Australia) to 1 in 800,000 (France) donations before NAT (EIA 3.0) to approximately 1 in 1 million (Australia, Italy) to 1 in 10 million (France) after NAT (Table 2). Although HBV risk estimates need to be evaluated with some caution (see below), data from these reports suggest that of the major viral TTIs evaluated, HBV remains most likely to be transmissible by transfusion, with residual risk estimates of 1 in 75,000 (Spain) to 1 in 520,000 (Australia) donations (Table 3). Mini-pool HBV NAT testing only reduces the WP by about 10 days<sup>20</sup> because the ramp-up viremia

This work was supported by NHLBI contracts N01-HB-97077 (superseded by N01-HB-47114), -97078, -97079, -97080, -97081, and -97082.

TRANSFUSION 2002;42:966-972.

**TABLE 1. HIV risks in RPT donors and NAT yield by country**

Country	Period of observation	Allogeneic collections		IR/10 <sup>5</sup> person years (95% CI)	WP (days)		Residual risk/10 <sup>6</sup> RPT donations		NAT yield/10 <sup>6</sup> donations	
		Donations ×10 <sup>6</sup> /year (n)	Study coverage (%)		Sero	NAT	Serology estimate	Serology/NAT estimate	Predicted RPT*/all†	Observed
Australia <sup>15</sup>	7/00-6/01	1.0	100	0.6 (0.1-2.0)	22	11	0.3	0.2	0.2*/0.2†	0.0‡ p§ = 1.00
Europe										
EPFA <sup>16,  </sup>										
Central <sup>17</sup>	1/97-NA	NA	NA	0.7 (0.5-1.1)	22	11	0.4	0.2	0.2*/0.3†	0.6‡ p§ = 0.50
France <sup>12</sup>	1/98-12/00	3.0	60	1.2 (0.7-2.0)	22	11	0.7	0.4	0.4*/0.4†	0.6‡ p§ = 1.00
Italy <sup>13</sup>	1/96-12/00	2.4	20	3.8 (2.8-5.0)	22	11	2.3	1.1	1.1*	
Spain <sup>11</sup>	1/97-12/99	1.4	71	3.2 (2.2-4.5)	22	11	1.9	1.0	1.0*	
US	1/91-12/93 <sup>8</sup>	13.2 <sup>18</sup>	8	3.4 (2.2-4.8)	22	11	2.0	1.0	1.0*	
	1/96-12/00††	13.2 <sup>18</sup>	8	1.6 (1.0-2.3)	22	11	1.0	0.5	0.5*	
	1/00-12/01 <sup>14</sup>	13.2 <sup>18</sup>	50	1.6 (1.2-2.0)	16**	11	0.7	0.5	0.2*	0.2†† p§ = 1.00

\* Predicted yield for RPT donations obtained by multiplying RPT IRs (5th column) by the detectable RNA+ window (11 days).

† Predicted yield for all donations obtained by multiplying the estimated IR in all donors by the detectable RNA+ window (11 days); we assumed that 10 percent (Australia) or 20 percent (EPFA and France) of donations were FT.

‡ Observed yield per 10<sup>6</sup> donations (FT and RPT). There was one HIV yield case per 1.8 10<sup>6</sup> donations in France (written communication, Josiane Pilonel, June 2002).

§ Two-tailed two-sample exact Poisson test comparing predicted and observed yield donations.

|| Representation from Denmark, England and Wales, Finland, Germany, and Switzerland. IR for EPFA estimated from the residual risk obtained by the Müller-Breitkreutz mathematical model and used as an estimate for Germany and Austria to allow for comparison of predicted versus observed NAT yield in Central Europe.

¶ REDS data.

\*\* Accounts for HIV-1 p24Ag testing.

†† Observed yield per 10<sup>6</sup> RPT donations.

**TABLE 2. HCV risks in RPT donors and NAT yield by country**

Country	Period of observation	Allogeneic collections		IR/10 <sup>5</sup> person years (95% CI)	WP (days)		Residual risk/10 <sup>6</sup> RPT donations		NAT yield/10 <sup>6</sup> donations	
		Donations ×10 <sup>6</sup> /year (n)	Study coverage (%)		Sero	NAT	Serology estimate	Serology/NAT estimate	Predicted RPT*/all†	Observed
Australia <sup>15</sup>	7/00-6/01	1.0	100	4.7 (2.7-7.5)	66	7‡	8.5	0.9	7.6*/8.4†	1.0§ p   = 0.02
Europe										
Central <sup>17</sup>	1/97-NA	NA	NA	2.3 (0.3-4.7) <sup>19,¶</sup>	66	7‡	4.2	0.4	3.8*/4.5†	1.7§ p   = 0.13
EPFA <sup>16,**</sup>	1/97-12/97	NA	NA	0.9 (0.7-1.2)	66	7‡	1.6	0.2	1.4*	
France <sup>12</sup>	1/98-12/00	3.0	60	0.6 (0.3-1.3)	66	7	1.2	0.1	1.0*/1.2†	0.6§ p   = 0.76
Italy <sup>13</sup>	1/96-12/00	2.4	20	4.1 (3.0-5.3)	70	12	7.9	1.3	6.6*	
Spain <sup>11</sup>	1/97-12/99	1.4	71	3.7 (2.6-5.1)	66	7‡	6.7	0.7	6.0*	
US	1/91-12/93 <sup>8</sup>	13.2 <sup>18</sup>	8	4.3 (2.4-6.9)	82	10‡	9.7	1.2	NA	
	1/96-12/00††	13.2 <sup>18</sup>	8	2.9 (2.1-3.9)	70	8	5.6	0.6	4.9*	
	1/00-12/01 <sup>14</sup>	13.2 <sup>18</sup>	50	1.9 (1.5-2.4)	70	10	3.6	0.5	3.1*	2.6‡‡ p   = 0.43

\* Predicted yield for RPT donations obtained by multiplying RPT IR (5th column) by the detectable RNA+ window (58-62 days).

† Predicted yield for all donations was calculated by multiplying the IR in all donors by the detectable RNA+ window (58-62 days); 10 percent (Australia) or 20 percent (Central Europe and France) of donations were assumed to be FT.

‡ The current HCV mini-pool NAT WP was assumed to be between 7 and 12 days, based upon the best current estimates for sensitivity of mini-pool HCV NAT. Any estimate in the original publication longer than 12 days was changed to reflect this assumption.

§ Observed yield/10<sup>6</sup> donations (FT, RPT). There was one HCV yield case/1.8 10<sup>6</sup> donations in France (written communication, Josiane Pilonel, June 2002).

|| Two-tailed two-sample exact Poisson test comparing predicted and observed yield donations.

¶ IR based on 1994-95 second and third generation anti-HCV EIA data for Germany and was used as an estimate of the IR in Germany and Austria to allow for comparison of predicted versus observed NAT yield in Central Europe. We assumed that the WP for the HCV serology assay used in Central Europe<sup>17</sup> in 1997 onward was 66 days.

\*\* Representation from Denmark, England and Wales, Finland, Germany, and Switzerland. IR estimated from the residual risk obtained by the Müller-Breitkreutz mathematical model.

†† REDS data.

‡‡ Observed yield/10<sup>6</sup> RPT donations.

**TABLE 3. HBV risk and predicted NAT yield in RPT donors by country**

Country	Period of observation	Allogeneic collections		IR/10 <sup>5</sup> person years (95% CI)	WP (days)		Residual risk/10 <sup>6</sup> RPT donations		NAT yield/10 <sup>6</sup> RPT donations
		Donations ×10 <sup>6</sup> /year (n)	Study coverage (%)		Sero	NAT	Serology estimate	Serology/NAT estimate	Predicted
Australia <sup>15</sup>	7/00-6/01	1.0	100	1.6 (0.3-4.6)	45	34	1.9	1.5	0.5
Europe									
France <sup>12</sup>	1/98-12/00	3.0	60	1.4 (0.9-2.2)	56	46	2.1	1.8	0.4
Spain <sup>11</sup>	1/97-12/99	1.4	71	8.4 (5.2-12.6)	59	34	13.5	7.8	5.7
US	1/91-12/93 <sup>8</sup>	13.2 <sup>18</sup>	8	9.8 (6.7-13.4)	59	34	15.8	9.1	6.7
	1/96-12/00†	13.2 <sup>18</sup>	8	6.0 (4.0-8.7)	59	49	9.7	8.1	1.6
	1/00-12/01 <sup>14</sup>	13.2 <sup>18</sup>	50	3.0 (2.3-3.9)	59	49	4.9	4.0	0.8

\* HBV IRs in RPT donors (adjusted from HBsAg) are presented.

† Data from REDS.

phase for HBV is characterized by a much slower doubling time (2.6 days) than for HIV (20.5 hours)<sup>21</sup> or HCV (14.9 hours).<sup>22</sup> The difficulty in decreasing HBV risks also stems from the fact that no risk factor can be identified in 30 to 40 percent of HBV cases<sup>23</sup> and the consequent inability to develop donor-screening criteria that are sensitive enough to effectively screen out HBV-infected donors and specific enough to avoid unnecessary deferral of healthy donors. Widespread implementation of HBV immunization programs should, in the future, help reduce risks of HBV transfusion transmission by decreasing the number of infectious donors and susceptible transfusion recipients.

### ASSUMPTIONS OF THE IR/WP MODEL

To better interpret the data at hand and be able to compare IRs (and consequently residual risk estimates) obtained by various countries, the reader should become familiar with the IR/WP model's basic assumptions, key components, and major nuances in method. The basic assumptions of the model are that 1) WP donations represent the most dominant source of risk; 2) overall IR in repeat (RPT) donors is constant throughout the period of the study; 3) a recently infected donor is as likely to donate in the early interval after infection as he or she is to donate subsequently; and 4) if no further adjustment is made, first-time (FT) donors have the same IR as RPT donors. The first assumption appears to be robust for HIV and HCV with WP risk probably accounting for 90 percent or more of the total risk and other sources of risk (viral variants, chronic seronegative carriers, and testing error) for 10 percent or less.<sup>7</sup> This is also the case for HBV in countries performing anti-HBc screening, whereas HbsAg, anti-HBc-reactive chronic carriers likely contribute a level of risk equal or greater to WP risk in the absence of anti-HBc screening.<sup>24,25</sup> We have no reason to doubt the validity of the second assumption if the period of the study is short (or, for longer periods, if there does

not appear to be changes in rates through time that can result from changes in the epidemiology of the agent in the population or revision of screening criteria).

The third assumption may be more problematic because more recent studies have noted that the seroconversion interval may be longer than expected for HIV<sup>26</sup> or HCV incident cases.<sup>15,16</sup> This would imply that some donors delay their return around the time of infection and suggests that the model (which assumes a steady-state rate of donation) may tend to overestimate the yield of tests that detect infection earlier (i.e., NAT or HIV p24Ag). Consistent with this concern, the IR/WP model predicted that eight HCV NAT-only donations would be detected in Australia (of 0.97 million donations screened) compared with the one observed ( $p = 0.02$ , Table 2), and Seed et al.<sup>15</sup> reported that the duration of the seroconversion interval for HCV seroconverters was much longer than for the other viruses. Similarly, an apparent difference in observed versus expected HCV yield was noted for Central Europe, although these results were not significant ( $p = 0.13$ ) and need to be viewed with caution because regional incidence data (Germany, 1994-95)<sup>19</sup> may not accurately estimate NAT yield data that would be obtained throughout Central Europe after 1996. There was no significant difference in predicted versus observed HCV yield data in France or in the US. This is compatible with the finding of no delay in return for HCV seroconverters in the US.<sup>26</sup> Further, no significant differences were found in any study between observed (0.0-0.6/10<sup>6</sup> donations) and predicted (0.2-0.4/10<sup>6</sup> donations) yields for HIV (Table 1). (Although NAT yield data are now becoming available for several other countries,<sup>27</sup> we were not able to evaluate whether observed and expected yields differed for countries for which we could not find published IR data.) Thus, the IR/WP model seems to predict reasonably well the yield of new assays unless donors significantly delay their return around the time of infection and seroconversion.

Finally, the recent development of a sensitive/less-sensitive HIV-1 assay<sup>28</sup> and evaluation of NAT-yield

data<sup>14</sup> now permit assessment of IRs in FT donors. Jansen et al.<sup>28</sup> reported a 2.4-fold higher IR in FT donors (7.18/100,000 person-years) than in RPT donors (2.95/100,000 person-years) using the sensitive/less-sensitive HIV-testing strategy. Using NAT-yield data, Dodd et al.<sup>14</sup> confirm this 2-fold differential in IRs between FT and RPT donors for HIV ( $2.1\times$ ) and extend this finding to HCV ( $2.4\times$ ). It is therefore now possible to estimate IRs in all donation if the relative proportion of FT and RPT donors is known. Most developed countries report that their donation base consists of about 20 percent FT and 80 percent RPT donors, which implies that the overall donor IR can be approximated by multiplying the IR obtained in RPT donors by 1.2 ( $[0.2 \times 2 \times \text{IR in RPT donors}] + [0.8 \times \text{IR in RPT donors}]$ ).

### ESTIMATING THE IR

The IR/WP model can only be applied if two key statistics, the IR and the WP, are known with accuracy. Let us first turn to IR estimation because this statistic armed with a 95 percent CI allows for comparison of risks between countries that rely on similar testing systems and, therefore, similar WPs (95% CI around residual risks are usually unavailable because data is often too limited to allow calculation of variability around the WP estimate). IRs are derived by dividing the numbers of known confirmed incident cases (detected by serologic or NAT screening) by the number of person-years or the sum of the periods during which donors are at risk (i.e., are not infected yet). This derivation is confined to RPT donors because this population is followed for a certain period, allowing us to evaluate how many donors are incident cases (i.e., first test negative on all assays and then test positive for a particular marker on a subsequent donation). The minimum statistics required for IR calculations are the number of incident cases (the numerator), the total number of RPT donations, and the mean interdonation interval length (multiplication of the last two entities provides the number of person-years or denominator). Alternatively, if donation histories are available for all RPT donors, we can derive person-years by summing up the length of all interdonation intervals. Because a number of adjustments can be made to both the number of incident cases and to the number of person-years, the reader should assess the following factors to determine if studies are directly comparable:

1. Was there an adjustment for false-positive confirmatory tests? Some studies (e.g., Alvarez et al.,<sup>11</sup> REDS), upon further review of their laboratory results, have excluded false-positive donations, thereby providing a more accurate count of true sero- or NAT-incident cases.
2. Was the full length of the seroconversion intervals included in the person-years calculation or just a

portion (often half) to try to adjust for when the infection probably occurred during the interval? In our experience, this adjustment does not have any demonstrable impact on the magnitude of the IR estimate.

3. Was there an adjustment for nontransfused seronegative and seroconversion units? Studies (e.g., Pillonel et al.,<sup>12</sup> Velati et al.,<sup>13</sup> REDS) may choose to exclude interdonation intervals that start with a donation that would not have been transfused because it was reactive for another marker or, when applicable, because the donor used the Confidential Unit Exclusion process. This may lead to a slight decrease in the number of incident cases and person-years.
4. Were all interdonation intervals that start with a donation tested by serology alone and end by a donation tested by both serology and NAT (or p24Ag for HIV) lengthened by the reduction in the WP? The IRs may be overestimated if this adjustment is not made, although this adjustment will only have a significant impact when the reduction in WP is substantial. In Tables 1 through 3, this adjustment was only applied to REDS IRs calculated for the period 1996-2000 because the US implemented NAT for HIV and HCV in 1999.
5. Was the measured HBsAg IR taken as the HBV incidence, or was an adjustment factor calculated to account for the transient nature of HBs-antigenemia? All IRs reported in Table 3 refer to HBV, with most having been adjusted from HBsAg using the method developed by Korelitz et al.<sup>29</sup> More recently, REDS investigators have used a different mathematical derivation to estimate HBV incidence<sup>30,31</sup> that results in similar, albeit slightly higher, estimates.

With these caveats in mind, we can first note that the IRs in RPT donors are small (Tables 1-3). When compared with rates in US blood donors, rates in the US general population are about 9 times higher for HIV (15.0 vs. 1.6), 7 times higher for HCV (13.4 vs. 1.9), and nearly 40 times higher for HBV (114.4 vs. 3.0).<sup>32</sup> Second, some variation exists between countries. Southern Europe (Italy, Spain) had higher HIV IRs than other European countries, the US, or Australia (Table 1). IRs for HCV were highest in Australia (in marked contrast to HIV), followed by Southern Europe (Italy, Spain) and the US (Table 2). Lowest HCV estimates were obtained for the European Plasma Fractionation Association (EPFA)-European countries and France (Table 2). For HBV, rates appeared similar in Australia and France, intermediate in the US, and highest in Southern Europe (Table 3). In the US, the HBV rate obtained at REDS centers was higher than reported by Dodd et al.<sup>14</sup> and probably reflects REDS geographic representation. There appears to have been a decrease in IRs in RPT donors over the last decade in some countries, as noted by Pillonel et al.<sup>12</sup> in France (all markers) and by

Dodd et al.<sup>14</sup> (HBV only) in the US. REDS has also noted that IRs for HIV and HBV have decreased in the US between the early and the late 1990s.

### ESTIMATING THE INFECTIOUS WP

The second key statistic needed to estimate residual risk is the length of the WP for each test. WP estimates for HCV and HBV used in the model are generally based upon analyses of data on the time of exposure by transfusion to the development of a positive test. In contrast, the WP estimate for HIV is a theoretical number derived from mathematical modeling of TTIs occurring from HIV-seronegative units donated by persons who subsequently seroconverted.<sup>1,7</sup> WP estimates for HIV and HCV have been relatively well characterized;<sup>7</sup> however, the WP for HBV is more uncertain because the mean of 59 days is based upon a series of only seven evaluative transfusion-transmitted cases.<sup>33</sup> Although the model uses these WP estimates, what we are really interested in is the period during which a donation is infectious; this infectivity period may or may not be the same as the length of the WP commonly used to estimate residual risks (Tables 1-3). For example, it is unclear whether a donation given in the early stages after exposure (i.e., during the so-called eclipse [RNA-] pre-ramp-up viremia [intermittent RNA+] and very early ramp-up viremia [low viral load] phases) contains enough virus to be infectious.<sup>7</sup> In particular, the significance and infectivity of the low-level intermittent blips of RNA or DNA recently detected in plasma donor seroconversion panels in the pre-ramp-up viremia phase are uncertain and have not been included in the model for residual risk.<sup>22,34</sup> Further studies including additional animal transmission model systems,<sup>35</sup> donor lookback studies as described by Roth et al.<sup>17,25</sup> (where prior units from sero- or NAT-positive RPT donors are traced and recipients of these units tested), and recipient traceback studies<sup>36</sup> (investigation by recall or testing of stored donation specimens of donors whose blood products were transfused to recipients who subsequently acquired an infection after transfusion) are clearly needed to answer these questions and permit better estimation of risks. For example, Weusten et al.<sup>37</sup> used minimum chimpanzee infectious doses data to statistically predict human infectivity, relative to viral load and NAT assay performance, in a recently published mathematical modeling of WP risk. Although this is a clear conceptual advance and seems like a reasonable approach, the animal infectivity data employed in the model were limited, and it is unclear whether the animal data is directly transferable to humans. Further work in this area is called for.

### TTIs: PREDICTED VERSUS OBSERVED

It is difficult to assess whether the residual risk estimates that have been reported here are overestimated as a re-

sult of this uncertainty in WP estimates. In the US, the model would predict that about 7 potentially infectious HIV and HCV donations and 65 potentially infectious HBV donations may be released yearly (of approx. 13.2 million donations in the US per year<sup>18</sup>). Even these rates seem high relative to the paucity of TTI reports. Indeed, only a handful of HIV<sup>38</sup> and HCV<sup>39</sup> transmissions have been documented since NAT-screening was introduced in the late 1990s. For HBV, reports of post-transfusion infections are rare, with the exception of in regions endemic for HBV infection (e.g., Japan).<sup>36</sup> However, the rarity of confirmed transfusion cases should not be interpreted to discount these risk projections for the following reasons: 1) some available allogeneic units are not transfused (7.5% in the US);<sup>18</sup> 2) most countries do not have mandatory surveillance of recipients for TTI, and clinical case reporting would depend upon accurate diagnosis, which may be difficult if symptoms never develop or occur after many years, by which time the association with the original transfusion is difficult to substantiate (e.g., HBV is asymptomatic in 50-70% of adults and spontaneously resolves with no long-term consequences in the majority of cases);<sup>40,41</sup> 3) 24 to 34 percent of RBC recipients may not survive more than 1 year after their transfusion, and 36 to 49 percent may not survive more than 40 months after transfusion, thereby preventing diagnosis of transfusion transmission if it occurred;<sup>42</sup> 4) some recipients may already carry the infection (recent REDS data indicate that 0.25, 1.75, and 8.5% of recipients are positive for HIV, HCV, and anti-HBc, respectively, before surgery and transfusion); and 5) some recipients may not be susceptible to acquiring the infection if, for example, they have been immunized for HBV.

### ESTIMATING RESIDUAL RISKS IN RESOURCE-RESTRICTED COUNTRIES

The estimated residual risks of viral TTIs, although probably conservative in magnitude, are now very low in developed countries. Further incremental reduction in major viral risk (e.g., from implementation of individual donation NAT or pathogen reduction) is predicted to achieve exceedingly small safety benefits at high cost.<sup>7,43,44</sup> These countries can now afford to turn their attention to reducing other risks of transfusion, such as those caused by bacterial contamination or transfusion errors. However, this is not the situation in most developing countries faced with high TTI prevalence and IRs, and lacking the resources or the infrastructure to implement appropriate serologic screening, let alone NAT screening. The WHO estimates that at least 13 million of the 75 million units collected in the world each year are not completely tested using basic serologic assays, including up to 45 percent of blood donations in developing countries.<sup>45,46</sup> Even when testing is performed, the

sensitivity of assays may be poor and inadvertent release of test-positive units may occur.<sup>46,47</sup> Measuring risks of TTIs in these settings also becomes extremely difficult. As discussed, application of the IR/WP model requires collection of accurate information on the number of incident cases, RPT donations, and average interdonation interval for RPT donors (or seroconversion interval length for all incident cases).<sup>16</sup> This level of testing and data management may not be achievable in some developing countries. It may actually be easier for some countries to test a sample of HIV+ donations by the sensitive/less-sensitive HIV-1 assay or to perform mini-pool NAT for HCV to estimate risks in their donor population than to record donation histories or FT and RPT status. Hence, alternative approaches to modeling risk need to be developed to assist developing countries in monitoring the safety of their blood supply.

**Simone A. Glynn**

*E-mail: Simoneglynn@westat.com*

**Steven H. Kleinman**

**David J. Wright**

*Westat, Rockville, MD*

**Michael P. Busch**

*Blood Centers of the Pacific and*

*University of California, San Francisco, CA*

**for the NHLBI Retrovirus Epidemiology Donor Study**

## REFERENCES

1. Kleinman SH, Busch MP. Risks of transfusion transmitted infection: direct estimation and mathematical modeling. In: Contreras M., ed. New aspects of blood transfusion. Bailliere's best practice and research in clinical haematology. London: Bailliere-Tindall, 2000;13:631-49.
2. Centers for Disease Control and Prevention. Persistent lack of detectable HIV-1 antibody in a person with HIV infection - Utah, 1995. MMWR 1996;45:181-5.
3. Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. The Sentinel Counties Chronic non-A, non-B Hepatitis Study Team. New Eng J Med 1992;327:1899-905.
4. Loussert-Ajaka I, Ly TD, Chaix ML, et al. HIV-1/HIV-2 seronegativity in HIV-1 subtype O infected patients. Lancet 1994;343:1393-4.
5. Jongerius JM, Wester M, Cuypers HTM, et al. New hepatitis B virus mutant form in a blood donor that is undetectable in several hepatitis B surface antigen screening assays. Transfusion 1998;38:56-9.
6. Busch MP, Watanabe KK, Smith JW et al. False-negative testing errors in routine viral marker screening of blood donors. For the Retrovirus Epidemiology Donor Study. Transfusion 2000;40:585-9.
7. Busch MP. Closing the windows on viral transmission by blood transfusion. In: Stramer SL, ed. Blood safety in the new millenium. Bethesda, MD: American Association of Blood Banks, 2001:45.
8. Schreiber GB, Busch MP, Kleinman SH, Korelitz JJ. The risk of transfusion-transmitted viral infections. The Retrovirus Epidemiology Donor Study. N Engl J Med 1996; 334: 1685-90.
9. Kleinman S, Busch MP, Korelitz JJ, Schreiber GB. The incidence/window period model and its use to assess the risk of transfusion-transmitted human immunodeficiency virus and hepatitis C virus infection. Transfus Med Rev 1997;11:155-72.
10. Lackritz EM, Satten GA, Aberle-Grasse J, et al. Estimated risk of transmission of the human immunodeficiency virus by screened blood in the United States. N Engl J Med 1995;333:1721-5.
11. Alvarez M, Oyonarte S, Rodriguez PM, Hernandez JM. Estimated risk of transfusion-transmitted viral infections in Spain. Transfusion 2002;42:994-8.
12. Pillonel J, Laperche S, Saura C, Desenclos JC, Courouce AM. Trends in residual risk of transfusion-transmitted viral infection in France between 1992 and 2000. Transfusion 2002;42:980-8.
13. Velati C, Romano L, Baruffi L, et al. Residual risk of transfusion-transmitted HCV and HIV infections by antibody-screened blood in Italy. Transfusion 2002;42:989-93.
14. Dodd RY, Notari IV EP, Stramer SL. Current prevalence and incidence of infectious disease markers and estimated window period risk in the American Red Cross blood donor population. Transfusion 2002;42:975-9.
15. Seed CR, Cheng A, Ismay SL, et al. Assessing the accuracy of three viral risk models in predicting the outcome of implementing HIV & HCV NAT donor screening in Australia and the implications for future HBV NAT. Transfusion 2002;in press.
16. Müller-Breitkreutz K. Results of viral marker screening of unpaid blood donations and probability of window period donations in 1997. Vox Sang 2000;78:149-57.
17. Roth WK, Weber M, Buhr S, et al. Yield of HCV and HIV-1 NAT after screening of 3.6 million blood donations in Central Europe. Transfusion 2002;42:862-8.
18. Comprehensive report on blood collection and transfusion in the United States in 1999. Bethesda, MD: The National Blood Data Resource Center, 2001.
19. Koerner K, Cardoso M, Dengler Th, Kerowgan M, Kubanek B. Estimated risk of transmission of hepatitis C virus by blood transfusion. Vox Sang 1998;74:213-6.
20. Biswas R, Busch M, Hsia C, Laycock M, Hirschhorn D, Tabor E. Comparative sensitivity of HBV NAT and HBsAg donor testing (abstract). Transfusion 2001;41:8S.
21. Fiebig EW, Heldebrandt C, Smith R, et al. HIV viremia preceding antibody (ab) seroconversion (sc): detection by p24Antigen minipool nucleic acid amplification tests (mp-NAT) and individual donation (ID) NAT (abstract). Transfusion 2000;40:25S.
22. Busch MP, Giachetti C, Conrad A, et al. Dynamics of HCV viremia during early HCV infection: implications for

- minipool (MP) vs individual donation (ID) nucleic acid amplification tests (NAT) (abstract). *Transfusion* 2000;40:25S.
23. Alter MJ, Mast EE. The epidemiology of viral hepatitis in the United States. *Gastroenterol Clin North Am* 1994;23:437-55.
24. Kleinman SH, Busch MP. HBV: amplified and back in the blood safety spotlight. *Transfusion* 2001;41:1081-5.
25. Roth WK, Weber M, Petersen D, et al. Nucleic acid testing (NAT) for hepatitis B virus and anti-HBc testing increase blood safety. *Transfusion* 2002;42:869-75.
26. Schreiber GB, Glynn SA, Satten GA, et al. HIV seroconverting donors delay their return: screening test implications. *Transfusion* 2002;42:414-21.
27. International Forum. Implementation of donor screening for infectious agents transmitted by blood by nucleic acid technology. *Vox Sang* 2002; 82:87-111.
28. Janssen RS, Satten GA, Stramer SL, et al. New testing strategy to detect early HIV-1 infection for use in incidence estimates and for clinical and prevention purposes. *JAMA* 1998;280:42-8.
29. Korelitz JJ, Busch MP, Kleinman SH, et al. A method for estimating hepatitis B virus incidence rates in volunteer blood donors. *Transfusion* 1997;37:634-40.
30. Wright DJ, Glynn SA, Busch M, et al. A proposed new method to estimate hepatitis B incidence rates based on HBsAg marker (abstract). *Transfusion* 1999;39:106S.
31. Glynn SA, Kleinman SH, Schreiber GB, et al. Trends in incidence and prevalence of major transfusion-transmissible viral infections in US blood donors, 1991 to 1996. *JAMA* 2000;284: 229-35.
32. Dodd RY. In: Stramer SL, ed. Blood safety in the new millennium. Bethesda, MD: AABB, 2001;42:97-122.
33. Mimms LT, Mosley JW, Hollinger FB, et al. Effect of concurrent acute infection with hepatitis C virus on acute hepatitis B virus infection. *BMJ* 1993;307:1095-7.
34. Nübling CM, Unger G, Chudy M, Raia S, Löwer J. Sensitivity of HCV core antigen and RNA detection in the early infection phase. *Transfusion* 2002;42:1037-45.
35. Murthy KK, Henrad DR, Eichberg JW, et al. Redefining the HIV-infectious window period in the chimpanzee model: evidence to suggest that viral nucleic acid testing can prevent blood-borne transmission. *Transfusion* 1999;39:688-93.
36. Matsumoto C, Tadokoro K, Fujimura K, et al. Analysis of HBV infection after blood transfusion in Japan through investigation of a comprehensive donor specimen repository. *Transfusion* 2001; 41:878-84.
37. Weusten JJ, van Drimmelen AA, Lelie PN. Mathematical modeling of the risk of HBV, HCV, and HIV transmission by window phase donations not detected by NAT. *Transfusion* 2002;42:537-48.
38. Delwart E, Kalmin N, Jones S, et al. First case of HIV transmission by an RNA screened blood donation. Poster #768-W. Presented at the 9th Conference on Retroviruses & Opportunistic Infections. Seattle WA, February 24-28, 2002. Available at <http://www.retroconference.org>.
39. Schüttler CG, Caspari G, Jursch CA, et al. Hepatitis C virus transmission by a blood donation negative in nucleic acid amplification tests for viral RNA [letter]. *Lancet* 2000; 355:41-2.
40. Moyer LA, Mast EE. Hepatitis B: virology, epidemiology, disease and prevention, and an overview of viral hepatitis. *Am J Prev Med* 1994;10S:45-55.
41. Shapiro CN. Epidemiology of hepatitis B. *Ped Infect Dis J* 1993;12:433-7.
42. Vamvakas EC. Sources and uses of data on long-term survival after blood transfusion. In: Vamvakas EC, ed. Evidence-based practice of transfusion medicine. Bethesda, MD: AABB, 2001:201-19.
43. O'Grady MJ. The impact of regulatory and technological changes on the cost of inpatient blood and blood products: implications for Medicare. Bethesda, MD: The Project HOPE Center for Health Affairs, 2001.
44. Council of Europe Expert Committee in Blood Transfusion Study Group on Pathogen Inactivation of Labile Blood Components. Pathogen inactivation of labile blood products. *Transfus Med* 2001;11:149-75.
45. WHO and International Federation of Red Cross and Red Crescent Societies. Safe blood starts with me. No. 12. Geneva: WHO, 2000.
46. Chamberland ME, Lackritz EM, Busch MP. HIV screening of the blood supply in developed and developing countries. *AIDS Rev* 2001;3:24-35.
47. Moore A, Herrera G, Nyamongo J, et al. Estimated risk of HIV transmission by blood transfusion in Kenya. *Lancet* 2001;358:657-60. ■