

PLANTCOM meeting, Dijon

Quantitative genetics of plant–plant interactions: case of crop mixtures

Jemay Salomon

UMR GQE Le Moulon

Université Paris–Saclay, INRAE, CNRS, AgroParisTech

Outline

Context and motivation

Case of crop mixtures

- Material and methods

- Results

- Conclusions and perspectives

Outline

Context and motivation

Case of crop mixtures

Material and methods

Results

Conclusions and perspectives

Intra-plot diversification: major agroecological lever

- ▶ caveat: not used in breeding

Intra-plot diversification: major agroecological lever

- ▶ caveat: not used in breeding

Challenge: combinatorial explosion

- ▶ strategy (here): quantitative genetics model to predict unobserved mixtures
- ▶ *joint* analysis of monovarietal and mixed stands to *gradually* introduce mixing ability in breeding programs

Intra-plot diversification: major agroecological lever

- ▶ caveat: not used in breeding

Challenge: combinatorial explosion

- ▶ strategy (here): quantitative genetics model to predict unobserved mixtures
- ▶ *joint* analysis of monovarietal and mixed stands to *gradually* introduce mixing ability in breeding programs

Gaps of knowledge:

- ▶ Which experimental designs?
- ▶ Which magnitude of genetic (co)variances?
- ▶ Which genetic architectures?

Outline

Context and motivation

Case of crop mixtures

- Material and methods

- Results

- Conclusions and perspectives

Outline

Context and motivation


Case of crop mixtures

Material and methods

Results

Conclusions and perspectives

Genetic model



Social intra-genotypic value: $SIGV := SBV^{SC} + (DBV \times SBV)^{SC}$

Genetic model

Intercrop of i (species s_1) and j (species s_2):

$$Y_{i(j)} = \mu^{s_1, IC} + \textcolor{red}{DBV}_i + SBV_j^{IC} + (DBV \times SBV)_{ij}^{IC}$$

$$Y_{j(i)} = \mu^{s_2, IC} + DBV_j + SBV_i^{IC} + (DBV \times SBV)_{ji}^{IC}$$

Sole crop of i and sole crop of j :

► $Y_i = \mu^{s_1, SC} + \textcolor{red}{DBV}_i + SIGV_i$

► $Y_j = \mu^{s_2, SC} + DBV_j + SIGV_j$

Simulation and inference

- ▶ 200 wheat genotypes + 2 pea genotypes (testers)

Simulation and inference

- ▶ 200 wheat genotypes + 2 pea genotypes (testers)
- ▶ 3 experimental designs of 400 plots each:
 - Sole_only (complete): sole cropping system (SC)
 - Inter_only (sparse): intercropping system (IC)
 - Sole_inter_50 (sparse): combination of SC and IC

Simulation and inference

- ▶ 200 wheat genotypes + 2 pea genotypes (testers)
- ▶ 3 experimental designs of 400 plots each:
 - Sole_only (complete): sole cropping system (SC)
 - Inter_only (sparse): intercropping system (IC)
 - Sole_inter_50 (sparse): combination of SC and IC
- ▶ Parameter values: Moutier et al. (2022); Haug et al. (2023)

Simulation and inference


- ▶ 200 wheat genotypes + 2 pea genotypes (testers)
- ▶ 3 experimental designs of 400 plots each:
 - Sole_only (complete): sole cropping system (SC)
 - Inter_only (sparse): intercropping system (IC)
 - Sole_inter_50 (sparse): combination of SC and IC
- ▶ Parameter values: Moutier et al. (2022); Haug et al. (2023)
- ▶ Varying proportions of $\text{var}(\text{SIGV})$ relative to $\text{var}(\text{DBV})$

Simulation and inference

- ▶ 200 wheat genotypes + 2 pea genotypes (testers)
- ▶ 3 experimental designs of 400 plots each:
 - Sole_only (complete): sole cropping system (SC)
 - Inter_only (sparse): intercropping system (IC)
 - Sole_inter_50 (sparse): combination of SC and IC
- ▶ Parameter values: Moutier et al. (2022); Haug et al. (2023)
- ▶ Varying proportions of $\text{var}(\text{SIGV})$ relative to $\text{var}(\text{DBV})$
- ▶ Software implementation: `plantmix` (R package)

Panel and field trial

- 200 wheat varieties
 - ▶ 395k SNPs
- 2 pea varieties (testers)
- All sole and intercrops in 2 complete blocks
- Traits per species: grain yield, TKW, final height, SLA, soil cover



Panel and field trial



Le Moulon, Saclay, 2025

Outline

Context and motivation


Case of crop mixtures

Material and methods


Results

Conclusions and perspectives


Accuracy of parameter estimation



Accuracy of parameter estimation




Accuracy of parameter estimation




- Experimental design, genetic effects, and the ratio $\sigma_{SIGV}^2 / \sigma_{DBV}^2$ influence accuracy.

Accuracy of parameter estimation



- Experimental design, genetic effects, and the ratio $\sigma_{SIGV}^2 / \sigma_{DBV}^2$ influence accuracy.
- Inter_only provides the most accurate estimates.

Accuracy of parameter estimation



- **Experimental design, genetic effects, and the ratio $\sigma_{SIGV}^2 / \sigma_{DBV}^2$ influence accuracy.**
- **Inter_only provides the most accurate estimates.**
- **Sole_inter_50 helps decoupling SIGV and DBV, improving their estimation.**

Variance estimates for grain yield

σ^2_{DBV}	σ^2_{SBV}	σ^2_{SIGV}	$\sigma^2_{\text{DBV} \times \text{SBV}}$	$\sigma^2_{\text{e-pur}(w)}$	$\Sigma^2_{\text{e-mix}(w)}$	$\Sigma^2_{\text{e-mix}(p)}$
10.1 [8.01, 11.39]	1.25 [0.17, 2.47]	2.49 [0, 5.23]	2.19 [0.14, 5.76]	16.6 [12.00, 20.00]	12.78 [10.61, 13.98]	18.53 [15.56, 21.30]

Values are shown as the estimate (first line), followed by the 95% confidence interval in brackets (second line), calculated across 500 replicates using bootstrapping.


- Social genetic variances < variance of direct breeding values

Correlation estimates for grain yield

$\rho_{\text{DBV-SBV}}$	$\rho_{\epsilon\text{-mix}}$
0.24	-0.19
[0.01, 1]	[-0.29, -0.07]

- ▶ Unexpected positive correlation observed between DBV and SBV
⇒ favorable wheat–pea interactions

Meta-GWAS on grain yield [DBV, SBV, SIGV]



H_0 rejected when a SNP is associated with
at least two traits (FDR controlled at 5%)

- **9 pleiotropic regions detected**
 - Not detected in single-trait GWAS.
 - Further analyses underway for validation.

Outline

Context and motivation

Case of crop mixtures

Material and methods

Results

Conclusions and perspectives

Take-Home Messages

- **Joint modeling of sole and intercrops + incomplete design + genomic relationship matrix: accurate estimations of breeding values.**

Take-Home Messages

- **Joint modeling of sole and intercroops + incomplete design + genomic relationship matrix: accurate estimations of breeding values.**
- **Field data analyses indicated that social genetic variances were small, but still contribute significantly**

Take-Home Messages

- **Joint modeling of sole and intercrops + incomplete design + genomic relationship matrix: accurate estimations of breeding values.**
- **Field data analyses indicated that social genetic variances were small, but still contribute significantly**
- **Methodology allowing breeders to fine-tune their program depending on the proportion of ressources they want to allocate to sole vs intercrops.**

Take-Home Messages

- **Joint modeling of sole and intercroops + incomplete design + genomic relationship matrix: accurate estimations of breeding values.**
- **Field data analyses indicated that social genetic variances were small, but still contribute significantly**
- **Methodology allowing breeders to fine-tune their program depending on the proportion of ressources they want to allocate to sole vs intercroops.**
- **Preliminary GWAS identified several regions associated with at least two types of breeding values.**


Take-Home Messages

- **Joint modeling of sole and intercroops + incomplete design + genomic relationship matrix: accurate estimations of breeding values.**
- **Field data analyses indicated that social genetic variances were small, but still contribute significantly**
- **Methodology allowing breeders to fine-tune their program depending on the proportion of ressources they want to allocate to sole vs intercroops.**
- **Preliminary GWAS identified several regions associated with at least two types of breeding values.**
- **Analyze the other traits, notably those from drone imaging (PhD V. Freitas).**

Acknowledgments

- ▶ **GQE:** J. Enjalbert, T. Flutre, C. Bourhis-Lézier, V. Freitas
 - ▶ M. Lu, P. Briens, S. Roty, T. Linares, F. Legendre, J. Hélie, J. Le Gall, L. Salze
 - ▶ E. Akaffou, L. Wang, M. Mesnil, D. Wang, F. Petit, R. Diallo, U. Louis, C. Coste, F. Alléhaut
- ▶ **UEVS:** C. Bédard, F. Barriuso, D. Sowamber, J. Cannesson, O. Minguy, A. Belkian
- ▶ **ECOSYS:** J.-M. Gilliot
- ▶ **BIOGER:** T. Vidal, M. Delrieu
- ▶ **IGEPP:** N. Moutier, R. Perronne
- ▶ **AGAP:** A. Baranger

Supp 1: meta-GWAS on grain yield [DBV, SBV, SIGV]



Local scores along chromosome 4B. The boxes represent the significant zones identified. The threshold was set to $\xi = 2$, and the nominal FDR level is fixed at 0.05.