


Species Distribution Models

Foundations and applied modelling in R



What does a species distribution model do?



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

It links observed records to environmental conditions, then projects that relationship across space.

Records

locations where the
species was observed



Environment

temperature, rainfall,
seasonality, habitat



Suitability

a mapped
hypothesis,
not proof of presence

Suitability is not the same as distribution



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

Species may fail to occupy all suitable habitat, and records may miss places where the species occurs.

- Abiotic conditions can limit survival or reproduction.
- Biotic interactions can shape realised distributions.
- Dispersal barriers can prevent occupation of suitable environments.
- Sampling effort changes what we observe.

SDM output

environmental suitability

a hypothesis for
interpretation

Worked example: *Aedes aegypti* in Brazil




Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

After the foundation, the workshop uses a dengue vector as a practical modelling case.

- GBIF gives reusable occurrence records.
- Brazil provides a national-scale projection exercise.
- The model estimates vector suitability, not dengue risk.
- The same script can be adapted to other species.



Occurrence data: useful, messy, teachable




Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

The first modelling decision is whether the records are good enough for the question.

Cleaned *Aedes aegypti* occurrence records
Brazil, GBIF records with coordinates



Before modelling

- Remove records without coordinates.
- Check the study boundary.
- Thin duplicate raster cells.
- Keep a clean backup for live teaching.


Predictors define the environmental lens



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

We keep four climate predictors so the first model is fast enough and interpretable enough for a 90-minute workshop.



Workshop predictors

- Annual mean temperature
- Temperature seasonality
- Annual precipitation
- Precipitation seasonality

Presence-only data require pseudo-absences



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

Pseudo-absences are comparison points. They are not confirmed absences.

Presences

known occurrence
records



Pseudo-absences

background contrast



Formatted data

ready for biomod2

In the script: `BIOMOD_FormatingData()`

Three algorithms, three modelling assumptions



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

We use multiple algorithms to make model behaviour visible.

GLM

**transparent
statistical trend**

simpler shape

GBM

**boosted sequence
of trees**

flexible shape

RF

**many trees voting
together**

robust shape

Evaluation asks whether predictions generalise



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

Good scores make the model worth inspecting. They do not make the map true.

Training data



Model



Test data

TSS

threshold-based balance of
sensitivity and specificity

AUCroc

threshold-independent ranking
of presences above absences


Variable importance: from output to questions



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

Importance is model sensitivity, not biological proof.



Ask

- Which predictors changed predictions most?
- Do GLM, GBM, and RF agree?
- Could this be ecology, sampling, or correlation?


Ensembles and projections make choices visible




Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

The map is a synthesis of model behaviour across Brazil.



individual model average



ensemble projection


A good SDM should sharpen the next conversation



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

CERI
Centre for Epidemic
Response and Innovation

Now we build the model line by line in R.



Before acting on a map, ask:

- Where is suitability high or low?
- Where are the data thin?
- What could be sampling bias?
- Which predictor story is plausible?
- What dengue surveillance data would you add?