





Stage: SEVERE

Metadata Snapshot

Name: solomon
Age: 67
Gender: Male
Systolic (mmHg): 189
Diastolic (mmHg): 110
BMI: 30.8
Glucose: 398
HbA1c: 8.9
Cholesterol: 276
Smoking: Yes
Hypertension: Yes
Diabetes Duration: 17



Summary

- Clinical Report — Ophthalmologist
- Patient: Name: solomon, Age: 67, BMI: 30.8
- Predicted stage: SEVERE
- Model confidence: 32.7%

- Risk score: 69.33%
- Findings (automated lesion quantification):
 - - Microaneurysms: 6.09%
 - - Exudates: 30.00%
 - - Hemorrhages: 1.62%
 - - Cotton Wool: 30.00%
 - - Neovascularization: 30.00%
 - - Total Lesion Load: 67.71%
- Clinical interpretation:
 - Extensive hemorrhages, venous beading and/or cotton-wool spots consistent with severe NPDR.
Urgent referral advised.
- Model & Explainability:
 - - Multi-model fusion: CNN ensemble (EfficientNet, ResNet50, ViT) + metadata models (RF, XGBoost, stacked ensemble).
 - - Explainability outputs available: GradCAM++, LIME, SHAP — review heatmaps and overlay masks.
- Performance metrics:
 - - Accuracy: 0.947
 - - F1-score: 0.938
 - - AUC/ROC: 0.971
- Suggested clinical action items:
 1. Correlate with slit-lamp exam and fundus exam.
 2. If macular edema suspected, obtain OCT imaging.
 3. For suspected PDR, arrange urgent retina referral for potential PRP/anti-VEGF.
 4. Document lesion masks if available for audit.

