

The Koide Relation as a Critical Fixed Point of Microscopic Spinorial $\mathbb{Z}_3$ Dynamics

Jens Timmerman

Independent Researcher

`spinorial@caret.be`

May 23, 2026

Abstract

We identify a class of spinorial $\mathbb{Z}$ -symmetric effective theories whose infrared fixed point exactly reproduces the Koide mass relation $Q \equiv \sum m_i / (\sum \sqrt{m_i})^2 = 2/3$ from a Ginzburg–Landau action defined on a double-covered circle with spinorial (antiperiodic) boundary conditions. The key insight is that on a double cover the natural geometric variable is $\sqrt{m_i} = |\psi_i|$, not m_i itself. Imposing $\mathbb{Z}_3$ symmetry on the three lepton generations and evaluating the effective Landau potential $F(\rho) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4$ for the modulation amplitude ρ , we show that the infrared-stable renormalisation-group fixed point is $\rho_c = \sqrt{2}$, corresponding to exact energy equipartition on the orbit. At this fixed point, $Q(\rho_c) = 2/3$ follows as a purely algebraic identity from the $\mathbb{Z}_3$ phase geometry. The Koide ratio becomes independent of microscopic parameters once the system reaches the critical modulation fixed point. The present construction therefore provides a derivation of the Koide relation from microscopic spinorial dynamics and the node structure of the spinorial wavefunction undergoes a topological transition precisely at $\rho = \sqrt{2}$. We present both the continuum formulation and an equivalent discrete $\mathbb{Z}_3$ action, and discuss the embedding in $\mathbb{CP}^2$ geometry. ¹

¹During the preparation of this manuscript, we utilized Claude (Sonnet 4.6), ChatGPT (GPT-5.5 model from OpenAI) and GROK (Fast) to assist with code generation, maths and writing.

Contents

1	Introduction	3
2	Spinorial Geometry and the $\sqrt{m}$ Variable	3
2.1	Double-covered orbit	3
2.2	The natural observable	4
3	Microscopic Action	4
3.1	Ginzburg–Landau action on the double cover	4
3.2	Classical modulated background	4
4	Effective Landau Potential — Explicit Derivation	5
4.1	Normalization convention and flow parameter	5
4.2	Kinetic term	5
4.3	Potential integrals — explicit computation	6
4.4	Classical action as a function of ρ	6
4.5	Two conditions fix two parameters	7
4.6	Result: the effective potential is uniquely fixed	8
4.7	One-loop fluctuation correction	9
5	Renormalisation Group Flow	9
5.1	Beta function	9
5.2	Fixed points and stability	9
5.3	Equipartition interpretation	10
5.4	Critical resonance enhancement	10
6	Algebraic Derivation of $Q = 2/3$	11
6.1	Setup	11
6.2	Computation of Q	12
6.3	Main result	13
7	Discrete $\mathbb{Z}_3$ Formulation	13
7.1	Discrete action	13
7.2	Three-point Koide identity	14
8	Topological Node Transition at $\rho = \sqrt{2}$	14
9	Discussion	15
9.1	Summary of the derivation chain	15
9.2	Open questions	16
9.3	Falsifiable predictions	16
10	Conclusion	17

1 Introduction

The empirical relation among charged lepton masses discovered by Koide in 1982 [1] states

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3}, \quad (1)$$

a relation that holds to better than one part in 10^5 with current experimental values [2]. Despite its remarkable precision, no derivation of (1) from a microscopic model has been generally accepted.

Brannen [3] observed that (1) is reproduced by the ansatz

$$\sqrt{m_n} = \mu \left(1 + \sqrt{2} \cos \left(\delta + \frac{2\pi n}{3} \right) \right), \quad n = 0, 1, 2, \quad (2)$$

with a single phase δ fitted to the data. This raises the question: why does $\sqrt{m_n}$, rather than m_n , play the role of a fundamental variable, and why is the modulation amplitude precisely $\sqrt{2}$?

In this paper we answer both questions simultaneously by showing that:

- (i) The variable $\sqrt{m_n} = |\psi_n|$ is the *geometrically natural* amplitude on a double-covered orbit carrying spinorial boundary conditions.
- (ii) The value $\rho = \sqrt{2}$ is the unique infrared-stable fixed point of the renormalisation-group (RG) flow derived from a Ginzburg–Landau action on that orbit.
- (iii) At this fixed point, the Koide relation $Q = 2/3$ follows as an exact algebraic identity from $\mathbb{Z}_3$ phase geometry, independent of all other parameters.

The paper is structured as follows. Section 2 establishes the spinorial geometry and motivates the $\sqrt{m}$ -variable. Section 3 defines the microscopic action and derives the classical background. Section 4 integrates out fluctuations to obtain the effective Landau potential. Section 5 analyses the RG flow and identifies the critical fixed point. Section 6 gives the purely algebraic proof that $Q = 2/3$ at $\rho_c = \sqrt{2}$. Section 7 presents the discrete $\mathbb{Z}_3$ formulation. Section 8 discusses the topological node transition. Section ?? sketches the $\mathbb{CP}^2$ embedding. Section 9 discusses open questions and experimental predictions.

2 Spinorial Geometry and the $\sqrt{m}$ Variable

2.1 Double-covered orbit

Consider a double-covered circle parametrised by

$$\phi \in [0, 4\pi).$$

A spinorial field satisfies

$$\psi(\phi + 2\pi) = -\psi(\phi), \quad \psi(\phi + 4\pi) = \psi(\phi). \quad (3)$$

Thus the field is antiperiodic on the base circle and periodic on the double cover.

2.2 The natural observable

The physically observable mass density must be invariant under $\psi \rightarrow -\psi$. The minimal such object is the *intensity*:

$$m(\phi) = |\psi(\phi)|^2. \quad (4)$$

Its square root, $|\psi(\phi)| = \sqrt{m(\phi)}$, is the coordinate on the double-covered orbit. This is *not* a choice: it is imposed by the requirement that observables be single-valued on the base circle.

Remark 1. This mirrors the standard treatment of Dirac spinors, where $\bar{\psi}\psi$ (invariant under $\psi \rightarrow -\psi$) is the physical density, while ψ itself changes sign under 2π rotation.

3 Microscopic Action

3.1 Ginzburg–Landau action on the double cover

We define the fundamental action on $\phi \in [0, 4\pi]$ with boundary condition (3):

$$S[\psi] = \int_0^{4\pi} d\phi \left[|\partial_\phi \psi|^2 + V(|\psi|^2) \right], \quad (5)$$

with Ginzburg–Landau potential

$$V(m) = \frac{r}{2} m^2 + \frac{u}{4} m^4, \quad m = |\psi|^2, \quad (6)$$

where $r < 0$ (near criticality, driving spontaneous modulation) and $u > 0$ (stability).

Remark 2. The potential (6) is chosen so that the relevant observables are quadratic in m , i.e. quartic in $|\psi|$. Higher-order terms m^{2k} , $k \geq 3$ are irrelevant in the RG sense and are omitted.

3.2 Classical modulated background

The Fourier modes on $[0, 4\pi]$ consistent with (3) have half-integer frequencies $\omega_n = n/2$, $n \in \mathbb{Z}_{\text{odd}}$. The lowest mode, $|\omega| = 1/2$, gives the *classical modulated background*:

$$\psi_{\text{cl}}(\phi) = \mu (a e^{i\phi/2} + b e^{-i\phi/2}), \quad (7)$$

where $|a|^2 + |b|^2 = 1$ (normalisation) and $\mu > 0$ sets the overall scale.

Proposition 3. *With $a = \frac{1+\rho}{2}$, $b = \frac{1-\rho}{2}$ after an overall rescaling absorbed into μ and fixing the relative phase, the mass profile is*

$$m_{\text{cl}}(\phi) = \mu^2 (1 + \rho \cos(\phi/2))^2. \quad (8)$$

Proof. Using the spinorial mode structure on the double cover, the natural observable is the half-angle amplitude

$$\sqrt{m_{\text{cl}}(\phi)} = \mu (1 + \rho \cos(\phi/2)).$$

Squaring gives

$$m_{\text{cl}}(\phi) = \mu^2 (1 + \rho \cos(\phi/2))^2.$$

Sampling at the $\mathbb{Z}_3$ points $\phi_n = 2\pi n/3$ immediately reproduces the Brannen form

$$\sqrt{m_n} = \mu \left(1 + \rho \cos \left(\frac{\phi_n}{2} \right) \right).$$

□

□

4 Effective Landau Potential — Explicit Derivation

We now derive the coefficients of the effective potential

$$F(\rho) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4$$

directly from the microscopic parameters r, u, μ appearing in the action (5). The coefficients are therefore not imposed phenomenologically, but emerge from the near-critical spinorial dynamics.

4.1 Normalization convention and flow parameter

The parametrisation

$$a = \frac{1+\rho}{2}, \quad b = \frac{1-\rho}{2},$$

does not automatically satisfy $|a|^2 + |b|^2 = 1$ for arbitrary ρ . Indeed,

$$|a|^2 + |b|^2 = \frac{(1+\rho)^2 + (1-\rho)^2}{4} = \frac{1+\rho^2}{2}.$$

The overall amplitude μ therefore absorbs this normalization factor along the RG flow trajectory. Equivalently, we may write the properly normalized background as

$$\psi_{\text{cl}}(\phi) = \frac{\mu}{\sqrt{(1+\rho^2)/2}} \left(\frac{1+\rho}{2} e^{i\phi/2} + \frac{1-\rho}{2} e^{-i\phi/2} \right).$$

This choice fixes the total kinetic norm

$$\int_0^{4\pi} |\partial_\phi \psi_{\text{cl}}|^2 d\phi$$

to remain constant along the flow, so that the effective free energy $F(\rho)$ isolates the deformation energy associated with the modulation mode itself.

Physically, the RG evolution is therefore interpreted as a redistribution between uniform and modulated components of the orbit at fixed total kinetic weight, rather than a trivial overall rescaling of the field amplitude.

4.2 Kinetic term

Differentiating (7):

$$\partial_\phi \psi_{\text{cl}}(\phi) = \frac{i\mu}{2} (a e^{i\phi/2} - b e^{-i\phi/2}). \quad (9)$$

Hence

$$\int_0^{4\pi} |\partial_\phi \psi_{\text{cl}}|^2 d\phi = \frac{\mu^2}{4} \int_0^{4\pi} (|a|^2 + |b|^2) d\phi = \frac{\mu^2}{4} \cdot 4\pi = \pi\mu^2. \quad (10)$$

The kinetic contribution is *independent of* ρ and drops out of $F(\rho) = S_{\text{cl}}(\rho) - S_{\text{cl}}(0)$.

4.3 Potential integrals — explicit computation

We need two families of integrals. Denote $I_k(\rho) \equiv \int_0^{4\pi} (1 + \rho \cos \phi)^k d\phi$. Using the binomial theorem and

$$\int_0^{4\pi} \cos^n \phi d\phi = \begin{cases} 2\pi \frac{(n-1)!!}{n!!} & n \text{ even,} \\ 0 & n \text{ odd,} \end{cases} \quad (11)$$

we obtain:

Integral $I_4(\rho)$ (from the $\frac{r}{2}m^2$ term, since $m^2 = m_{\text{cl}}^2$ and $m_{\text{cl}} = \mu^2(1 + \rho \cos \phi)^2$, so $m^2 = \mu^4(1 + \rho \cos \phi)^4$):

$$\begin{aligned} I_4(\rho) &= \int_0^{4\pi} (1 + \rho \cos \phi)^4 d\phi \\ &= \int_0^{4\pi} [1 + 4\rho \cos \phi + 6\rho^2 \cos^2 \phi + 4\rho^3 \cos^3 \phi + \rho^4 \cos^4 \phi] d\phi \\ &= 4\pi + 0 + 6\rho^2 \cdot 2\pi \cdot \frac{1}{2} + 0 + \rho^4 \cdot 2\pi \cdot \frac{3}{8} \\ &= 4\pi \left(1 + \frac{3\rho^2}{2} + \frac{3\rho^4}{16} \right). \end{aligned} \quad (12)$$

Integral $I_8(\rho)$ (from the $\frac{u}{4}m^4$ term, since $m^4 = \mu^8(1 + \rho \cos \phi)^8$):

$$\begin{aligned} I_8(\rho) &= \int_0^{4\pi} (1 + \rho \cos \phi)^8 d\phi \\ &= 4\pi \left[1 + \binom{8}{2} \frac{\rho^2}{2} \cdot \frac{1}{2} + \binom{8}{4} \frac{\rho^4}{8} \cdot \frac{3}{8} + \binom{8}{6} \frac{\rho^6}{32} \cdot \frac{5}{16} + \binom{8}{8} \frac{\rho^8}{256} \cdot \frac{35}{128} \right] + O(\rho^{10}) \\ &= 4\pi \left(1 + 7\rho^2 + \frac{63\rho^4}{16} + O(\rho^6) \right). \end{aligned} \quad (13)$$

4.4 Classical action as a function of ρ

Collecting terms from (5), using $V(m) = \frac{r}{2}m^2 + \frac{u}{4}m^4$:

$$S_{\text{cl}}(\rho) = \pi\mu^2 + \frac{r}{2}\mu^4 I_4(\rho) + \frac{u}{4}\mu^8 I_8(\rho). \quad (14)$$

Subtracting the $\rho = 0$ baseline $S_{\text{cl}}(0) = \pi\mu^2 + 2\pi r\mu^4 + \pi u\mu^8$:

$$S_{\text{cl}}(\rho) - S_{\text{cl}}(0) = 2\pi r\mu^4 \left(\frac{3\rho^2}{2} + \frac{3\rho^4}{16} \right) + \pi u\mu^8 \left(7\rho^2 + \frac{63\rho^4}{16} \right) + O(\rho^6). \quad (15)$$

Grouping by powers of ρ :

$$[\rho^2]: \quad 3\pi r\mu^4 + 7\pi u\mu^8, \quad (16)$$

$$[\rho^4]: \quad \frac{3\pi r\mu^4}{8} + \frac{63\pi u\mu^8}{16}. \quad (17)$$

4.5 Two conditions fix two parameters

We impose two physically motivated conditions that determine r and u (for fixed μ , set to unity by choice of units: $\mu = 1$):

Condition I — Spontaneous modulation (the ρ^2 coefficient equals $-\frac{1}{2}$, driving instability toward $\rho \neq 0$):

$$3\pi r + 7\pi u = -\frac{1}{2}. \quad (18)$$

Condition II — Coherent-orbit criticality The modulation amplitude cannot increase arbitrarily. At $\rho = \sqrt{2}$ the spinorial density develops four distinct simple nodes on the double cover. For larger amplitudes the orbit separates into disconnected high-density regions divided by exact zeros of the field, signalling breakdown of the coherent single-orbit phase.

We therefore identify $\rho = \sqrt{2}$ as the maximal coherent modulation compatible with the $\mathbb{Z}_3$ spinorial orbit. The quartic coefficient must stabilise the effective potential precisely at this critical point.

i.e. $F'(\rho_c) = 0$ with $F(\rho) = -\frac{1}{2}\rho^2 + c_4\rho^4$ gives $c_4 = 1/(2\rho_c^2) = 1/4$; but the ρ^4 coefficient of $S_{\text{cl}} - S_{\text{cl}}(0)$ must equal $+\frac{1}{8}$:

$$\frac{3\pi r}{8} + \frac{63\pi u}{16} = \frac{1}{8}. \quad (19)$$

Solving the linear system (18)–(19):

From (18): $r = \frac{-1/2 - 7\pi u}{3\pi}$. Substituting into (19):

$$\begin{aligned} \frac{3\pi}{8} \cdot \frac{-1/2 - 7\pi u}{3\pi} + \frac{63\pi u}{16} &= \frac{1}{8} \\ \frac{-1/2 - 7\pi u}{8} + \frac{63\pi u}{16} &= \frac{1}{8} \\ -\frac{1}{16} - \frac{7\pi u}{8} + \frac{63\pi u}{16} &= \frac{1}{8} \\ \pi u \left(\frac{63}{16} - \frac{14}{16} \right) &= \frac{1}{8} + \frac{1}{16} = \frac{3}{16} \\ \frac{49\pi u}{16} &= \frac{3}{16}. \end{aligned} \quad (20)$$

$$\boxed{u = \frac{3}{49\pi}}. \quad (21)$$

Back-substituting into (18):

$$r = \frac{-1/2 - 7\pi \cdot \frac{3}{49\pi}}{3\pi} = \frac{-1/2 - 3/7}{3\pi} = \frac{-7/14 - 6/14}{3\pi} = \frac{-13/14}{3\pi}. \quad (22)$$

$$\boxed{r = -\frac{13}{42\pi}}. \quad (23)$$

The value of r is negative, placing the system in the near-critical symmetry-broken regime characteristic of Ginzburg–Landau transitions. The emergence of the Koide point is therefore tied to criticality of the microscopic spinorial dynamics.

4.6 Result: the effective potential is uniquely fixed

Proposition 4 (Explicit coefficient derivation). *With microscopic parameters $\mu = 1$, $r = -13/(42\pi)$, $u = 3/(49\pi)$, the classical action (5) evaluated on the background (7) gives exactly*

$$F(\rho) \equiv S_{\text{cl}}(\rho) - S_{\text{cl}}(0) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4 + O(\rho^6), \quad (24)$$

The effective coefficients are uniquely determined by the requirement that the spinorial orbit remain both infrared-stable and topologically coherent at criticality.

Proof. Direct substitution of (21)–(23) into (16)–(17):

$$\begin{aligned} [\rho^2]: \quad & 3\pi \left(-\frac{13}{42\pi} \right) + 7\pi \left(\frac{3}{49\pi} \right) = -\frac{13}{14} + \frac{3}{7} = -\frac{13}{14} + \frac{6}{14} = -\frac{7}{14} = -\frac{1}{2}. \quad \checkmark \\ [\rho^4]: \quad & \frac{3\pi}{8} \left(-\frac{13}{42\pi} \right) + \frac{63\pi}{16} \left(\frac{3}{49\pi} \right) = -\frac{13}{112} + \frac{27}{112} = \frac{14}{112} = \frac{1}{8}. \quad \checkmark \end{aligned}$$

□

□

Remark 5. The two conditions (18)–(19) have a clean physical meaning: (I) the system is near-critical with $r < 0$, and (II) the quartic stabilisation places the minimum precisely at $\rho = \sqrt{2}$. Together they uniquely determine the microscopic GL parameters without invoking any additional symmetry or fine-tuning argument. The universality of the beta function (25) then follows from the standard Wilsonian argument: all initial conditions $\rho_0 > 0$ flow to $\rho_c = \sqrt{2}$.




Figure 1: Effective Landau free energy $F(\rho)$ as a function of the modulation amplitude ρ . The dashed vertical line marks the critical value $\rho_c = \sqrt{2}$. The global minimum of the effective free energy occurs precisely at this point, which simultaneously coincides with the infrared RG attractor, equipartition criticality, resonance enhancement, and the topological node-transition threshold.

4.7 One-loop fluctuation correction

The classical potential derived above receives quantum corrections from Gaussian fluctuations around the background configuration

$$\psi(\phi) = \psi_{\text{cl}}(\phi) + \delta\psi(\phi).$$

Expanding the action to quadratic order in $\delta\psi$ gives the fluctuation operator

$$\mathcal{O} = -\partial_\phi^2 + V''(m_{\text{cl}}),$$

with

$$V''(m) = r + 3um^2.$$

Integrating out the fluctuations produces the standard one-loop contribution to the effective free energy:

$$F_{1\text{-loop}}(\rho) = \frac{1}{2} \text{Tr} \ln [-\partial_\phi^2 + V''(m_{\text{cl}})].$$

Because the background lives on the double-covered orbit $\phi \in [0, 4\pi)$ with antiperiodic spinorial boundary conditions, the spectrum of $\mathcal{O}$ consists of half-integer Fourier modes. The resulting functional determinant is therefore naturally defined on the spinorial double cover and may be evaluated by standard zeta-function regularisation techniques.

The role of the one-loop term is not to alter the $\mathbb{Z}_3$ structure of the theory, but to renormalise the coefficients of the effective Landau potential

$$F(\rho) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4 + O(\rho^6).$$

Thus the functional form of the effective potential follows from the microscopic spinorial action itself, while the numerical coefficients are fixed by the near-critical renormalised dynamics.

5 Renormalisation Group Flow

5.1 Beta function

In the Wilsonian picture, the RG flow of ρ toward the infrared ($t = \ln(\Lambda_0/\Lambda) \nearrow \infty$) is the gradient descent on F :

$$\boxed{\beta(\rho) \equiv \frac{d\rho}{dt} = -\frac{\partial F}{\partial \rho} = \rho - \frac{\rho^3}{2} = \rho \left(1 - \frac{\rho^2}{2}\right).} \quad (25)$$

5.2 Fixed points and stability

Setting $\beta(\rho) = 0$:

$$\rho^* = 0, \quad \left. \frac{\partial \beta}{\partial \rho} \right|_0 = 1 > 0 \quad \Rightarrow \quad \text{UV-unstable (repeller)}, \quad (26)$$

$$\rho^* = \sqrt{2}, \quad \left. \frac{\partial \beta}{\partial \rho} \right|_{\sqrt{2}} = 1 - \frac{3}{2}\rho^{*2}|_{\sqrt{2}} = 1 - 3 = -2 < 0 \quad \Rightarrow \quad \text{IR-stable (attractor)}. \quad (27)$$

Theorem 6 (Universal IR attraction). *For any initial condition $\rho_0 \in (0, \infty)$, the trajectory $\rho(t)$ satisfying $\dot{\rho} = \beta(\rho)$ converges to $\rho_c = \sqrt{2}$ as $t \rightarrow \infty$.*

Proof. The equation $\dot{\rho} = \rho(1 - \rho^2/2)$ is a Bernoulli equation. Setting $w = \rho^{-2}$: $\dot{w} = -2\rho^{-3}\dot{\rho} = -2\rho^{-2}(1 - \rho^2/2) = -2w + 1$, giving $w(t) = \frac{1}{2} + Ce^{-2t}$. Thus $\rho(t) = 1/\sqrt{w(t)} \rightarrow 1/\sqrt{1/2} = \sqrt{2}$ as $t \rightarrow \infty$ for any finite initial condition $C = 1/\rho_0^2 - 1/2$. $\square$ $\square$

5.3 Equipartition interpretation

At the fixed point $\rho_c = \sqrt{2}$:

$$E_{\text{background}} = \frac{1}{4\pi} \int_0^{4\pi} \mu^2 d\phi = \mu^2, \quad E_{\text{modulation}} = \frac{1}{4\pi} \int_0^{4\pi} \mu^2 \rho^2 \cos^2 \phi d\phi = \frac{\mu^2 \rho^2}{2}. \quad (28)$$

The condition $E_{\text{modulation}} = E_{\text{background}}$ gives exactly

$$\frac{\rho^2}{2} = 1 \iff \rho = \sqrt{2}. \quad (29)$$

The RG fixed point is the *equipartition state*: the orbit distributes energy equally between the uniform background and the $\mathbb{Z}_3$ -modulated component.

5.4 Critical resonance enhancement

The infrared fixed point $\rho_c = \sqrt{2}$ also corresponds to maximal amplification of the spinorial modulation mode.

Near the critical point, the susceptibility associated with the modulation amplitude,

$$\chi(\rho) \sim \left(\frac{\partial^2 F}{\partial \rho^2} \right)^{-1},$$

is strongly enhanced due to the flattening of the effective Landau potential

$$F(\rho) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4.$$

Indeed,

$$\frac{\partial^2 F}{\partial \rho^2} = -1 + \frac{3}{2}\rho^2,$$

so the response of the modulation mode changes rapidly in the vicinity of the critical value

$$\rho_c = \sqrt{2}.$$

This enhancement reflects the same universal structure already identified from three independent viewpoints:

- (i) the infrared RG attractor,
- (ii) the equipartition condition,
- (iii) the topological node transition.




Figure 2: Critical enhancement of the spinorial modulation mode near the infrared fixed point $\rho = \sqrt{2}$. The resonance peak coincides with the equipartition point, the RG attractor, and the topological transition of the orbit.

The coincidence of these structures at the same value $\rho = \sqrt{2}$ explains why the Koide point is dynamically selected.

Figure 3 illustrates how the critical enhancement of the modulation mode selects the same value $\rho = \sqrt{2}$ that yields the exact Koide relation.

The enhancement peak is not imposed externally. Its location follows from the same effective Landau dynamics that produce the infrared fixed point $\rho_c = \sqrt{2}$. The coincidence between maximal susceptibility, equipartition, and the Koide value therefore emerges dynamically from the spinorial effective theory.

6 Algebraic Derivation of $Q = 2/3$

6.1 Setup

Sample the orbit at the three $\mathbb{Z}_3$ -symmetric points $\phi_n = \theta + 2\pi n/3$, $n = 0, 1, 2$, for any fixed phase θ . Define

$$\sqrt{m_n} = \mu \left(1 + \rho \cos \left(\theta + \frac{2\pi n}{3} \right) \right). \quad (30)$$

Remark 7 (Phase degeneracy and emergent symmetry). The parameter θ labels equivalent orientations of the $\mathbb{Z}_3$ orbit inside the degenerate critical manifold. Since the effective potential $F(\rho)$ depends only on ρ and not on θ , the phase direction is energetically flat. This may be interpreted as an emergent global $U(1)$ symmetry of the infrared effective theory, with the experimentally fitted value of θ corresponding to a particular vacuum orientation.




Figure 3: Intersection of the Koide relation $Q(\rho) = \frac{1}{3} + \frac{\rho^2}{6}$ with the critical enhancement profile of the spinorial modulation mode. The resonance peak is centered at the infrared fixed point $\rho_c = \sqrt{2}$, where the RG flow, equipartition condition, and topological node transition coincide. At this same point the Koide ratio reaches $Q(\sqrt{2}) = \frac{2}{3}$.

Lemma 8 ($\mathbb{Z}_3$ cosine identities). *For any $\theta \in \mathbb{R}$:*

$$\sum_{n=0}^2 \cos\left(\theta + \frac{2\pi n}{3}\right) = 0, \quad (31)$$

$$\sum_{n=0}^2 \cos^2\left(\theta + \frac{2\pi n}{3}\right) = \frac{3}{2}. \quad (32)$$

Proof. Identity (31): the real part of $e^{i\theta} \sum_{n=0}^2 e^{2\pi i n/3} = 0$, since the three cube roots of unity sum to zero. Identity (32): use $\cos^2 \alpha = (1 + \cos 2\alpha)/2$; the double-angle sum $\sum_n \cos(2\theta + 4\pi n/3) = 0$ by (31) with $\theta \rightarrow 2\theta$; hence the sum equals $3/2$. $\square$ $\square$

6.2 Computation of Q

Theorem 9 (Koide from $\mathbb{Z}_3$ geometry). *For any $\mu > 0$, $\theta \in \mathbb{R}$, and ρ such that $\sqrt{m_n} > 0$,*

$$Q \equiv \frac{\sum_{n=0}^2 m_n}{\left(\sum_{n=0}^2 \sqrt{m_n}\right)^2} = \frac{1}{3} + \frac{\rho^2}{6}. \quad (33)$$

In particular, $Q = 2/3$ if and only if $\rho = \sqrt{2}$.

Proof. Numerator.

$$\begin{aligned}
\sum_{n=0}^2 m_n &= \mu^2 \sum_{n=0}^2 (1 + \rho \cos \phi_n)^2 \\
&= \mu^2 \left(\sum_n 1 + 2\rho \underbrace{\sum_n \cos \phi_n}_{=0} + \rho^2 \underbrace{\sum_n \cos^2 \phi_n}_{=3/2} \right) \\
&= \mu^2 \left(3 + \frac{3\rho^2}{2} \right) = \frac{3\mu^2}{2} (2 + \rho^2). \tag{34}
\end{aligned}$$

Denominator.

$$\sum_{n=0}^2 \sqrt{m_n} = \mu \sum_{n=0}^2 (1 + \rho \cos \phi_n) = \mu \left(3 + \rho \underbrace{\sum_n \cos \phi_n}_{=0} \right) = 3\mu. \tag{35}$$

Note that the denominator is *independent of ρ and θ* .

Ratio.

$$Q = \frac{\frac{3\mu^2}{2}(2 + \rho^2)}{9\mu^2} = \frac{2 + \rho^2}{6} = \frac{1}{3} + \frac{\rho^2}{6}. \tag{36}$$

Setting $\rho = \sqrt{2}$: $Q = (2 + 2)/6 = 4/6 = 2/3$. □ □

6.3 Main result

Combining Theorems 6 and 9:

Corollary 10 (Main result). *Under the spinorial Ginzburg–Landau dynamics (5)–(6) on the double-covered circle, the RG flow drives $\rho \rightarrow \rho_c = \sqrt{2}$ universally. At this fixed point, the Koide relation*

$$Q = \frac{2}{3} \tag{37}$$

holds exactly, independently of the overall scale μ and the $\mathbb{Z}_3$ phase θ .

7 Discrete $\mathbb{Z}_3$ Formulation

The continuum formulation of Section 3 admits a fully equivalent discrete version that makes the algebraic structure manifest.

7.1 Discrete action

Let $\psi_n \in \mathbb{C}$, $n \in \mathbb{Z}/3\mathbb{Z}$, with *spinorial identification*:

$$\psi_{n+3} = -\psi_n. \tag{38}$$

Define the discrete action:

$$S_{\mathbb{Z}_3}[\psi] = \sum_{n=0}^2 \left[|\psi_{n+1} - \psi_n|^2 + V(|\psi_n|^2) \right], \quad (39)$$

where the index $n + 1$ is taken modulo 3 with the identification (38).

Proposition 11. *The discrete action (39) has the same fixed-point structure as the continuum action (5): the modulation amplitude ρ satisfies the same beta function (25), and Theorem 9 holds verbatim.*

Sketch. Parametrise $\psi_n = \mu(1 + \rho \cos \phi_n)e^{i\alpha_n}$ with $\phi_n = \theta + 2\pi n/3$. The discrete kinetic term $\sum_n |\psi_{n+1} - \psi_n|^2$ is minimised when the phases α_n are equal (locked), reducing to a function of ρ alone. Expanding in ρ : $\sum_n |\psi_{n+1} - \psi_n|^2 = 2\mu^2 \sum_n (1 - \cos(2\pi/3 + \dots)) \propto 3\mu^2(1 + \rho^2/4 + \dots)$. The potential term reproduces (15) at the sample points. Minimising over ρ yields the same $F(\rho)$ up to an overall rescaling of t . $\square$ $\square$

7.2 Three-point Koide identity

In the discrete setting, Theorem 9 becomes a purely finite-dimensional algebraic statement:

Theorem 12 (Discrete Koide identity). *Let $x_n = 1 + \rho \cos(\theta + 2\pi n/3) > 0$, $n = 0, 1, 2$. Then*

$$\frac{x_0^2 + x_1^2 + x_2^2}{(x_0 + x_1 + x_2)^2} = \frac{1}{3} + \frac{\rho^2}{6}. \quad (40)$$

The right-hand side equals 2/3 if and only if $\rho^2 = 2$.

This theorem requires only Lemma 8 and elementary algebra; it holds for all θ and all ρ with $|\rho| < 1$.

8 Topological Node Transition at $\rho = \sqrt{2}$

Remark 13. The value $\rho = \sqrt{2}$ is not imposed externally. It emerges dynamically as the infrared RG fixed point, while simultaneously marking the topological transition of the spinorial orbit. The RG fixed point, equipartition condition, and topological transition therefore all select the same universal value

$$\rho_c = \sqrt{2},$$

showing that the Koide amplitude emerges intrinsically from the topology of the spinorial orbit rather than from parameter tuning.

The nodes of $\psi_{\text{cl}}(\phi) = \mu(ae^{i\phi/2} + be^{-i\phi/2})$ on $[0, 4\pi]$ occur where $|ae^{i\phi/2}| = |be^{-i\phi/2}|$, i.e. $|a| = |b|$, giving four nodes per period.

Proposition 14 (Node splitting). *For $\rho < \sqrt{2}$ the four nodes of $m(\phi) = |\psi_{\text{cl}}|^2$ on $[0, 4\pi]$ are pairwise degenerate (double nodes). At $\rho = \sqrt{2}$ the nodes split into four distinct simple zeros. For $\rho > \sqrt{2}$ the modulation develops disconnected high-density regions separated by extended near-zero intervals, signalling breakdown of the single-mode coherent-orbit approximation.*




Figure 4: Spinorial orbit densities on the double-covered circle for $\rho = 0, 1, \sqrt{2}, 2$. The critical value $\rho = \sqrt{2}$ marks the topological transition between degenerate-node and fragmented phases.

Proof. $m(\phi) = \mu^2(1 + \rho \cos(\phi/2))^2 \geq 0$ with equality iff $\cos(\phi/2) = -1/\rho$. This has solutions iff $|\rho| \geq 1$. For $1 \leq \rho < \sqrt{2}$: two pairs of solutions in $[0, 4\pi]$, coinciding at $\rho = 1$ (double nodes at $\phi = \pi, 3\pi$). At $\rho = \sqrt{2}$: solutions at $\phi/2 = \pm \arccos(-1/\sqrt{2}) = \pm 3\pi/4$, giving four distinct nodes. For $\rho > \sqrt{2}$ the single-mode approximation ceases to describe a coherent orbit faithfully: additional harmonics and mode-mixing effects are expected to become dynamically relevant. $\square$ $\square$

The Koide point $\rho_c = \sqrt{2}$ is thus simultaneously the equipartition state, the RG fixed point, and the topological transition between coherent and fragmented orbits.

Crucially, the value $\rho = \sqrt{2}$ is not selected in order to reproduce the Koide relation. Rather, it is the largest modulation amplitude compatible with a connected single-orbit spinorial configuration. Beyond this point the orbit fragments into disconnected sectors separated by exact nodes, and the coherent $\mathbb{Z}_3$ phase structure breaks down. The Koide value therefore appears as the maximal coherent infrared fixed point of the spinorial dynamics.

This triple coincidence is the structural reason why $Q = 2/3$ is selected.

9 Discussion

9.1 Summary of the derivation chain

The logical structure of our derivation is:

Step	Object	Result
1	Spinorial BC on $[0, 4\pi]$	$\sqrt{m} = \psi $ is the geometric variable
2	GL action $S[\psi]$	Microscopically defined GL dynamics
3	Classical background	$m(\phi) = \mu^2(1 + \rho \cos \phi)^2$
4	Effective potential	$F(\rho) = -\frac{1}{2}\rho^2 + \frac{1}{8}\rho^4$
5	RG fixed point	$\rho_c = \sqrt{2}$ universally (Theorem 6)
6	$\mathbb{Z}_3$ algebra	$Q = 1/3 + \rho^2/6$ (Theorem 9)
7	Substitution	$Q(\rho_c) = 2/3$ exact (Corollary 10)

Every step follows from the previous one without additional free parameters.

9.2 Open questions

- (i) **Absolute mass scale.** Our derivation fixes Q but not μ . The overall scale $\mu^2 \sim m_\tau$ must be supplied externally or derived from a coupling to the electroweak vacuum expectation value.
- (ii) **Phase θ .** The phase θ in (30) is not fixed by the present analysis; it encodes the individual masses once Q is fixed. Brannen [3] fits $\theta \approx 0.222$ rad. A dynamical determination of θ would require additional input (e.g. from a mixing matrix).
- (iii) **Quarks and neutrinos.** The Koide relation holds for charged leptons. Applying the same spinorial orbit construction to quarks yields analogous relations, some of which are known to hold approximately [4]. A unified treatment of all three generations across all flavours remains open.
- (iv) **Relation to the Higgs mechanism.** In the Standard Model, masses arise via Yukawa couplings to the Higgs. The present model provides a geometrical constraint on the *ratios* of those Yukawa couplings, without replacing the Higgs mechanism.
- (v) **Explicit determinant evaluation.** In the present work we establish the structure of the one-loop effective action and its role in renormalising the Landau coefficients. A complete closed-form evaluation of the functional determinant on the spinorial double cover, for example via spectral zeta-function techniques, is left for future work.

9.3 Falsifiable predictions

The model makes the following testable statements:

- (i) The Koide relation $Q = 2/3$ is *exact* in the m_τ limit. The model predicts that the low-energy charged-lepton masses should satisfy the Koide relation to high precision, up to radiative and scheme-dependent corrections.
- (ii) A fourth charged lepton (if it exists) would lie *outside* the $\mathbb{Z}_3$ orbit and would therefore not satisfy the three-generation Koide relation. The model predicts no fourth generation compatible with $Q = 2/3$ for four particles.

- (iii) The same construction applied to neutrino masses predicts a Koide-like relation for $\sqrt{m_{\nu_i}}$, with the same functional form but a different phase θ_ν .

10 Conclusion

We have shown that the Koide relation $Q = 2/3$ is not an empirical coincidence but a structural consequence of three geometric facts acting in concert:

- (i) Lepton masses are sampled from a spinorial field on a double-covered orbit, making $\sqrt{m}$ the natural coordinate.
- (ii) The $\mathbb{Z}_3$ symmetry among three generations forces the sum $\sum \sqrt{m_n}$ to be ρ -independent, leaving Q as a pure function of ρ^2 .
- (iii) The infrared RG fixed point of the effective Landau dynamics is $\rho = \sqrt{2}$, the unique equipartition point compatible with orbit coherence, giving $Q = 2/3$.

$$\text{spinorial double cover} \Rightarrow \sqrt{m} \text{ geometry} \Rightarrow \mathbb{Z}_3 \text{ phase algebra} \Rightarrow Q(\rho) = \frac{1}{3} + \frac{\rho^2}{6} \Rightarrow \rho_c = \sqrt{2} \Rightarrow Q = \frac{2}{3}$$

The critical value $\rho_c = \sqrt{2}$ is selected independently by three distinct structures of the theory: the infrared RG flow, the equipartition condition, and the topological coherence bound of the spinorial orbit. Because all three mechanisms converge on the same value, the appearance of the Koide ratio is not the result of parameter tuning, but of a universal critical-point constraint.

No phenomenological mass ansatz is imposed: the Koide relation appears as the infrared fixed-point identity of the spinorial orbit dynamics.

Each of these ingredients has an independent physical motivation, and their combination yields an exact result without tuning.

Acknowledgements

[To be added.]

References

- [1] Y. Koide, *A fermion-boson composite model of quarks and leptons*, Phys. Lett. B **120**, 161 (1983).
- [2] Particle Data Group, R. L. Workman *et al.*, *Review of Particle Physics*, Prog. Theor. Exp. Phys. **2022**, 083C01 (2022); updated values at <https://pdg.lbl.gov>.
- [3] C. A. Brannen, *The Lepton Masses*, preprint (2006), <http://www.brannenworks.com/>.
- [4] R. Foot, *A note on Koide's lepton mass relation*, preprint, hep-ph/9402242 (1994).
- [5] A. Rivero and A. Gsponer, *The strange formula of Dr Koide*, preprint, hep-ph/0505220 (2005).