

Title

Alexander F. Jercher^{1,2,*} and Sebastian Steinhaus^{1,†}

¹*Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena,
Max-Wien-Platz 1, 07743 Jena, Germany, EU*

²*corresponding author*

(Dated: October 27, 2025)

I. INTRODUCTION

$8\pi G_N = 1?$, spacetime signature $(+, -, -, -)$

II. SETUP

A. Lorentzian Regge calculus

Regge calculus [1] is a discrete gravitational theory on simplicial manifolds Δ with the length of edges $\{l_e\}$ dynamical. In a Lorentzian setting, a simplicial manifold consists of intrinsically flat Lorentzian 4-simplices $\sigma \in \Delta$. Curvature is located at triangles $t \in \Delta$ and captured by deficit angles $\delta_t(\{l_e\})$, lying in the space orthogonal to t . For t spacelike (timelike), this space is isomorphic to $\mathbb{R}^{1,1}$ ($\mathbb{R}^2$).¹ Lorentzian deficit angles are defined as $\delta_t^{L,\pm} = \mp i2\pi - \sum_{\sigma \supset t} \psi_{\sigma,t}^{L,\pm}$, following the conventions of [2, 3], where the $\psi_{\sigma,t}$ are dihedral angles. “ $\pm$ ” indicates a choice of sign which becomes important for causally irregular configurations [2–4] discussed below. The bulk action of Lorentzian Regge calculus is given by [1–3]

$$S_R[\{l_e\}] = \sum_{t \in \Delta: \text{tl}} |A_t(\{l_e\})| \delta_t^E(\{l_e\}) + \sum_{t \in \Delta: \text{sl}} |A_t(\{l_e\})| \delta_t^{L,\pm}(\{l_e\}), \quad (1)$$

with $|A_t|$ the absolute value of the triangle area. In the presence of a boundary, $\partial\Delta \neq \emptyset$, S_R attains boundary terms such that overall additivity is ensured. The equations of motion are given by $\frac{\partial S_R}{\partial l_e} = 0$ supplemented by the Schläfli identity, $\sum_t |A_t| \frac{\partial \delta_t}{\partial l_e} = 0$.

Regge calculus can be formulated in terms of different variables [5, 6], with area variables [7–9] playing a particularly important role here due to their connection to LQG and spin-foams. The relation between area and length variables is in general not invertible, and area variables need to be further constrained to yield the dynamical equations of length Regge calculus [3, 6, 9]. However,

in the symmetry reduced setting introduced below, the relation of area and length is globally invertible and the aforementioned constraints trivialize. For the remainder, we study the classical dynamics of area Regge calculus which will be used in future work as a comparison to results of the effective spin-foam path integral.

B. Lorentzian 4-frusta

Lorentzian 4-dimensional frusta are ideal to capture spatially flat, homogeneous and isotropic discrete geometries [10, 11]. The boundary of a 4-frustum, depicted in Fig. 1, contains two spacelike 3-cubes and six 3-frusta that are either spacelike or timelike. Gluing 4-frusta along boundary 3-frusta in spatial direction and along boundary 3-cubes in temporal direction, one obtains an extended cosmological spacetime of hypercubical combinatorics $\mathcal{X}_V^{(4)}$ with slices labelled by $n \in \{0, \dots, \mathcal{V}_T\}$ discretized by $\mathcal{V}_S^3$ spacelike 3-cubes. We say that a 4-frustum lies in a “slab” between slices n and $n+1$.

The entire geometry of a slab is captured by three variables describing the 3-cubes and 3-frusta which can be either areas or lengths. The former set of variables consists of the area of squares contained in the 3-cubes, j_0 and j_1 , and the area of the trapezoids contained in the 3-frusta, k . The latter consists of the edge length contained in the 3-cubes, s_0 and s_1 , and the height of the 4-frustum, H_0 , which implicitly determines the trapezoids and 3-frusta.² In the highly symmetry restricted setting considered here, there is a one-to-one map between area variables, (j_0, j_1, k_0) , and length variables, (s_0, s_1, H_0) , given explicitly in Appendix A. We will exploit this relation in Sec. IV using length variables for analytical computations and clear interpretation, especially because the height is directly related to the lapse function N in the continuum. When solving the Regge equations numerically thereafter, we will use area variables instead to provide a classical benchmark for the effective spin-foam

* alexander.jercher@uni-jena.de

† sebastian.steinhaus@uni-jena.de

¹ For the remainder, we neglect lightlike edges, triangles and tetrahedra. See [2] for the definition of deficit angles associated to $(d-1)$ -dimensional null cells.

² Strictly speaking, the height H_0 is a derived quantity that is not a length associated to a 1-dimensional sub-cell of the cellular complex. Such a quantity would instead be given by the length l_0 associated to the strut connecting two neighboring spacelike slices. However, the relation between strut length and height is invertible for non-null struts, which assume here.

$$\text{sub-cells} \left| -\frac{\Delta j^2}{8} < k^2 < -\frac{\Delta j^2}{16} \right| -\frac{\Delta j^2}{16} < k^2 < 0 \left| 0 < k^2 < \frac{\Delta j^2}{16} \right| \frac{\Delta j^2}{16} < k^2$$

TABLE I: ...




FIG. 1: Boundary of a 4-frustum given by two 3-cubes and six equal 3-frusta.

As discussed in [11, 13–16], two of these sectors I and II exhibit an irregular light cone structure, also referred to as causality violations, located at spacelike sub-cells. Examples of such irregularities are given by the trouser singularity and the yarmulke singularity associated to spatial topology change [17–19]. For causality violations at $(d - 2)$ -dimensional spacelike hinges the imaginary parts of the dihedral angles do not sum up to $\mp i2\pi$, leading to a complex valued Regge action. Choosing “+” in the convention of Lorentzian angles, the amplitude e^{iS_R} is exponentially suppressed for trouser and exponentially enhanced for yarmulke configurations, vice versa for “−” [3]. Without a complexification of the vari-

calculations of [12]. The general notions of Regge calculus introduced above are straightforwardly adapted to either of the two sets of variables.

The squared volume of subcells connecting neighboring slices, i.e. 3-frusta, trapezoids and edges, which are detailed in Appendix A, carries information on the causal structure. If it is positive (negative), the corresponding subcell is timelike (spacelike). In the symmetry reduced setting, the ratio $\frac{|j_n - j_{n+1}|}{4k_n}$ fully characterizes the causal character of subcells suggesting separating the 4-frusta configurations into three sectors summarized in Tab. I.

ables, the Regge equations of motion for causality violating configurations are complex equations with real variables and therefore overconstrained. Moreover, the results of [11, 13] show that in the respective settings, no real solutions exist. Since this applies also to the setting used in this work we henceforth focus on the causally regular sector III.

C. Lorentzian Regge action for spatially flat cosmology

For a single 4-frustum between, say slice n and $n + 1$, the causally regular Regge action is given by

$$S_{\text{III}}^{(n)} = 6(j_n - j_{n+1}) \sinh^{-1} \left(\frac{j_{n+1} - j_n}{\sqrt{16k_n^2 + (j_n - j_{n+1})^2}} \right) + 12k_n \left[\frac{\pi}{2} - \cos^{-1} \left(\frac{(j_n - j_{n+1})^2}{16k_n^2 + (j_n - j_{n+1})^2} \right) \right] - \Lambda V^{(4)}, \quad (2)$$

where Λ is a cosmological constant and $V^{(4)}$ is the 4-volume defined in Eq. (A4) as an equation of (j_n, j_{n+1}, k_n) . By construction, the 4-frustum action is additive such that on a discretization $\mathcal{X}_V^{(4)}$ with $\mathcal{V}_S^3$ vertices in spatial direction and $\mathcal{V}_T + 1$ spatial slices, the action is given by $S_R = \mathcal{V}_S^3 \sum_{n=0}^{\mathcal{V}_T-1} S_R^{(n)}$. Additivity of the action per slab implies that the equations of motion for a given geometric quantity, being either j_n or k_n , will only depend on neighboring geometric labels.

In the following, we study the discrete dynamics of spatially flat cosmology as governed by the Regge equations. To that end, we consider first the case of a single time step, $\mathcal{V}_T = 1$, in Sec. III and generalize then to an arbitrary number of time steps in Sec. IV.

III. RECAP: ONE TIME STEP

The classical and quantum dynamics of a single timestep in discrete Lorentzian cosmology have been studied extensively in [4, 13, 14] for spatial curvature with cosmological constant and in [11, 16] for spatially flat spacetimes with a free scalar field. In this chapter we extend these studies to spatially flat cosmology with a cosmological constant, before then focussing on the case of arbitrarily many time steps.

For $\mathcal{V}_T = 1$, the square areas j_0, j_1 are boundary data and only the trapezoid area k_0 is dynamical. Thus, the Regge equations are simply given by

$$\frac{\partial S_{\text{III}}}{\partial k_0} = 0, \quad (3)$$

where due to the Schläfli identity, the first term in Eq. (2) drops out. Notice that this equation is transcendental and can therefore only be tackled numerically.

Vanishing cosmological constant. In the case of $\Lambda = 0$, the Regge action describes a spatially flat vacuum universe. If the boundary data is non-stationary, i.e. $j_0 \neq j_1$, Eq. (3) exhibits no solution for k_0 . In other words, a classical universe can neither expand nor contract if there is no matter present. This is in accordance with the continuum Friedmann equations in vacuum, $(\frac{\dot{a}}{a})^2 = 0$, with a the scale factor, which are solved only by $a = \text{const.}$ On the other hand, if $j_0 = j_1$, Eq. (3) is trivially satisfied and the trapezoid area k_0 is undetermined. In fact, the Regge action vanishes, $S_{\text{III}} = 0$ for all k_0 , and the 4-frusta reduce to 4-cuboids discretizing flat space. The arbitrariness of k is a common phenomenon for flat solutions in Regge calculus [20, 21] and signifies the restoration of diffeomorphism symmetry in the discrete [22, 23]. This symmetry amounts to moving spacelike slices (instead of individual vertices), see also the discussion in [24]. Note that for 4-cuboids, $k = \sqrt{j}H$ with H the height of the cuboid which is directly related to the lapse function N in the continuum [11, 13]. The lapse is the gauge variable of the residual diffeomorphism symmetry after symmetry reduction, encoding time reparametrization invariance.

Non-vanishing cosmological constant. For a non-vanishing cosmological constant, $\Lambda \neq 0$, and stationary boundary conditions, $j_0 = j_1$, the only solution to Eq. (3) is $k_0 = 0$, corresponding to degenerate 4-frusta with vanishing height. Thus, a stationary universe for $\Lambda \neq 0$ only exists if no time passes, i.e. if there is no actual time evolution. The more interesting case is $j_0 \neq j_1$ where a non-trivial solution k_0^* for Eq. (3) exists. More precisely, a unique solution k_0^* is found for any $\Lambda > 0$ and any $j_0 \neq j_1$. The on-shell trapezoid area $k_0^*(\Lambda)$ as a function of Λ behaves as a monomial decay $k_0^* \sim c\Lambda^{-\alpha}$ with $c, \alpha > 0$ constants depending on j_0, j_1 . For $\Lambda \neq 0$, the solution diverges and ceases to exist at $\Lambda = 0$ in lines with the discussion above. In the limit $\Lambda \rightarrow \infty$, Eq. (3) is dominated by the Λ -term, i.e.

$$\Lambda \frac{(j_0 + j_1)^2 k}{4V^{(4)}} \approx 0, \quad (4)$$

solved by $k_0^* = 0$.

Note that the restriction on the sign of Λ is in accordance with the continuum Friedmann equations, $(\frac{\dot{a}}{a})^2 = N^2 \frac{\Lambda}{3}$, where real solutions for the scale factor require $\Lambda > 0$.

For completeness, let us contrast these results with the existence of solutions in the spatially spherical case in Lorentzian [13] and Euclidean [25, 26] signature.³ In the

former setting, note first that for $\Lambda = 0$, no solutions exist. Furthermore, for $\Lambda > 0$, solutions exist only if $j_0 > j_\Lambda$ and $j_1 > j_\Lambda$, where j_Λ is a Λ -dependent minimal area. This is in accordance with the continuum where the scale factor typically follows a cosh-function, attaining a minimal value a_Λ . In Euclidean signature, solutions exist for $\Lambda = 0$. These correspond, in the continuum, to solutions where $\dot{a} = \text{const.}$ For $\Lambda > 0$, solutions exist only if $j_0 < j_\Lambda$ and $j_1 < j_\Lambda$ where j_Λ is the same as in the Lorentzian case, however now representing an upper bound. Again, this behavior agrees with continuum results where a Euclidean universe with positive spatial curvature can only reach a maximum value a_Λ .

IV. ARBITRARY NUMBER OF TIME STEPS

In this section, we study the Regge equations of spatially flat Lorentzian cosmology for an arbitrary number of time steps $\mathcal{V}_T$. This amounts to solving $2\mathcal{V}_T - 1$ coupled transcendental equations for the bulk variables $k_0, j_1, \dots, j_{\mathcal{V}_T-1}, k_{\mathcal{V}_T-1}$ for given boundary data $j_0, j_{\mathcal{V}_T}$ and cosmological constant Λ ,

$$\frac{\partial S_{\text{III}}}{\partial k_n} = 0, \quad \frac{\partial S_{\text{III}}}{\partial j_m} = 0, \quad (5)$$

with $n \in \{0, \dots, \mathcal{V}_T - 1\}$ and $m \in \{1, \dots, \mathcal{V}_T - 1\}$. Since the Regge action S_{III} is additive with respect to the slices, i.e. $S_{\text{III}} = \mathcal{V}_S^3 \sum_n S_{\text{III}}^{(n)}$, Eqs. (5) simplify significantly,

$$\begin{aligned} 0 &= \frac{\partial S_{\text{III}}}{\partial k_n} = \mathcal{V}_S^3 \frac{\partial S_{\text{III}}^{(n)}}{\partial k_n}, \\ 0 &= \frac{\partial S_{\text{III}}}{\partial j_n} = \mathcal{V}_S^3 \left(\frac{\partial S_{\text{III}}^{(n)}}{\partial j_n} + \frac{\partial S_{\text{III}}^{(n+1)}}{\partial j_n} \right). \end{aligned} \quad (6)$$

Thus, it is sufficient to consider the single slab action $S_{\text{III}}^{(n)}$, given in Eq. (3), and its derivatives. An explicit form of these equations is given in Appendix ??.

A priori, the number of cubes in each spacelike slice, $\mathcal{V}_S^3$, enters the total Regge action only as a multiplicative factor due to the symmetry restriction. Thus, in the following two sections, we consider $\mathcal{V}_S = 1$ for simplicity. However, let us mention here that in Sec. IV C, the parameter $\mathcal{V}_S$ turns out to be crucial to obtain the correct continuum limit. We tackle the problem of finding solutions to Eq. (5) first analytically in the subsequent section and develop numerical methods thereafter.

Add explicit equations to the appendix, mention simple dependence, additivity

A. Analytical considerations

For vanishing cosmological constant, $\Lambda = 0$, and stationary boundary data, $j_0 = j_{\mathcal{V}_T}$, the results for a single

³ The construction of the spatially spherical Regge action differs from the construction here, in particular because there exist only finitely many homogeneous triangulations of the 3-sphere, see [13, 26] for details.

time step obtained in the previous section straightforwardly generalize. That is, if $\Lambda = 0$, the Regge equations are satisfied if and only if the boundary data satisfies $j_0 = j_{\mathcal{V}_T}$. Then, the bulk areas $j_n = j_0$ for all $n \in \{1, \dots, \mathcal{V}_T - 1\}$, and the trapezoid areas are undetermined, marking the restoration of time reparametrization invariance of flat space. On the other hand, for $\Lambda \neq 0$ and boundary data $j_0 = j_{\mathcal{V}_T}$, no non-degenerate solution to the Regge equations is found. A unique solution exist for $j_0 \neq j_{\mathcal{V}_T}$ when $\Lambda > 0$ which is in accordance with the continuum Friedmann equations. To summarize, a spatially flat vacuum universe is flat and stationary while a positive cosmological will necessarily lead to curved non-stationary solutions. For the remainder, we assume $\Lambda > 0$ and $j_0 \neq j_{\mathcal{V}_T}$.

Eq. (5) has been analyzed further (coupling a scalar field rather than a cosmological constant) in [11]. Therein, the length variables (s_n, s_{n+1}, H_n) have been employed in which the monotonicity of solutions and the continuum limit are derived most easily. The results of [11] transferred to the present setting with $\Lambda \neq 0$ are briefly summarized in the following.

Combining the Regge equations

$$\frac{\partial S_{\text{III}}}{\partial H_{n-1}} = 0, \quad \frac{\partial S_{\text{III}}}{\partial H_n} = 0, \quad \frac{\partial S_{\text{III}}}{\partial s_n} = 0, \quad (7)$$

one can straightforwardly derive the relation [11]

$$\frac{s_n - s_{n-1}}{H_{n-1}} = \frac{s_{n+1} - s_n}{H_n}, \quad (8)$$

for all $n \in \{1, \dots, \mathcal{V}_T - 1\}$. That is, the difference of spatial edge length with respect to the height of the 4-frusta remains constant. Introducing

$$\tau_0 = \sum_{n=0}^{\mathcal{V}_T-1} H_n, \quad (9)$$

which we will later identify as the discrete elapsed proper time, one can show that Eq. (8) implies

$$\frac{s_{n+1} - s_n}{H_n} = \frac{s_{\mathcal{V}_T} - s_0}{\tau_0}, \quad (10)$$

for all n . Since $H_n, \tau_0 > 0$, this implies $\text{sgn}(s_{n+1} - s_n) = \text{sgn}(s_{\mathcal{V}_T} - s_0)$, that is, solutions evolve monotonically and the boundary data $s_0, s_{\mathcal{V}_T}$ selects whether the universe is expanding, $s_{\mathcal{V}_T} > s_0$, or contracting, $s_{\mathcal{V}_T} < s_0$.

Following the procedure outlined in [11], an analytical continuum limit of the Regge equations can be taken in two steps. First, one performs a linearization of $\frac{\partial S_{\text{III}}}{\partial H_n} = 0$ in the variable $\eta_n = \frac{s_{n+1} - s_n}{H_n}$. In this approximation, corresponding to small deficit angles, one finds

$$\frac{(s_n - s_{n+1})^2}{s_n^2 + s_{n+1}^2} = H_n^2 \frac{\Lambda}{6}. \quad (11)$$

Then, the discrete variables (s_n, s_{n+1}, H_n) need to be identified with the continuum scale factor $a(t)$ and the

lapse function $N(t)$ parametrized in some coordinate time t . In the continuum limit, the height H_n corresponds to the infinitesimal proper time, $H_n \rightarrow N dt$. The discrete length on a given slice is mapped to the scale factor, $s_n \rightarrow a(t)$, while the length of a neighboring edge attains an additional time derivative term, $s_{n+1} \rightarrow a(t) + \dot{a} dt$. Then, at leading order in dt , one finds indeed that Eq. (11) approaches

$$\left(\frac{\dot{a}}{a}\right)^2 = N^2 \frac{\Lambda}{3}, \quad (12)$$

which is the Friedmann equation for a spatially flat universe with cosmological constant. Note that configurations with spacelike trapezoids or spacelike 3-frusta satisfy $\eta_n^2 > 2$ and therefore bear no straightforward continuum limit [11].

The cosmological Regge equations have been shown to capture the Friedmann equations in a suitable continuum limit in previous works [10, 11, 13, 27]. Still, Eq. (5) remain to be solved explicitly in the discrete for many time steps $\mathcal{V}_T$. This represents however an important step since the solutions serve as a classical and discrete benchmark for identifying a semi-classical continuum regime in discrete path integral approaches such as effective spin-foams. In the following section, we tackle this issue employing numerical methods.

B. Numerical implementation

Solving the Regge equations explicitly requires numerical methods. In this work, we employ the package `NLSolve`⁴ in `Julia`, designed for solving non-linear equations. Note that while Eqs. (5) define a real-valued function in real variables, `NLSolve` can also be applied to complex numbers **which may be relevant for the complex critical point method** []. The algorithm we developed for this work specifically tailored to Eq. (5) amounts to the `solve_eom`⁵ method which we explain in the following.

First, note that the method `nsolve` of the `NSolve` package takes as input a vector valued function $\vec{F}$, a vector of initial variables $\vec{x}_0$, and parameters for the number of iterations as well as the error tolerance for solutions. If the problem is sufficiently well-behaved, `nsolve` finds a solution to the equation $\vec{F}(\vec{x}) = \vec{0}$ using $\vec{x}_0$ as a starting guess of the solving algorithm. In our case,

⁴ A documentation of the package can be found at github.com/JuliaNLSolvers/NLSolve.jl. Note that also the `NSolve` method in `Mathematica` can be used, but is expected have lower performance in terms of accuracy and computation time.

⁵ See this github repository for the source code.

$\vec{F} : \mathbb{R}^{2\mathcal{V}_T-1} \rightarrow \mathbb{R}^{2\mathcal{V}_T-1}$, with

$$\vec{F} = \left(\frac{\partial S_{\text{III}}}{\partial k_0}, \frac{\partial S_{\text{III}}}{\partial j_1}, \dots, \frac{\partial S_{\text{III}}}{\partial j_{\mathcal{V}_T-1}}, \frac{\partial S_{\text{III}}}{\partial k_{\mathcal{V}_T-1}} \right), \quad (13)$$

$$\vec{x} = (k_0, j_1, \dots, j_{\mathcal{V}_T-1}, k_{\mathcal{V}_T-1}). \quad (14)$$

Therefore, adapting Eq. (5) to be solved by `nsolve` is done in two steps: 1) set up the function $\vec{F}$ for fixed boundary areas $j_0, j_{\mathcal{V}_T}$ and cosmological constant Λ , and 2) propose a starting guess $\vec{x}_0$. While step 1) is straightforward, step 2) requires particular care because the starting guess $\vec{x}_0$ is crucial for convergence and good performance of the solving algorithm. In particular, as $\mathcal{V}_T$ grows, the number of variables and equation grows and the algorithm because more sensitive to a well-chosen $\vec{x}_0$.

By trial and error, one can find sufficient starting guesses for `nsolve` to converge given various small $\mathcal{V}_T$.⁶ For the solutions $\vec{x}^* = (k_0^*, j_1^*, \dots, j_{\mathcal{V}_T-1}^*, k_{\mathcal{V}_T-1}^*)$, the following patterns are being observed. First, the j_n^* can be approximated as

$$j_n^* \approx j_0 + n\Delta j, \quad n \in \{1, \dots, \mathcal{V}_T - 1\}, \quad (15)$$

with $\Delta j = (j_{\mathcal{V}_T} - j_0)/\mathcal{V}_T$. Second, let k_0^* be the solution of the trapezoid area for $\mathcal{V}_T = 1$. Then, for $\mathcal{V}_T > 1$, the k_n^* are approximated by

$$k_n^* \approx k_0^*/\mathcal{V}_T, \quad n \in \{0, \dots, \mathcal{V}_T - 1\}. \quad (16)$$

That is, to find a good initial guess for the trapezoid areas, one only needs to find a solution k_0^* for a single time step. Crucially, for a single equation, `nsolve` converges for a broad range of initial guesses, so k_0^* can be easily found. In summary, finding a good initial guess $\vec{x}_0$ amounts to finding the one time step solution k_0^* and then setting

$$\vec{x}_0 = \left(\frac{k_0^*}{\mathcal{V}_T}, j_0 + \Delta j, \dots, j_0 + (\mathcal{V}_T - 1)\Delta j, \frac{k_0^*}{\mathcal{V}_T} \right). \quad (17)$$

solver summary

To understand the stability of the solver under perturbations of the initial guess $\vec{x}_0$, consider a vector $\vec{\delta} = \frac{1}{\sqrt{2\mathcal{V}_T-1}}(\delta, \dots, \delta)$. Applying the solver with $\vec{x}_0 + \vec{\delta}$ as initial guess one finds the following behavior:

- For $\mathcal{V}_T = 1$, the solver yields consistent results as long as the entries of $\vec{x}_0 + \vec{\delta}$ are positive. For negative entries, there are non-trivial sign changes in the Regge equation, and in this case, it not clear if solutions even exist. Choosing δ such that the entries are positive, the solutions are stable even for large $\delta > 100$. Also, the number of iterations does




FIG. 2: Fit with $\hat{\beta} = 4.23$ and $\hat{\alpha} = 6.67$. $t(V) = 10^\alpha V^\beta$.

only increase slightly. This is important since setting up $\vec{x}_0$ requires finding k_0^* first. If this first step would be extremely sensitive, the proposal above would not be viable.

- For $\mathcal{V}_T > 1$, the number of iterations needed to find convergence is generically larger, even for $\delta = 0$. Choosing δ such that the entries of $\vec{x}_0 + \vec{\delta}$ are positive, one finds that solutions are stable for small $|\delta|$ ($|\delta| \lesssim 10$ for $\mathcal{V}_T = 2$) with little fluctuations in the required iterations. However, for larger δ , the solutions suddenly jump to unreasonably large or to negative values and the iterations also increase slightly. This effect increases with $\mathcal{V}_T$. For the solver, this means that 1) one should stick with the proposal of $\vec{x}_0$ given above, and 2) for very large $\mathcal{V}_T$, one needs to come up with a refined proposal of initial guesses as $\vec{x}_0$. In this work, the values we consider for $\mathcal{V}_T$ are small enough for $\vec{x}_0$ to yield satisfactory results.

, one observes that for

- output
- convergence monitor
- scaling of computation time as function of $\mathcal{V}$

C. Continuum limit

Facilitated by the numerical tools developed above, we study in this section the continuum limit of Regge calculus at the level of the solutions, rather than at the level of the equations. Quantifying whether the discrete solutions provide a satisfactory approximation to the continuum solutions requires two ingredients: 1) For comparison, one needs to consider appropriate continuum quantities which serve as limiting points of the discrete solutions. 2) One needs to identify the right parameters to tune. Here, the discreteness is parametrized by the number of slices, $\mathcal{V}_T$, as well as the number of 3-cubes per

⁶ We tried using the continuum solution to find a good initial guess. However, we found that the prescription given here ensures fast convergence more reliably.

slice, $\mathcal{V}_S^3$. Thus, the continuum is expected to be reached by suitably scaling $\mathcal{V}_T, \mathcal{V}_S \rightarrow \infty$.

1) *Continuum quantities.* One difficulty in comparing discrete and continuum solutions is that in the discrete, the trapezoid areas are dynamical while in the continuum, the lapse function N is a gauge variable, and thus, any time parametrization corresponds to a gauge choice. That is, when performing the limit, which time parametrization in the continuum emerges? Clearly, the labels $n \in \{0, \dots, \mathcal{V}_T\}$ do not serve as a time as this would imply constant distance between any two neighboring spacelike slices. Instead, recall that the height of 4-frusta was mapped to infinitesimal proper time in the analytical continuum limit of Sec. ???. Therefore, the sum of heights, $\mathcal{T} = \sum_n H_n$, serves as a discrete approximation of the continuum proper time elapsed, defined as $\int dt N(t)$. The advantage of considering proper time is that it is gauge invariant. Thus, one does not run into the issue of having to pick a parametrization in the continuum to compare to the discrete solutions. Using proper time, we will then compare the evolution of the square area j_n^* as a function of the accumulated height $\sum_{m=0}^n H_m$ with the squared scale factor $A = a^2$ as a function of proper time.

2) *The right scaling.* A priori, the parameter $\mathcal{V}_S$ enters the Regge action only as a multiplicative factor. Therefore, it might be tempting to simply set $\mathcal{V}_S = 1$, as done in the analytical part in Sec. ???. However, Fig. ?? demonstrates that leaving $\mathcal{V}_S = 1$ while increasing $\mathcal{V}_T$, the accumulated height $\tau = \sum H$ does not converge to the value of the continuum proper time τ_0 . To obtain the correct τ_0 , one therefore needs to tune $\mathcal{V}_S$ and $\mathcal{V}_T$ simultaneously. Here, a second subtlety needs to be noticed. The continuum proper time only depends on Λ and the initial and final squared scale factor A_0, A_1 . Setting $j_0 = A_0$ and $j_{\mathcal{V}_T} = A_1$ and increasing $\mathcal{V}_S$ i

$\mathcal{V}_T$ and $\mathcal{V}_S$ can be scaled to larger values individually. In particular, from Eq. (5), one can see that

Continuum limit also in the analytical sense? Consider $s_n \rightarrow s_n/\mathcal{V}_S$.

V. CONCLUSION

Extension to $k = \pm 1$.

Acknowledgements The authors thank Seth Asante, Bianca Dittrich and Steffen Gielen for helpful discussions and Tim Stötzl for support with the numerical implementation. AFJ, JDS and SSSt gratefully acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Grant No 422809950. AFJ and JDS acknowledge support by the DFG under Grant No 406116891 within the Research Training Group RTG 2522/1. AFJ is grateful for the generous financial support by the MCQST via the seed funding Aost 862981-8 granted to Jibril Ben Achour by the DFG under Germany's Excellence Strategy – EXC-2111 – 390814868.

Appendix A: Frusta geometry

In area variables, the geometry of a slab is fully characterized by (j_0, j_1, k) , the square area of the cubes in

slice 0 and 1, respectively, and the trapezoid area k . We provide the formulae of the squared volumes of relevant sub-cells in the following. Here, the square areas j_0, j_1 are assumed to be the absolute values of the signed areas and therefore positive. For the squared trapezoid area, we write $\underline{k}^2$ for the signed squared area and k^2 for its absolute value.

A crucial quantity is the squared height of the 4-frusta,

$$H^2 = \frac{8\underline{k}^2 + (j_0 - j_1)^2}{2(\sqrt{j_0} + \sqrt{j_1})^2}, \quad (\text{A1})$$

due to its direct relation to the proper time in the continuum. Furthermore, it gives a bound $8\underline{k}^2 > -(j_0 - j_1)^2$ for the 4-frusta to be embeddable in Minkowski space. Configurations that violate this bound correspond to Euclidean frusta.

The squared length of struts, which connect neighboring spacelike slices is

$$l^2 = \frac{16\underline{k}^2 - (j_0 - j_1)^2}{4(\sqrt{j_0} + \sqrt{j_1})^2}. \quad (\text{A2})$$

Thus, configurations with $16\underline{k}^2 > (j_0 - j_1)^2$ have timelike struts. If equality holds instead, struts are lightlike. Otherwise, they are spacelike. The squared height of 3-frusta is

$$h^2 = \frac{16\underline{k}^2 + (j_0 - j_1)^2}{4(\sqrt{j_0} + \sqrt{j_1})^2}, \quad (\text{A3})$$

so 3-frusta are timelike if $16\underline{k}^2 > -(j_0 - j_1)^2$, lightlike if equality holds and spacelike otherwise.

The 4-volume enters the Regge action with a cosmological constant and is the discrete analogon of the volume element $d^4x \sqrt{-g}$,

$$V_{(4)}^2 = \frac{(j_0 + j_1)^2}{32} (8\underline{k}^2 + (j_0 - j_1)^2). \quad (\text{A4})$$

In this work, we focus exclusively on the causally regular sector, that is, timelike trapezoids, $\underline{k}^2 > 0$. The two types of deficit angles, those located at trapezoids and those located at squares, are respectively given by

$$\delta^E = \frac{\pi}{2} - \cos^{-1} \left(\frac{(j_n - j_{n+1})^2}{16\underline{k}^2 + (j_n - j_{n+1})^2} \right), \quad (\text{A5})$$

and

$$\delta^L = \sinh^{-1} \left(\frac{j_1 - j_0}{\sqrt{16\underline{k}^2 + (j_n - j_{n+1})^2}} \right). \quad (\text{A6})$$

δ^E is a Euclidean angle since the orthogonal subspace to a timelike trapezoid is $\mathbb{R}^2$. In contrast, δ^L is a Lorentzian angle as the orthogonal subspace to a spacelike square is $\mathbb{R}^{1,1}$.

Let us repeat here the Regge action of the causally regular sector,

$$S_{\text{III}}^{(n)} = 6(j_n - j_{n+1}) \sinh^{-1} \left(\frac{j_{n+1} - j_n}{\sqrt{16k_n^2 + (j_n - j_{n+1})^2}} \right) + 12k_n \left[\frac{\pi}{2} - \cos^{-1} \left(\frac{(j_n - j_{n+1})^2}{16k_n^2 + (j_n - j_{n+1})^2} \right) \right] - \Lambda V^{(4)}. \quad (\text{A7})$$

The equations of motion are given by

$$\frac{\partial S_{\text{III}}}{\partial k_n} = 0, \quad \frac{\partial S_{\text{III}}}{\partial j_n} = 0, \quad (\text{A8})$$

which can be simplified using the Schläfli identity and

the additivity of the action. That is, the above equations can be reformulated as

$$\frac{\partial S_{\text{III}}^{(n)}}{\partial k_n} = 0, \quad \frac{\partial S_{\text{III}}^{(n-1)}}{\partial j_n} + \frac{\partial S_{\text{III}}^{(n)}}{\partial j_n} = 0, \quad (\text{A9})$$

with the derivative terms explicitly given by

$$\begin{aligned} \frac{\partial S_{\text{III}}^{(n)}}{\partial k_n} &= 12 \left[\frac{\pi}{2} - \cos^{-1} \left(\frac{(j_n - j_{n+1})^2}{16k_n^2 + (j_n - j_{n+1})^2} \right) \right] - \Lambda \frac{(j_n + j_{n+1})^2 k_n}{4V_{(4)}}, \\ \frac{\partial S_{\text{III}}^{(n)}}{\partial j_n} &= 6 \sinh^{-1} \left(\frac{j_{n+1} - j_n}{\sqrt{16k_n^2 + (j_n - j_{n+1})^2}} \right) - \Lambda \frac{j_n(j_n - j_{n+1}) + 4k_n^2}{2\sqrt{(j_n - j_{n+1})^2 + 8k_n^2}}. \end{aligned} \quad (\text{A10})$$

Appendix B: Cosmology in the continuum

The line element of a spatially flat, homogeneous and isotropic universe is given by

$$ds^2 = N^2(t) dt^2 - a^2(t) d\mathbf{x}^2, \quad (\text{B1})$$

with $a(t)$ the scale factor and $d\mathbf{x}^2$ the 3-dimensional Euclidean line element. The lapse function, $N(t)$, corresponds to a choice of gauge emerging from the time reparametrization invariance of the system.

Dynamics of the cosmological system are governed by the Friedmann equations which, in the case where matter only consists of a cosmological constant, are simply given by

$$\left(\frac{\dot{a}}{a} \right)^2 = N^2 \frac{\Lambda}{3}. \quad (\text{B2})$$

This is straightforwardly solved by

$$a(t) = a_0 \exp \left(\pm \frac{\Lambda}{3} \int_{t_0}^t dt' N(t') \right), \quad (\text{B3})$$

where we identify the gauge-invariant elapsed proper time as

$$\tau^{\text{cont}}(t_0, t) = \int_{t_0}^t dt' N(t'). \quad (\text{B4})$$

The proper time plays an important role in the comparison between discrete and continuum equations as the cumulative height of frusta converges to the proper time in an appropriate continuum limit.

In the discrete, as well as for effective spin-foams in ensuing studies, one is not given an initial scale factor and an initial time but rather an initial and final scale factor (corresponding to $\sqrt{j_0}$ and $\sqrt{j_{V_T}}$, respectively). If, in the continuum, a_0, a_1 and Λ are given, the proper time is fixed to

$$\tau_0^{\text{cont}} = \log \left(\frac{a_1}{a_0} \right) \sqrt{\frac{\Lambda}{3}}. \quad (\text{B5})$$

This quantity serves as a limiting value of $\tau = \sum_n H_n$ in the continuum limit described in Sec. ??.

-
- [1] T.E. Regge, *General relativity without coordinates*, *Nuovo Cimento* **19** (1961) 558.
 - [2] R.D. Sorkin, *Lorentzian angles and trigonometry including lightlike vectors*, 1908.10022.
 - [3] S.K. Asante, B. Dittrich and J. Padua-Argüelles, *Effective spin foam models for Lorentzian quantum gravity*, *Class. Quant. Grav.* **38** (2021) 195002 [2104.00485].
 - [4] S.K. Asante, B. Dittrich and J. Padua-Argüelles,

Complex actions and causality violations: applications to Lorentzian quantum cosmology, *Class. Quant. Grav.* **40** (2023) 105005 [2112.15387].

- [5] J.W. Barrett, *First order Regge calculus*, *Class. Quant. Grav.* **11** (1994) 2723 [hep-th/9404124].
- [6] B. Dittrich and S. Speziale, *Area-angle variables for general relativity*, *New J. Phys.* **10** (2008) 083006 [0802.0864].
- [7] S.K. Asante, B. Dittrich and H.M. Haggard, *The*

- Degrees of Freedom of Area Regge Calculus: Dynamics, Non-metricity, and Broken Diffeomorphisms*, *Class. Quant. Grav.* **35** (2018) 135009 [1802.09551].
- [8] J. Makela and R.M. Williams, *Constraints on area variables in Regge calculus*, *Class. Quant. Grav.* **18** (2001) L43 [gr-qc/0011006].
- [9] J.W. Barrett, M. Rocek and R.M. Williams, *A Note on area variables in Regge calculus*, *Class. Quant. Grav.* **16** (1999) 1373 [gr-qc/9710056].
- [10] B. Bahr, S. Klöser and G. Rabuffo, *Towards a Cosmological subsector of Spin Foam Quantum Gravity*, *Phys. Rev. D* **96** (2017) 086009 [1704.03691].
- [11] A.F. Jercher and S. Steinhaus, *Cosmology in Lorentzian Regge calculus: causality violations, massless scalar field and discrete dynamics*, *Class. Quant. Grav.* **41** (2024) 105008 [2312.11639].
- [12] A. Jercher and S. Steinhaus, *Transfer matrix for effective spin-foams (wip)*, **25x.xxxxx**.
- [13] B. Dittrich, S. Gielen and S. Schander, *Lorentzian quantum cosmology goes simplicial*, *Class. Quant. Grav.* **39** (2022) 035012 [2109.00875].
- [14] B. Dittrich and J. Padua-Argüelles, *Lorentzian Quantum Cosmology from Effective Spin Foams*, *Universe* **10** (2024) 296 [2306.06012].
- [15] A.F. Jercher, J.D. Simão and S. Steinhaus, *Partial absence of cosine problem in 3d Lorentzian spin foams*, **2404.16943**.
- [16] A.F. Jercher, J.D. Simão and S. Steinhaus, (2 + 1) *Lorentzian quantum cosmology from spin-foams: opportunities and obstacles for semi-classicality*, *Class. Quant. Grav.* **42** (2025) 085015 [2411.08109].
- [17] G.T. Horowitz, *Topology change in classical and quantum gravity*, *Class. Quant. Grav.* **8** (1991) 587.
- [18] J. Louko and R.D. Sorkin, *Complex actions in two-dimensional topology change*, *Class. Quant. Grav.* **14** (1997) 179 [gr-qc/9511023].
- [19] H.F. Dowker, R.S. Garcia and S. Surya, *Morse index and causal continuity: A Criterion for topology change in quantum gravity*, *Class. Quant. Grav.* **17** (2000) 697 [gr-qc/9910034].
- [20] M. Rocek and R.M. Williams, *QUANTUM REGGE CALCULUS*, *Phys. Lett. B* **104** (1981) 31.
- [21] M. Rocek and R.M. Williams, *The Quantization of Regge Calculus*, *Z. Phys. C* **21** (1984) 371.
- [22] B. Dittrich, *Diffeomorphism Symmetry in Quantum Gravity Models*, *Adv. Sci. Lett.* **2** (2009) 151 [0810.3594].
- [23] B. Bahr and B. Dittrich, *Improved and Perfect Actions in Discrete Gravity*, *Phys. Rev. D* **80** (2009) 124030 [0907.4323].
- [24] B. Bahr and S. Steinhaus, *Investigation of the Spinfoam Path integral with Quantum Cuboid Intertwiners*, *Phys. Rev. D* **93** (2016) 104029 [1508.07961].
- [25] F. Hilpert, *Simplicial cosmology of a closed vacuum universe with positive cosmological constant in riemannian and lorentzian regge calculus*, Master's thesis, Friedrich-Schiller-Universität Jena, 2024.
- [26] P. Collins and R.M. Williams, *Dynamics of the Friedmann Universe Using Regge Calculus*, *Phys. Rev. D* **7** (1973) 965.
- [27] L. Brewin, *Friedmann cosmologies via the regge calculus*, *Classical and Quantum Gravity* **4** (1999) 899.