

Title

Alexander F. Jercher^{1,2,*} and Sebastian Steinhaus^{1,†}

¹*Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena,
Max-Wien-Platz 1, 07743 Jena, Germany, EU*

²*corresponding author*

(Dated: October 13, 2025)

I. INTRODUCTION

$8\pi G_N = 1?$, spacetime signature $(+, -, -, -)$

II. SETUP

A. Lorentzian Regge calculus

Regge calculus [1] is a discrete gravitational theory on simplicial manifolds Δ with the length of edges $\{l_e\}$ dynamical. In a Lorentzian setting, a simplicial manifold consists of intrinsically flat Lorentzian 4-simplices $\sigma \in \Delta$. Curvature is located at triangles $t \in \Delta$ and captured by deficit angles $\delta_t(\{l_e\})$, lying in the space orthogonal to t . For t spacelike (timelike), this space is isomorphic to $\mathbb{R}^{1,1}$ ($\mathbb{R}^2$).¹ Lorentzian deficit angles are defined as $\delta_t^{L,\pm} = \mp i2\pi - \sum_{\sigma \supset t} \psi_{\sigma,t}^{L,\pm}$, following the conventions of [2, 3], where the $\psi_{\sigma,t}$ are dihedral angles. “ $\pm$ ” indicates a choice of sign which becomes important for causally irregular configurations [2–4] discussed below. The bulk action of Lorentzian Regge calculus is given by [1–3]

$$S_R[\{l_e\}] = \sum_{t \in \Delta: \text{tl}} |A_t(\{l_e\})| \delta_t^E(\{l_e\}) + \sum_{t \in \Delta: \text{sl}} |A_t(\{l_e\})| \delta_t^{L,\pm}(\{l_e\}), \quad (1)$$

with $|A_t|$ the absolute value of the triangle area. In the presence of a boundary, $\partial\Delta \neq \emptyset$, S_R attains boundary terms such that overall additivity is ensured. The equations of motion are given by $\frac{\partial S_R}{\partial l_e} = 0$ supplemented by the Schläfli identity, $\sum_t |A_t| \frac{\partial \delta_t}{\partial l_e} = 0$.

Regge calculus can be formulated in terms of different variables [5, 6], with area variables [7–9] playing a particularly important role here due to their connection to LQG and spin-foams. The relation between area and length variables is in general not invertible, and area variables need to be further constrained to yield the dynamical equations of length Regge calculus [3, 6, 9]. However,




FIG. 1: Boundary of a 4-frustum given by two 3-cubes and six equal 3-frusta.

in the symmetry reduced setting introduced below, the relation of area and length is globally invertible and the aforementioned constraints trivialize. For the remainder, we study the classical dynamics of area Regge calculus which will be used in future work as a comparison to results of the effective spin-foam path integral.

B. Lorentzian 4-frusta

Lorentzian 4-dimensional frusta are ideal to capture spatially flat, homogeneous and isotropic discrete geometries [10, 11]. The boundary of 4-frusta, depicted in Fig. 1, contains two spacelike 3-cubes and six 3-frusta that are either spacelike or timelike. Its entire geometry is captured by the area of squares contained in the 3-cubes, j_0 and j_1 , and the area of the trapezoids contained in the 3-frusta, k . The general notions of Regge calculus introduced above are straightforwardly adapted to this setting. Gluing 4-frusta along boundary 3-frusta in spatial direction and along boundary 3-cubes in temporal direction, one obtains an extended cosmological spacetime of hypercubical combinatorics $\mathcal{X}_V^{(4)}$ with slices labelled by $n \in \{0, \dots, \mathcal{V}_T\}$ discretized by $\mathcal{V}_S^3$ spacelike 3-cubes. We say that a 4-frustum lies in a “slab” between slices n and $n + 1$.

The squared volume of subcells connecting neighboring slices, i.e. 3-frusta, trapezoids and edges, which are detailed in Appendix ??, carries information on the causal structure. If it is positive (negative), the corresponding subcell is timelike (spacelike). In the symmetry reduced setting, the ratio $\frac{|j_n - j_{n+1}|}{4k_n}$ fully characterizes the causal character of subcells suggesting separating the 4-frusta configurations into three sectors summarized in Tab. ??.

* alexander.jercher@uni-jena.de

† sebastian.steinhaus@uni-jena.de

¹ For the remainder, we neglect lightlike edges, triangles and tetrahedra. See [2] for the definition of deficit angles associated to $(d - 1)$ -dimensional null cells.

As discussed in [11–15], two of these sectors I and II exhibit an irregular light cone structure, also referred to as causality violations, located at spacelike sub-cells. Examples of such irregularities are given by the trouser singularity and the yarmulke singularity associated to spatial topology change [16–18]. For causality violations at $(d - 2)$ -dimensional spacelike hinges the imaginary parts of the dihedral angles do not sum up to $\mp i2\pi$, leading to a complex valued Regge action. Choosing “+” in the convention of Lorentzian angles, the amplitude e^{iS_R} is exponentially suppressed for trouser and exponentially enhanced for yarmulke configurations, vice versa for “−” [3]. Without a complexification of the vari-

ables, the Regge equations of motion for causality violating configurations are complex equations with real variables and therefore overconstrained. Moreover, the results of [11, 12] show that in the respective settings, no real solutions exist. Since this applies also to the setting used in this work we henceforth focus on the causally regular sector III.

C. Lorentzian Regge action for spatially flat cosmology

For a single 4-frustum between, say slice n and $n + 1$, the causally regular Regge action is given by

$$S_{\text{III}}^{(n)} = 6(j_n - j_{n+1}) \sinh^{-1} \left(\frac{j_{n+1} - j_n}{\sqrt{16k_n^2 + (j_n - j_{n+1})^2}} \right) + 12k_n \left[\frac{\pi}{2} - \cos^{-1} \left(\frac{(j_n - j_{n+1})^2}{16k_n^2 + (j_n - j_{n+1})^2} \right) \right] - \Lambda V^{(4)}, \quad (2)$$

where Λ is a cosmological constant and $V^{(4)}$ is the 4-volume defined in Eq. (??) as an equation of (j_n, j_{n+1}, k_n) . By construction, the 4-frustum action is additive such that on a discretization $\mathcal{X}_V^{(4)}$ with $\mathcal{V}_S^3$ vertices in spatial direction and $\mathcal{V}_T + 1$ spatial slices, the action is given by $S_R = \mathcal{V}_S^3 \sum_{n=0}^{\mathcal{V}_T-1} S_R^{(n)}$. Additivity of the action per slab implies that the equations of motion for a given geometric quantity, being either j_n or k_n , will only depend on neighboring geometric labels.

In the following, we study the discrete dynamics of spatially flat cosmology as governed by the Regge equations. To that end, we consider first the case of a single time step, $\mathcal{V}_T = 1$, in Sec. III and generalize then to an arbitrary number of time steps in Sec. IV.

III. RECAP: ONE TIME STEP

The classical and quantum dynamics of a single timestep in discrete Lorentzian cosmology have been studied extensively in [4, 12, 13] for spatial curvature with cosmological constant and in [11, 15] for spatially flat spacetimes with a free scalar field. In this chapter we extend these studies to spatially flat cosmology with a cosmological constant, before then focussing on the case of arbitrarily many time steps.

For $\mathcal{V}_T = 1$, the square areas j_0, j_1 are boundary data and only the trapezoid area k_0 is dynamical. Thus, the Regge equations are simply given by

$$\frac{\partial S_{\text{III}}}{\partial k_0} = 0, \quad (3)$$

where due to the Schläfli identity, the first term in Eq. (2) drops out. Notice that this equation is transcendental and can therefore only be tackled numerically.

Vanishing cosmological constant. In the case of $\Lambda = 0$, the Regge action describes a spatially flat vacuum universe. If the boundary data is non-stationary, i.e. $j_0 \neq j_1$, Eq. (3) exhibits no solution for k_0 . In other words, a classical universe can neither expand nor contract if there is no matter present. This is in accordance with the continuum Friedmann equations in vacuum, $(\frac{\dot{a}}{a})^2 = 0$, with a the scale factor, which are solved only by $a = \text{const}$. On the other hand, if $j_0 = j_1$, Eq. (3) is trivially satisfied and the trapezoid area k_0 is undetermined. In fact, the Regge action vanishes, $S_{\text{III}} = 0$ for all k_0 , and the 4-frusta reduce to 4-cuboids discretizing flat space. The arbitrariness of k is a common phenomenon for flat solutions in Regge calculus [19, 20] and signifies the restoration of diffeomorphism symmetry in the discrete [21, 22]. This symmetry amounts to moving spacelike slices (instead of individual vertices), see also the discussion in [23]. Note that for 4-cuboids, $k = \sqrt{j}H$ with H the height of the cuboid which is directly related to the lapse function N in the continuum [11, 12]. The lapse is the gauge variable of the residual diffeomorphism symmetry after symmetry reduction, encoding time reparametrization invariance.

Non-vanishing cosmological constant. For a non-vanishing cosmological constant, $\Lambda \neq 0$, and stationary boundary conditions, $j_0 = j_1$, the only solution to Eq. (3) is $k_0 = 0$, corresponding to degenerate 4-frusta with vanishing height. Thus, a stationary universe for $\Lambda \neq 0$ only exists if no time passes, i.e. if there is no actual time evolution. The more interesting case is $j_0 \neq j_1$ where a non-trivial solution k_0^* for Eq. (3) exists. More precisely, a unique solution k_0^* is found for any $\Lambda > 0$ and any $j_0 \neq j_1$. The on-shell trapezoid area $k_0^*(\Lambda)$ as a function of Λ behaves as a monomial decay $k_0^* \sim c\Lambda^\alpha$ with $c, \alpha > 0$ constants depending on j_0, j_1 . For $\Lambda \neq 0$, the solution diverges and ceases to exist at $\Lambda = 0$ in lines with the discussion above. In the limit $\Lambda \rightarrow \infty$, Eq. (3)

is dominated by the Λ -term, i.e.

$$\Lambda \frac{(j_0 + j_1)^2 k}{4V^{(4)}} \approx 0, \quad (4)$$

solved by $k_0^* = 0$.

Note that the restriction on the sign of Λ is in accordance with the continuum Friedmann equations, $(\frac{\dot{a}}{a})^2 = N^2 \frac{\Lambda}{3}$, where real solutions for the scale factor require $\Lambda > 0$.

In the spatially spherical case in Lorentzian (Euclidean) signature, where **scale factor is bounded from below (above), ... add contrasting info here.**

IV. ARBITRARY NUMBER OF TIME STEPS

In this section, we study the Regge equations of spatially flat Lorentzian cosmology for an arbitrary number of time steps $\mathcal{V}_T$. This amounts to solving $2\mathcal{V}_T - 1$ coupled transcendental equations for the bulk variables $k_0, j_1, \dots, j_{\mathcal{V}_T-1}, k_{\mathcal{V}_T-1}$ for given boundary data $j_0, j_{\mathcal{V}_T}$ and cosmological constant Λ ,

$$\frac{\partial S_{\text{III}}}{\partial k_n} = 0, \quad \frac{\partial S_{\text{III}}}{\partial j_m} = 0, \quad (5)$$

with $n \in \{0, \dots, \mathcal{V}_T - 1\}$ and $m \in \{1, \dots, \mathcal{V}_T - 1\}$. A priori, the number of cubes in each spacelike slice, $\mathcal{V}_S^3$, enters the total Regge action only as a multiplicative factor due to the symmetry restriction. Thus, in the following two sections, we consider $\mathcal{V}_S = 1$ for simplicity. However, let us mention here that in Sec. IV C, the parameter $\mathcal{V}_S$ turns out to be crucial to obtain the correct continuum limit. We tackle the problem of finding solutions to Eq. (5) first analytically in the subsequent section and develop numerical methods thereafter.

A. Analytical considerations

For vanishing cosmological constant, $\Lambda = 0$, and stationary boundary data, $j_0 = j_{\mathcal{V}_T}$, the results for a single time step obtained in the previous section straightforwardly generalize. That is, if $\Lambda = 0$, the Regge equations are satisfied if and only if the boundary data satisfies $j_0 = j_{\mathcal{V}_T}$. Then, the bulk areas $j_n = j_0$ for all $n \in \{1, \dots, \mathcal{V}_T - 1\}$, and the trapezoid areas are undetermined, marking the restoration of time reparametrization invariance of flat space. On the other hand, for $\Lambda \neq 0$ and boundary data $j_0 = j_{\mathcal{V}_T}$, no non-degenerate solution to the Regge equations is found. A unique solution exist for $j_0 \neq j_{\mathcal{V}_T}$ when $\Lambda > 0$ which is in accordance with the continuum Friedmann equations. To summarize, a spatially flat vacuum universe is flat and stationary while a positive cosmological will necessarily lead to curved non-stationary solutions. For the remainder, we assume $\Lambda > 0$ and $j_0 \neq j_{\mathcal{V}_T}$.

Refer to [11] and explain the monotonicity in the height variable picture.

- monotonicity
- analytical continuum limit

B. Numerical implementation

Solving the Regge equations explicitly requires numerical methods. In this work, we employ the package `NLSolve`² in `Julia`, designed for solving non-linear equations of the form $\vec{F}(\vec{x}) = \vec{0}$. Note that while Eqs. (5) define a function $\vec{F} : \mathbb{R}^{2\mathcal{V}_T-1} \rightarrow \mathbb{R}^{2\mathcal{V}_T-1}$, `NLSolve` can also be applied to complex numbers **which may be relevant for the complex critical point method** [1]. The algorithm we developed for this work specifically tailored to Eq. (5) comprises the `solve_eom`³ method which we explain in the following.

- input, output variables
- generating initial guess
- ignore $\mathcal{V}_S$ for the moment

To guarantee convergence and good performance of the solving algorithm, a good initial guess for the solutions needs to be provided. To that end,

C. Continuum limit

blabla

V. CONCLUSION

Acknowledgements The authors thank Seth Asante, Bianca Dittrich and Steffen Gielen for helpful discussions and Tim Stötzel for support with the numerical implementation. AFJ, JDS and SSt gratefully acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Grant No 422809950. AFJ and JDS acknowledge support by the DFG under Grant No 406116891 within the Research Training Group RTG 2522/1. AFJ is grateful for the generous financial support by the MCQST via the seed funding Aost 862981-8 granted to Jibril Ben Achour by the DFG under Germany's Excellence Strategy – EXC-2111 – 390814868.

² A documentation of the package can be found at github.com/JuliaNLSolvers/NLSolve.jl. Note that also the `NSolve` method in `Mathematica` can be used, but is expected have lower performance in terms of accuracy and computation time.

³ See this github repository for the source code.

Appendix A: Frusta geometry

Appendix B: Series convergence acceleration

Before we explicitly address the numerical evaluation of the partition function, let us digress briefly to introduce the Shanks transform and Wynn's ϵ -algorithm which have been applied to effective spin-foams⁴ in [13]. In the subsequent sections, we will frequently encounter unbounded sums of the form

$$\mathfrak{S} = \lim_{n \rightarrow \infty} \mathfrak{S}_n = \lim_{n \rightarrow \infty} \sum_{j=1}^n a_j \quad (\text{B1})$$

for some complex sequence $\{a_j\}$. Assuming the partial sum $\mathfrak{S}_n$ to be known and of the form [24]

$$\mathfrak{S}_n = \mathfrak{S} + \sum_{j=0}^{k-1} c_j \lambda_j^n, \quad (\text{B2})$$

with c_j coefficients and the $1 > |\lambda_0| > \dots > |\lambda_{k-1}|$ referred to as transients, there are $2k + 1$ unknowns given

by the limiting value $\mathfrak{S}$, the c_j and the λ_j . In order to solve for $\mathfrak{S}$, one utilizes $2k + 1$ consecutive sequence values $\mathfrak{S}_n, \mathfrak{S}_{n+1}, \dots, \mathfrak{S}_{n+2k}$ to obtain the k -th Shanks transform [26, 27] $e_k(\mathfrak{S}_n)$ as a ratio of determinants. For sufficiently large n , the Shanks transform $e_k(\mathfrak{S}_n)$ approximates $\mathfrak{S}$ faster than taking the limit $n \rightarrow \infty$ of the partial sums $\mathfrak{S}_n$.

In [28], Wynn introduced a non-linear recursive relation to efficiently compute the Shanks transform, commonly referred to as ϵ -algorithm. Given the partial sums $\mathfrak{S}_1, \dots, \mathfrak{S}_n$, define [24]

$$\epsilon_{-1}^{(n)} = 0, \quad \epsilon_0^{(n)} = \mathfrak{S}_n. \quad (\text{B3})$$

Then, ϵ 's of higher k are obtained via the relation

$$\epsilon_{k+1}^{(n)} = \epsilon_{k-1}^{(n+1)} + \frac{1}{\epsilon_k^{(n+1)} - \epsilon_k^{(n)}}. \quad (\text{B4})$$

Wynn has shown [28] that $\epsilon_{2k}^{(n)} = e_k(\mathfrak{S}_n)$ thus providing a fast algorithm for computing the k -th Shanks transform of a sequence $\mathfrak{S}_n$.⁵

Note that Wynn's ϵ -algorithm can be applied to sequences of more than one variable, which we discuss explicitly in Sec. ??.

-
- [1] T.E. Regge, *General relativity without coordinates*, *Nuovo Cimento* **19** (1961) 558.
- [2] R.D. Sorkin, *Lorentzian angles and trigonometry including lightlike vectors*, **1908.10022**.
- [3] S.K. Asante, B. Dittrich and J. Padua-Argüelles, *Effective spin foam models for Lorentzian quantum gravity*, *Class. Quant. Grav.* **38** (2021) 195002 [2104.00485].
- [4] S.K. Asante, B. Dittrich and J. Padua-Argüelles, *Complex actions and causality violations: applications to Lorentzian quantum cosmology*, *Class. Quant. Grav.* **40** (2023) 105005 [2112.15387].
- [5] J.W. Barrett, *First order Regge calculus*, *Class. Quant. Grav.* **11** (1994) 2723 [hep-th/9404124].
- [6] B. Dittrich and S. Speziale, *Area-angle variables for general relativity*, *New J. Phys.* **10** (2008) 083006 [0802.0864].
- [7] S.K. Asante, B. Dittrich and H.M. Haggard, *The Degrees of Freedom of Area Regge Calculus: Dynamics, Non-metricity, and Broken Diffeomorphisms*, *Class. Quant. Grav.* **35** (2018) 135009 [1802.09551].
- [8] J. Makela and R.M. Williams, *Constraints on area variables in Regge calculus*, *Class. Quant. Grav.* **18** (2001) L43 [gr-qc/0011006].
- [9] J.W. Barrett, M. Rocek and R.M. Williams, *A Note on area variables in Regge calculus*, *Class. Quant. Grav.* **16** (1999) 1373 [gr-qc/9710056].
- [10] B. Bahr, S. Klöser and G. Rabuffo, *Towards a Cosmological subsector of Spin Foam Quantum Gravity*, *Phys. Rev. D* **96** (2017) 086009 [1704.03691].
- [11] A.F. Jercher and S. Steinhaus, *Cosmology in Lorentzian Regge calculus: causality violations, massless scalar field and discrete dynamics*, *Class. Quant. Grav.* **41** (2024) 105008 [2312.11639].
- [12] B. Dittrich, S. Gielen and S. Schander, *Lorentzian quantum cosmology goes simplicial*, *Class. Quant. Grav.* **39** (2022) 035012 [2109.00875].
- [13] B. Dittrich and J. Padua-Argüelles, *Lorentzian Quantum Cosmology from Effective Spin Foams*, *Universe* **10** (2024) 296 [2306.06012].
- [14] A.F. Jercher, J.D. Simão and S. Steinhaus, *Partial absence of cosine problem in 3d Lorentzian spin foams*, **2404.16943**.
- [15] A.F. Jercher, J.D. Simão and S. Steinhaus, *(2 + 1) Lorentzian quantum cosmology from spin-foams: opportunities and obstacles for semi-classicality*, *Class. Quant. Grav.* **42** (2025) 085015 [2411.08109].
- [16] G.T. Horowitz, *Topology change in classical and quantum gravity*, *Class. Quant. Grav.* **8** (1991) 587.
- [17] J. Louko and R.D. Sorkin, *Complex actions in two-dimensional topology change*, *Class. Quant. Grav.* **14** (1997) 179 [gr-qc/9511023].
- [18] H.F. Dowker, R.S. Garcia and S. Surya, *Morse index and causal continuity: A Criterion for topology change*

⁴ A closely related series convergence acceleration, known as Aitken's Δ^2 -process [24] has been applied to infinite bulk variable summations in [25].

⁵ A Mathematica algorithm for convergence acceleration via Wynn's method can be found in the repository SequenceLimit. An implementation in JULIA can be found in <https://github.com/Jercheal/3d-cosmology>.

- in quantum gravity, *Class. Quant. Grav.* **17** (2000) 697 [gr-qc/9910034].
- [19] M. Rocek and R.M. Williams, *QUANTUM REGGE CALCULUS*, *Phys. Lett. B* **104** (1981) 31.
- [20] M. Rocek and R.M. Williams, *The Quantization of Regge Calculus*, *Z. Phys. C* **21** (1984) 371.
- [21] B. Dittrich, *Diffeomorphism Symmetry in Quantum Gravity Models*, *Adv. Sci. Lett.* **2** (2009) 151 [0810.3594].
- [22] B. Bahr and B. Dittrich, *Improved and Perfect Actions in Discrete Gravity*, *Phys. Rev. D* **80** (2009) 124030 [0907.4323].
- [23] B. Bahr and S. Steinhaus, *Investigation of the Spinfoam Path integral with Quantum Cuboid Intertwiners*, *Phys. Rev. D* **93** (2016) 104029 [1508.07961].
- [24] E.J. Weniger, *Nonlinear sequence transformations for the acceleration of convergence and the summation of divergent series*, *Computer Physics Reports* **10** (1989) 189 [math/0306302].
- [25] P. Donà and P. Frisoni, *Summing bulk quantum numbers with Monte Carlo in spin foam theories*, *Phys. Rev. D* **107** (2023) 106008 [2302.00072].
- [26] R.J. Schmidt, *Xxxiii. on the numerical solution of linear simultaneous equations by an iterative method*, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **32** (1941) 369.
- [27] D. Shanks, *Non-linear transformations of divergent and slowly convergent sequences*, *Journal of Mathematics and Physics* **34** (1955) 1 [<https://onlinelibrary.wiley.com/doi/pdf/10.1002/sapm195534>].
- [28] P. Wynn, *On a device for computing the $em(sn)$ transformation*, *Mathematical Tables and Other Aids to Computation* **10** (1956) 91.