

INTRO to DATA SCIENCE

SESSION 13 SUPPLEMENT: PROBABILITY

Rob Hall

DAT13 SF // April 20, 2015

I. PROBABILITY SUPPLEMENT

Q: What is a probability?

*Q: What is a **probability**?*

A: A number between 0 and 1 that characterizes the likelihood that some event will occur.

*Q: What is a **probability**?*

A: A number between 0 and 1 that characterizes the likelihood that some event will occur.

The probability of event A is denoted $P(A)$.

Q: What is the set of all possible events called?

Q: What is the set of all possible events called?

*A: This set is called the **sample space** Ω . Event A is a member of the sample space, as is every other event.*

Q: What is the set of all possible events called?

*A: This set is called the **sample space** Ω . Event A is a member of the sample space, as is every other event.*

The probability of the sample space $P(\Omega)$ is 1.

Q: Consider two events A & B . How can we characterize the intersection of these events?

Q: Consider two events A & B . How can we characterize the intersection of these events?

A: With the joint probability of A and B , written $P(AB)$.

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B ?*

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B ?*

A: The intersection of A & B divided by region B .

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B ?*

A: The intersection of A & B divided by region B .

NOTE

This information about B transforms the sample space.

Take a moment to convince yourself of this!

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B ?*

A: The intersection of A & B divided by region B .

*This is called the **conditional probability** of A given B , written $P(A|B) = P(AB) / P(B)$.*

NOTE

This information about B transforms the sample space.

Take a moment to convince yourself of this!

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B ?*

A: The intersection of A & B divided by region B .

*This is called the **conditional probability** of A given B , written $P(A|B) = P(AB) / P(B)$.*

*Notice, with this we can also write $P(AB) = P(A|B) * P(B)$.*

NOTE

This information about B transforms the sample space.

Take a moment to convince yourself of this!

Q: What does it mean for two events to be independent?

*Q: What does it mean for two events to be **independent**?*

A: Information about one does not affect the probability of the other.

*Q: What does it mean for two events to be **independent**?*

A: Information about one does not affect the probability of the other.

This can be written as $P(A|B) = P(A)$.

*Q: What does it mean for two events to be **independent**?*

A: Information about one does not affect the probability of the other.

This can be written as $P(A|B) = P(A)$.

Using the definition of the conditional probability, we can also write:

$$P(A|B) = P(AB) / P(B) = P(A) \rightarrow P(AB) = P(A) * P(B)$$

A motivating example: COOKIES!



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

In other words, we want: $P(\text{Bowl 1} \mid \text{vanilla})$ This is a conditional probability.

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

In other words, we want: $P(\text{Bowl 1} \mid \text{vanilla})$ This is a conditional probability.

How can we compute this?

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

In other words, we want: $P(\text{Bowl 1} \mid \text{vanilla})$

What about $P(\text{vanilla} \mid \text{Bowl 1})$?

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

In other words, we want: $P(\text{Bowl 1} \mid \text{vanilla})$

What about $P(\text{vanilla} \mid \text{Bowl 1})$? That's easy! $P(\text{vanilla} \mid \text{Bowl 1}) = 30/40 = 3/4$

Now suppose you choose one of the bowls at random and, without looking, select a cookie at random. The cookie is vanilla. What is the probability that it came from Bowl 1?



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

In other words, we want: $P(\text{Bowl 1} \mid \text{vanilla})$

But $P(\text{Bowl 1} \mid \text{vanilla})$ is NOT equal to $P(\text{vanilla} \mid \text{Bowl 1}) = 3/4$

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

$$P(AB) = P(A|B) * P(B)$$

from earlier slide

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

$$P(AB) = P(A|B) * P(B)$$

$$P(BA) = P(B|A) * P(A)$$

*from earlier slide
by substitution*

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

$$P(AB) = P(A|B) * P(B)$$

from earlier slide

$$P(BA) = P(B|A) * P(A)$$

by substitution

But $P(AB) = P(BA)$

since event $AB = \text{event } BA$

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

$$P(AB) = P(A|B) * P(B)$$

from earlier slide

$$P(BA) = P(B|A) * P(A)$$

by substitution

But $P(AB) = P(BA)$

since event $AB = \text{event } BA$

*$\rightarrow P(A|B) * P(B) = P(B|A) * P(A)$*

by combining the above

The way we get from $P(\text{Bowl 1} \mid \text{vanilla})$ to $P(\text{vanilla} \mid \text{Bowl 1})$ is as follows:

$$P(AB) = P(A|B) * P(B)$$

from earlier slide

$$P(BA) = P(B|A) * P(A)$$

by substitution

But $P(AB) = P(BA)$

since event $AB = \text{event } BA$

$$\rightarrow P(A|B) * P(B) = P(B|A) * P(A)$$

by combining the above

$$\rightarrow P(A|B) = P(B|A) * P(A) / P(B)$$

by rearranging last step

*This result is called **Bayes' theorem**.*

$$P(A|B) = P(A) * P(B|A) / P(B)$$

We want: $P(\text{Bowl 1} \mid \text{vanilla})$



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

$$P(A|B) = P(A) * P(B|A) / P(B)$$

What is $P(A)$?

What is $P(B)$?

What is $P(B|A)$?

We want: $P(\text{Bowl 1} \mid \text{vanilla})$



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

$$P(A|B) = P(A) * P(B|A) / P(B)$$

$$P(A) = 0.5$$

$$P(B) = 50 / 80 = 5/8$$

$$P(B|A) = 30/40 = 3/4$$

We want: $P(\text{Bowl 1} \mid \text{vanilla})$



*Bowl 1 contains:
30 vanilla cookies
10 chocolate chip cookies*



*Bowl 2 contains:
20 vanilla cookies
20 chocolate chip cookies*

$$P(A|B) = P(A) * P(B|A) / P(B) = 0.5 * 6/8 / 5/8 = \mathbf{3/5}$$

$$P(A) = 0.5$$

$$P(B) = 50 / 80 = 5/8$$

$$P(B|A) = 30/40 = 3/4$$

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

Some facts:

- This is a simple algebraic relationship using elementary definitions.*

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

Some facts:

- This is a simple algebraic relationship using elementary definitions.*
- It's interesting because it's kind of a "wormhole" between two different "interpretations" of probability.*

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

Some facts:

- This is a simple algebraic relationship using elementary definitions.*
- It's interesting because it's kind of a "wormhole" between two different "interpretations" of probability.*
- It's a very powerful computational tool.*