

RELATIVITY

37-1

WHAT IS PHYSICS?

One principal subject of physics is **relativity**, the field of study that measures events (things that happen): where and when they happen, and by how much any two events are separated in space and in time. In addition, relativity has to do with transforming such measurements (and also measurements of energy and momentum) between reference frames that move relative to each other. (Hence the name *relativity*.)

Transformations and moving reference frames, such as those we discussed in Sections 4-8 and 4-9, were well understood and quite routine to physicists in 1905. Then Albert Einstein (Fig. 37-1) published his **special theory of relativity**. The adjective *special* means that the theory deals only with **inertial reference frames**, which are frames in which Newton's laws are valid. (Einstein's *general theory of relativity* treats the more challenging situation in which reference frames can undergo gravitational acceleration; in this chapter the term *relativity* implies only inertial reference frames.)

Starting with two deceptively simple postulates, Einstein stunned the scientific world by showing that the old ideas about relativity were wrong, even though everyone was so accustomed to them that they seemed to be unquestionable common sense. This supposed common sense, however, was derived only from experience with things that move rather slowly. Einstein's relativity, which turns out to be correct for all physically possible speeds, predicted many effects that were, at first study, bizarre because no one had ever experienced them.

In particular, Einstein demonstrated that space and time are entangled; that is, the time between two events depends on how far apart they occur, and vice versa. Also, the entanglement is different for observers who move relative to each other. One result is that time does not pass at a fixed rate, as if it were ticked off with mechanical regularity on some master grandfather clock that controls the universe. Rather, that rate is adjustable: Relative motion can change the rate at which time passes. Prior to 1905, no one but a few daydreamers would have thought that. Now, engineers and scientists take it for granted because their experience with special relativity has reshaped their common sense. For example, any engineer involved with the Global Positioning System of the NAVSTAR satellites must routinely use relativity (both special relativity and general relativity) to determine the rate at which time passes on the satellites because that rate differs from the rate on Earth's surface. If the engineers failed to take relativity into account, GPS would become almost useless in less than one day.

Special relativity has the reputation of being difficult. It is not difficult mathematically, at least not here. However, it is difficult in that we must be very careful about *who* measures *what* about an event and just *how* that measurement is made—and it can be difficult because it can contradict routine experience.



Fig. 37-1 Einstein posing for a photograph as fame began to accumulate. (Corbis Images)

37-2 The Postulates

We now examine the two postulates of relativity, on which Einstein's theory is based:

- 1. The Relativity Postulate:** The laws of physics are the same for observers in all inertial reference frames. No one frame is preferred over any other.

Galileo assumed that the laws of *mechanics* were the same in all inertial reference frames. Einstein extended that idea to include *all* the laws of physics, especially those of electromagnetism and optics. This postulate does *not* say that the measured values of all physical quantities are the same for all inertial observers; most are not the same. It is the *laws of physics*, which relate these measurements to one another, that are the same.

- 2. The Speed of Light Postulate:** The speed of light in vacuum has the same value c in all directions and in all inertial reference frames.

We can also phrase this postulate to say that there is in nature an *ultimate speed* c , the same in all directions and in all inertial reference frames. Light happens to travel at this ultimate speed. However, no entity that carries energy or information can exceed this limit. Moreover, no particle that has mass can actually reach speed c , no matter how much or for how long that particle is accelerated. (Alas, the faster-than-light warp drive used in many science fiction stories appears to be impossible.)

Both postulates have been exhaustively tested, and no exceptions have ever been found.

The Ultimate Speed

The existence of a limit to the speed of accelerated electrons was shown in a 1964 experiment by W. Bertozzi, who accelerated electrons to various measured speeds and—by an independent method—measured their kinetic energies. He found that as the force on a very fast electron is increased, the electron's measured kinetic energy increases toward very large values but its speed does not increase appreciably (Fig. 37-2). Electrons have been accelerated in laboratories to at least 0.999 999 995 times the speed of light but—close though it may be—that speed is still less than the ultimate speed c .

This ultimate speed has been defined to be exactly

$$c = 299\,792\,458 \text{ m/s.} \quad (37-1)$$

Caution: So far in this book we have (appropriately) approximated c as $3.0 \times 10^8 \text{ m/s}$, but in this chapter we shall often use the exact value. You might want to store the exact value in your calculator's memory (if it is not there already), to be called up when needed.

Testing the Speed of Light Postulate

If the speed of light is the same in all inertial reference frames, then the speed of light emitted by a source relative to, say, a laboratory should be the same as the speed of light that is emitted by a source at rest in the laboratory. This claim has been tested directly, in an experiment of high precision. The “light source” was the *neutral pion* (symbol π^0), an unstable, short-lived particle that can be produced by collisions in a particle accelerator. It decays (transforms) into two gamma rays by the process

$$\pi^0 \rightarrow \gamma + \gamma. \quad (37-2)$$

Gamma rays are part of the electromagnetic spectrum (at very high frequencies)




Fig. 37-2 The dots show measured values of the kinetic energy of an electron plotted against its measured speed. No matter how much energy is given to an electron (or to any other particle having mass), its speed can never equal or exceed the ultimate limiting speed c . (The plotted curve through the dots shows the predictions of Einstein's special theory of relativity.)

and so obey the speed of light postulate, just as visible light does. (In this chapter we shall use the term light for any type of electromagnetic wave, visible or not.)

In 1964, physicists at CERN, the European particle-physics laboratory near Geneva, generated a beam of pions moving at a speed of $0.999\,75c$ with respect to the laboratory. The experimenters then measured the speed of the gamma rays emitted from these very rapidly moving sources. They found that the speed of the light emitted by the pions was the same as it would be if the pions were at rest in the laboratory, namely c .

37-3 Measuring an Event

An **event** is something that happens, and every event can be assigned three space coordinates and one time coordinate. Among many possible events are (1) the turning on or off of a tiny lightbulb, (2) the collision of two particles, (3) the passage of a pulse of light through a specified point, (4) an explosion, and (5) the sweeping of the hand of a clock past a marker on the rim of the clock. A certain observer, fixed in a certain inertial reference frame, might, for example, assign to an event A the coordinates given in Table 37-1. Because space and time are entangled with each other in relativity, we can describe these coordinates collectively as *spacetime* coordinates. The coordinate system itself is part of the reference frame of the observer.

A given event may be recorded by any number of observers, each in a different inertial reference frame. In general, different observers will assign different spacetime coordinates to the same event. Note that an event does not “belong” to any particular inertial reference frame. An event is just something that happens, and anyone in any reference frame may detect it and assign spacetime coordinates to it.

Making such an assignment can be complicated by a practical problem. For example, suppose a balloon bursts 1 km to your right while a firecracker pops 2 km to your left, both at 9:00 A.M. However, you do not detect either event precisely at 9:00 A.M. because at that instant light from the events has not yet reached you. Because light from the firecracker pop has farther to go, it arrives at your eyes later than does light from the balloon burst, and thus the pop will seem to have occurred later than the burst. To sort out the actual times and to assign 9:00 A.M. as the happening time for both events, you must calculate the travel times of the light and then subtract these times from the arrival times.

This procedure can be very messy in more challenging situations, and we need an easier procedure that automatically eliminates any concern about the travel time from an event to an observer. To set up such a procedure, we shall construct an imaginary array of measuring rods and clocks throughout the observer’s inertial frame (the array moves rigidly with the observer). This construction may seem contrived, but it spares us much confusion and calculation and allows us to find the coordinates, as follows.

Table 37-1	
Record of Event A	
Coordinate	Value
x	3.58 m
y	1.29 m
z	0 m
t	34.5 s

- 1. The Space Coordinates.** We imagine the observer’s coordinate system fitted with a close-packed, three-dimensional array of measuring rods, one set of rods parallel to each of the three coordinate axes. These rods provide a way to determine coordinates along the axes. Thus, if the event is, say, the turning on of a small lightbulb, the observer, in order to locate the position of the event, need only read the three space coordinates at the bulb’s location.
- 2. The Time Coordinate.** For the time coordinate, we imagine that every point of intersection in the array of measuring rods includes a tiny clock, which the observer can read because the clock is illuminated by the light generated by the event. Figure 37-3 suggests one plane in the “jungle gym” of clocks and measuring rods we have described.

The array of clocks must be synchronized properly. It is not enough to assemble a set of identical clocks, set them all to the same time, and then move them to their assigned positions. We do not know, for example, whether moving the clocks will change their rates. (Actually, it will.) We must put the clocks in place and *then* synchronize them.

If we had a method of transmitting signals at infinite speed, synchronization would be a simple matter. However, no known signal has this property. We therefore choose light (any part of the electromagnetic spectrum) to send out our synchronizing signals because, in vacuum, light travels at the greatest possible speed, the limiting speed c .

Here is one of many ways in which an observer might synchronize an array of clocks using light signals: The observer enlists the help of a great number of temporary helpers, one for each clock. The observer then stands at a point selected as the origin and sends out a pulse of light when the origin clock reads $t = 0$. When the light pulse reaches the location of a helper, that helper sets the clock there to read $t = r/c$, where r is the distance between the helper and the origin. The clocks are then synchronized.

- 3. The Spacetime Coordinates.** The observer can now assign spacetime coordinates to an event by simply recording the time on the clock nearest the event and the position as measured on the nearest measuring rods. If there are two events, the observer computes their separation in time as the difference in the times on clocks near each and their separation in space from the differences in coordinates on rods near each. We thus avoid the practical problem of calculating the travel times of the signals to the observer from the events.

We use this array to assign spacetime coordinates.




Fig. 37-3 One section of a three-dimensional array of clocks and measuring rods by which an observer can assign spacetime coordinates to an event, such as a flash of light at point A . The event's space coordinates are approximately $x = 3.6$ rod lengths, $y = 1.3$ rod lengths, and $z = 0$. The time coordinate is whatever time appears on the clock closest to A at the instant of the flash.

37-4 The Relativity of Simultaneity

Suppose that one observer (Sam) notes that two independent events (event Red and event Blue) occur at the same time. Suppose also that another observer (Sally), who is moving at a constant velocity $\vec{v}$ with respect to Sam, also records these same two events. Will Sally also find that they occur at the same time?

The answer is that in general she will not:

 If two observers are in relative motion, they will not, in general, agree as to whether two events are simultaneous. If one observer finds them to be simultaneous, the other generally will not.

We cannot say that one observer is right and the other wrong. Their observations are equally valid, and there is no reason to favor one over the other.

The realization that two contradictory statements about the same natural event can be correct is a seemingly strange outcome of Einstein's theory. However, in Chapter 17 we saw another way in which motion can affect measurement without balking at the contradictory results: In the Doppler effect, the frequency an observer measures for a sound wave depends on the relative motion of observer and source. Thus, two observers moving relative to each other can measure different frequencies for the same wave, and both measurements are correct.

We conclude the following:

 Simultaneity is not an absolute concept but rather a relative one, depending on the motion of the observer.

If the relative speed of the observers is very much less than the speed of light, then measured departures from simultaneity are so small that they are not noticeable. Such is the case for all our experiences of daily living; that is why the relativity of simultaneity is unfamiliar.

A Closer Look at Simultaneity

Let us clarify the relativity of simultaneity with an example based on the postulates of relativity, no clocks or measuring rods being directly involved. Figure 37-4 shows two long spaceships (the SS *Sally* and the SS *Sam*), which can serve as inertial reference frames for observers Sally and Sam. The two observers are stationed at the midpoints of their ships. The ships are separating along a common x axis, the relative velocity of *Sally* with respect to *Sam* being $\vec{v}$. Figure 37-4a shows the ships with the two observer stations momentarily aligned opposite each other.

Two large meteorites strike the ships, one setting off a red flare (event Red) and the other a blue flare (event Blue), not necessarily simultaneously. Each event leaves a permanent mark on each ship, at positions RR' and BB' .

Let us suppose that the expanding wavefronts from the two events happen to reach Sam at the same time, as Fig. 37-4b shows. Let us further suppose that, after the episode, Sam finds, by measuring the marks on his spaceship, that he was indeed stationed exactly halfway between the markers B and R on his ship when the two events occurred. He will say:

Sam Light from event Red and light from event Blue reached me at the same time. From the marks on my spaceship, I find that I was standing halfway between the two sources. Therefore, event Red and event Blue were simultaneous events.

As study of Fig. 37-4 shows, Sally and the expanding wavefront from event Red are moving *toward* each other, while she and the expanding wavefront from event Blue are moving in the *same direction*. Thus, the wavefront from event Red will reach Sally *before* the wavefront from event Blue does. She will say:

Sally Light from event Red reached me before light from event Blue did. From the marks on my spaceship, I found that I too was standing halfway between the two sources. Therefore, the events were not simultaneous; event Red occurred first, followed by event Blue.

These reports do not agree. Nevertheless, *both* observers are correct.

Note carefully that there is only one wavefront expanding from the site of each event and that *this wavefront travels with the same speed c in both reference frames*, exactly as the speed of light postulate requires.

It *might* have happened that the meteorites struck the ships in such a way that the two hits appeared to Sally to be simultaneous. If that had been the case, then Sam would have declared them not to be simultaneous.




Fig. 37-4 The spaceships of Sally and Sam and the occurrences of events from Sam's view. Sally's ship moves rightward with velocity $\vec{v}$. (a) Event Red occurs at positions RR' and event Blue occurs at positions BB' ; each event sends out a wave of light. (b) Sam simultaneously detects the waves from event Red and event Blue. (c) Sally detects the wave from event Red. (d) Sally detects the wave from event Blue.

37-5 The Relativity of Time

If observers who move relative to each other measure the time interval (or *temporal separation*) between two events, they generally will find different results. Why? Because the spatial separation of the events can affect the time intervals measured by the observers.

The time interval between two events depends on how far apart they occur in both space and time; that is, their spatial and temporal separations are entangled.


In this section we discuss this entanglement by means of an example; however, the example is restricted in a crucial way: *To one of two observers, the two events occur at the same location.* We shall not get to more general examples until Section 37-7.

Figure 37-5a shows the basics of an experiment Sally conducts while she and her equipment—a light source, a mirror, and a clock—ride in a train moving with constant velocity $\vec{v}$ relative to a station. A pulse of light leaves the light source B (event 1), travels vertically upward, is reflected vertically downward by the mirror, and then is detected back at the source (event 2). Sally measures a certain time interval Δt_0 between the two events, related to the distance D from source to mirror by

$$\Delta t_0 = \frac{2D}{c} \quad (\text{Sally}). \quad (37-3)$$

The two events occur at the same location in Sally's reference frame, and she needs only one clock C at that location to measure the time interval. Clock C is shown twice in Fig. 37-5a, at the beginning and end of the interval.

Consider now how these same two events are measured by Sam, who is standing on the station platform as the train passes. Because the equipment moves with the train during the travel time of the light, Sam sees the path of the light as shown in Fig. 37-5b. For him, the two events occur at different places in his reference frame, and so to measure the time interval between events, Sam must use *two* synchronized clocks, C_1 and C_2 , one at each event. According to Einstein's speed of light postulate, the light travels at the same speed c for Sam as



for Sally. Now, however, the light travels distance $2L$ between events 1 and 2. The time interval measured by Sam between the two events is

$$\Delta t = \frac{2L}{c} \quad (\text{Sam}), \quad (37-4)$$

in which

$$L = \sqrt{(\tfrac{1}{2}v \Delta t)^2 + D^2}. \quad (37-5)$$

From Eq. 37-3, we can write this as

$$L = \sqrt{(\tfrac{1}{2}v \Delta t)^2 + (\tfrac{1}{2}c \Delta t_0)^2}. \quad (37-6)$$

If we eliminate L between Eqs. 37-4 and 37-6 and solve for Δt , we find

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - (v/c)^2}}. \quad (37-7)$$

Equation 37-7 tells us how Sam's measured interval Δt between the events compares with Sally's interval Δt_0 . Because v must be less than c , the denominator in Eq. 37-7 must be less than unity. Thus, Δt must be greater than Δt_0 : Sam measures a *greater* time interval between the two events than does Sally. Sam and Sally have measured the time interval between the *same* two events, but the relative motion between Sam and Sally made their measurements *different*. We conclude that relative motion can change the *rate* at which time passes between two events; the key to this effect is the fact that the speed of light is the same for both observers.

We distinguish between the measurements of Sam and Sally in this way:



When two events occur at the same location in an inertial reference frame, the time interval between them, measured in that frame, is called the **proper time interval** or the **proper time**. Measurements of the same time interval from any other inertial reference frame are always greater.

Thus, Sally measures a proper time interval, and Sam measures a greater time interval. (The term *proper* is unfortunate in that it implies that any other measurement is improper or nonreal. That is just not so.) The amount by which a measured time interval is greater than the corresponding proper time interval is called **time dilation**. (To dilate is to expand or stretch; here the time interval is expanded or stretched.)

Often the dimensionless ratio v/c in Eq. 37-7 is replaced with β , called the **speed parameter**, and the dimensionless inverse square root in Eq. 37-7 is often replaced with γ , called the **Lorentz factor**:

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} = \frac{1}{\sqrt{1 - (v/c)^2}}. \quad (37-8)$$

With these replacements, we can rewrite Eq. 37-7 as

$$\Delta t = \gamma \Delta t_0 \quad (\text{time dilation}). \quad (37-9)$$

The speed parameter β is always less than unity, and, provided v is not zero, γ is always greater than unity. However, the difference between γ and 1 is not significant unless $v > 0.1c$. Thus, in general, “old relativity” works well enough for $v < 0.1c$, but we must use special relativity for greater values of v . As shown in Fig. 37-6, γ increases rapidly in magnitude as β approaches 1 (as v approaches c). Therefore, the greater the relative speed between Sally and Sam is, the greater will be the time interval measured by Sam, until at a great enough speed, the interval takes “forever.”

As the speed parameter goes to 1.0 (as the speed approaches c), the Lorentz factor approaches infinity.




Fig. 37-6 A plot of the Lorentz factor γ as a function of the speed parameter $\beta (= v/c)$.

You might wonder what Sally says about Sam's having measured a greater time interval than she did. His measurement comes as no surprise to her, because to her, he failed to synchronize his clocks C_1 and C_2 in spite of his insistence that he did. Recall that observers in relative motion generally do not agree about simultaneity. Here, Sam insists that his two clocks simultaneously read the same time when event 1 occurred. To Sally, however, Sam's clock C_2 was erroneously set ahead during the synchronization process. Thus, when Sam read the time of event 2 on it, to Sally he was reading off a time that was too large, and that is why the time interval he measured between the two events was greater than the interval she measured.

Two Tests of Time Dilation

1. Microscopic Clocks. Subatomic particles called *muons* are unstable; that is, when a muon is produced, it lasts for only a short time before it *decays* (transforms into particles of other types). The *lifetime* of a muon is the time interval between its production (event 1) and its decay (event 2). When muons are stationary and their lifetimes are measured with stationary clocks (say, in a laboratory), their average lifetime is $2.200\ \mu\text{s}$. This is a proper time interval because, for each muon, events 1 and 2 occur at the same location in the reference frame of the muon—namely, at the muon itself. We can represent this proper time interval with Δt_0 ; moreover, we can call the reference frame in which it is measured the *rest frame* of the muon.

If, instead, the muons are moving, say, through a laboratory, then measurements of their lifetimes made with the laboratory clocks should yield a greater average lifetime (a dilated average lifetime). To check this conclusion, measurements were made of the average lifetime of muons moving with a speed of $0.9994c$ relative to laboratory clocks. From Eq. 37-8, with $\beta = 0.9994$, the Lorentz factor for this speed is

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} = \frac{1}{\sqrt{1 - (0.9994)^2}} = 28.87.$$

Equation 37-9 then yields, for the average dilated lifetime,

$$\Delta t = \gamma \Delta t_0 = (28.87)(2.200\ \mu\text{s}) = 63.51\ \mu\text{s}.$$


The actual measured value matched this result within experimental error.

2. Macroscopic Clocks. In October 1971, Joseph Hafele and Richard Keating carried out what must have been a grueling experiment. They flew four portable atomic clocks twice around the world on commercial airlines, once in each direction. Their purpose was “to test Einstein’s theory of relativity with macroscopic clocks.” As we have just seen, the time dilation predictions of Einstein’s theory have been confirmed on a microscopic scale, but there is great comfort in seeing a confirmation made with an actual clock. Such macroscopic measurements became possible only because of the very high precision of modern atomic clocks. Hafele and Keating verified the predictions of the theory to within 10%. (Einstein’s *general* theory of relativity, which predicts that the rate at which time passes on a clock is influenced by the gravitational force on the clock, also plays a role in this experiment.)

A few years later, physicists at the University of Maryland carried out a similar experiment with improved precision. They flew an atomic clock round and round over Chesapeake Bay for flights lasting 15 h and succeeded in checking the time dilation prediction to better than 1%. Today, when atomic clocks are transported from one place to another for calibration or other purposes, the time dilation caused by their motion is always taken into account.

CHECKPOINT 1

Standing beside railroad tracks, we are suddenly startled by a relativistic boxcar traveling past us as shown in the figure. Inside, a well-equipped hobo fires a laser pulse from the front of the boxcar to its rear. (a) Is our measurement of the speed of the pulse greater than, less than, or the same as that measured by the hobo? (b) Is his measurement of the flight time of the pulse a proper time? (c) Are his measurement and our measurement of the flight time related by Eq. 37-9?



Sample Problem

Time dilation for a space traveler who returns to Earth

Your starship passes Earth with a relative speed of $0.9990c$. After traveling 10.0 y (your time), you stop at lookout post LP13, turn, and then travel back to Earth with the same relative speed. The trip back takes another 10.0 y (your time). How long does the round trip take according to measurements made on Earth? (Neglect any effects due to the accelerations involved with stopping, turning, and getting back up to speed.)

KEY IDEAS

We begin by analyzing the outward trip:

1. This problem involves measurements made from two (inertial) reference frames, one attached to Earth and the other (your reference frame) attached to your ship.
2. The outward trip involves two events: the start of the trip at Earth and the end of the trip at LP13.
3. Your measurement of 10.0 y for the outward trip is the proper time Δt_0 between those two events, because the events occur at the same location in your reference frame—namely, on your ship.

4. The Earth-frame measurement of the time interval Δt for the outward trip must be greater than Δt_0 , according to Eq. 37-9 ($\Delta t = \gamma \Delta t_0$) for time dilation.

Calculations: Using Eq. 37-8 to substitute for γ in Eq. 37-9, we find

$$\begin{aligned}\Delta t &= \frac{\Delta t_0}{\sqrt{1 - (v/c)^2}} \\ &= \frac{10.0 \text{ y}}{\sqrt{1 - (0.9990c/c)^2}} = (22.37)(10.0 \text{ y}) = 224 \text{ y}.\end{aligned}$$

On the return trip, we have the same situation and the same data. Thus, the round trip requires 20 y of your time but

$$\Delta t_{\text{total}} = (2)(224 \text{ y}) = 448 \text{ y} \quad (\text{Answer})$$

of Earth time. In other words, you have aged 20 y while the Earth has aged 448 y. Although you cannot travel into the past (as far as we know), you can travel into the future of, say, Earth, by using high-speed relative motion to adjust the rate at which time passes.

Sample Problem

Time dilation and travel distance for a relativistic particle

The elementary particle known as the *positive kaon* (K^+) is unstable in that it can *decay* (transform) into other particles. Although the decay occurs randomly, we find that, on average, a positive kaon has a lifetime of $0.1237 \mu\text{s}$ when stationary—that is, when the lifetime is measured in the rest frame of the kaon. If a positive kaon has a speed of $0.990c$ relative to a laboratory reference frame when the kaon is produced, how far can it travel in that frame during its lifetime according to *classical physics* (which is a reasonable approximation for speeds much less than c) and according to special relativity (which is correct for all physically possible speeds)?

KEY IDEAS

1. We have two (inertial) reference frames, one attached to the kaon and the other attached to the laboratory.
2. This problem also involves two events: the start of the kaon's travel (when the kaon is produced) and the end of that travel (at the end of the kaon's lifetime).
3. The distance traveled by the kaon between those two events is related to its speed v and the time interval for the travel by

$$v = \frac{\text{distance}}{\text{time interval}}. \quad (37-10)$$

With these ideas in mind, let us solve for the distance first with classical physics and then with special relativity.

Classical physics: In classical physics we would find the same distance and time interval (in Eq. 37-10) whether we measured them from the kaon frame or from the laboratory frame. Thus, we need not be careful about the frame in which the measurements are made. To find the kaon's travel distance d_{cp} according to classical physics, we first rewrite Eq. 37-10 as

$$d_{cp} = v \Delta t, \quad (37-11)$$

where Δt is the time interval between the two events in either frame. Then, substituting $0.990c$ for v and $0.1237 \mu\text{s}$ for Δt in Eq. 37-11, we find

$$\begin{aligned} d_{cp} &= (0.990c) \Delta t \\ &= (0.990)(299\,792\,458 \text{ m/s})(0.1237 \times 10^{-6} \text{ s}) \\ &= 36.7 \text{ m.} \end{aligned} \quad (\text{Answer})$$

This is how far the kaon would travel if classical physics were correct at speeds close to c .

Special relativity: In special relativity we must be very careful that both the distance and the time interval in Eq. 37-10 are measured in the *same* reference frame—especially when the speed is close to c , as here. Thus, to find the actual travel distance d_{sr} of the kaon *as measured from the laboratory frame* and according to special relativity, we rewrite Eq. 37-10 as

$$d_{sr} = v \Delta t, \quad (37-12)$$

where Δt is the time interval between the two events *as measured from the laboratory frame*.

Before we can evaluate d_{sr} in Eq. 37-12, we must find Δt . The $0.1237 \mu\text{s}$ time interval is a proper time because the two events occur at the same location in the kaon frame—namely, at the kaon itself. Therefore, let Δt_0 represent this proper time interval. Then we can use Eq. 37-9 ($\Delta t = \gamma \Delta t_0$) for time dilation to find the time interval Δt as measured from the laboratory frame. Using Eq. 37-8 to substitute for γ in Eq. 37-9 leads to

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - (v/c)^2}} = \frac{0.1237 \times 10^{-6} \text{ s}}{\sqrt{1 - (0.990c/c)^2}} = 8.769 \times 10^{-7} \text{ s.}$$

This is about seven times longer than the kaon's proper lifetime. That is, the kaon's lifetime is about seven times longer in the laboratory frame than in its own frame—the kaon's lifetime is dilated. We can now evaluate Eq. 37-12 for the travel distance d_{sr} in the laboratory frame as

$$\begin{aligned} d_{sr} &= v \Delta t = (0.990c) \Delta t \\ &= (0.990)(299\,792\,458 \text{ m/s})(8.769 \times 10^{-7} \text{ s}) \\ &= 260 \text{ m.} \end{aligned} \quad (\text{Answer})$$

This is about seven times d_{cp} . Experiments like the one outlined here, which verify special relativity, became routine in physics laboratories decades ago. The engineering design and the construction of any scientific or medical facility that employs high-speed particles must take relativity into account.




Additional examples, video, and practice available at WileyPLUS

37-6 The Relativity of Length

If you want to measure the length of a rod that is at rest with respect to you, you can—at your leisure—note the positions of its end points on a long stationary scale and subtract one reading from the other. If the rod is moving, however, you must note the positions of the end points *simultaneously* (in your reference frame) or your measurement cannot be called a length. Figure 37-7 suggests the difficulty of trying to measure the length of a moving penguin by locating its front and back at different times. Because simultaneity is relative and it enters into length measurements, length should also be a relative quantity. It is.

Fig. 37-7 If you want to measure the front-to-back length of a penguin while it is moving, you must mark the positions of its front and back simultaneously (in your reference frame), as in (a), rather than at different times, as in (b).



Let L_0 be the length of a rod that you measure when the rod is stationary (meaning you and it are in the same reference frame, the rod's rest frame). If, instead, there is relative motion at speed v between you and the rod *along the length of the rod*, then with simultaneous measurements you obtain a length L given by

$$L = L_0 \sqrt{1 - \beta^2} = \frac{L_0}{\gamma} \quad (\text{length contraction}). \quad (37-13)$$

Because the Lorentz factor γ is always greater than unity if there is relative motion, L is less than L_0 . The relative motion causes a *length contraction*, and L is called a *contracted length*. Because γ increases with speed v , the length contraction also increases with v .



The length L_0 of an object measured in the rest frame of the object is its **proper length** or **rest length**. Measurements of the length from any reference frame that is in relative motion *parallel* to that length are always less than the proper length.

Be careful: Length contraction occurs only along the direction of relative motion. Also, the length that is measured does not have to be that of an object like a rod or a circle. Instead, it can be the length (or distance) between two objects in the same rest frame—for example, the Sun and a nearby star (which are, at least approximately, at rest relative to each other).

Does a moving object *really* shrink? Reality is based on observations and measurements; if the results are always consistent and if no error can be determined, then what is observed and measured is real. In that sense, the object really does shrink. However, a more precise statement is that the object *is really measured* to shrink—motion affects that measurement and thus reality.

When you measure a contracted length for, say, a rod, what does an observer moving with the rod say of your measurement? To that observer, you did not locate the two ends of the rod simultaneously. (Recall that observers in motion relative to each other do not agree about simultaneity.) To the observer, you first located the rod's front end and then, slightly later, its rear end, and that is why you measured a length that is less than the proper length.

Proof of Eq. 37-13

Length contraction is a direct consequence of time dilation. Consider once more our two observers. This time, both Sally, seated on a train moving through a station, and Sam, again on the station platform, want to measure the length of the platform. Sam, using a tape measure, finds the length to be L_0 , a proper length because the platform is at rest with respect to him. Sam also notes that Sally, on the train, moves through this length in a time $\Delta t = L_0/v$, where v is the speed of the train; that is,

$$L_0 = v \Delta t \quad (\text{Sam}). \quad (37-14)$$

This time interval Δt is not a proper time interval because the two events that define it (Sally passes the back of the platform and Sally passes the front of the platform) occur at two different places, and therefore Sam must use two synchronized clocks to measure the time interval Δt .

For Sally, however, the platform is moving past her. She finds that the two events measured by Sam occur *at the same place* in her reference frame. She can time them with a single stationary clock, and so the interval Δt_0 that she measures is a proper time interval. To her, the length L of the platform is given by

$$L = v \Delta t_0 \quad (\text{Sally}). \quad (37-15)$$

If we divide Eq. 37-15 by Eq. 37-14 and apply Eq. 37-9, the time dilation equation, we have

$$\frac{L}{L_0} = \frac{v \Delta t_0}{v \Delta t} = \frac{1}{\gamma},$$

or

$$L = \frac{L_0}{\gamma}, \quad (37-16)$$

which is Eq. 37-13, the length contraction equation.

Sample Problem

Time dilation and length contraction as seen from each frame

In Fig. 37-8, Sally (at point *A*) and Sam's spaceship (of proper length $L_0 = 230$ m) pass each other with constant relative speed v . Sally measures a time interval of $3.57 \mu\text{s}$ for the ship to pass her (from the passage of point *B* in Fig. 37-8*a* to the passage of point *C* in Fig. 37-8*b*). In terms of c , what is the relative speed v between Sally and the ship?

KEY IDEAS

Let's assume that speed v is near c . Then:

1. This problem involves measurements made from two (inertial) reference frames, one attached to Sally and the other attached to Sam and his spaceship.
2. This problem also involves two events: the first is the passage of point *B* past Sally (Fig. 37-8*a*) and the second is the passage of point *C* past her (Fig. 37-8*b*).
3. From either reference frame, the other reference frame passes at speed v and moves a certain distance in the time interval between the two events:

$$v = \frac{\text{distance}}{\text{time interval}}. \quad (37-17)$$

Because speed v is assumed to be near the speed of light, we must be careful that the distance and the time interval in Eq. 37-17 are measured in the *same* reference frame. Otherwise, *speed* has no meaning.


Calculations: We are free to use either frame for the measurements. Because we know that the time interval Δt between the two events measured from Sally's frame is $3.57 \mu\text{s}$, let us also use the distance L between the two events measured from her frame. Equation 37-17 then becomes

$$v = \frac{L}{\Delta t}. \quad (37-18)$$

We do not know L , but we can relate it to the given L_0 : The distance between the two events as measured from Sam's frame is the ship's proper length L_0 . Thus, the distance L measured from Sally's frame must be less than L_0 , as given by Eq. 37-13 ($L = L_0/\gamma$) for length contraction. Substituting L_0/γ for L in Eq. 37-18 and then substituting Eq. 37-8 for γ , we find

$$v = \frac{L_0/\gamma}{\Delta t} = \frac{L_0 \sqrt{1 - (v/c)^2}}{\Delta t}.$$

Fig. 37-8 (a)–(b) Event 1 occurs when point *B* passes Sally (at point *A*) and event 2 occurs when point *C* passes her. (c)–(d) Event 1 occurs when Sally passes point *B* and event 2 occurs when she passes point *C*.



Solving this equation for v (notice that it is on the left and also buried in the Lorentz factor) leads us to

$$\begin{aligned} v &= \frac{L_0 c}{\sqrt{(c \Delta t)^2 + L_0^2}} \\ &= \frac{(230 \text{ m})c}{\sqrt{(299\,792\,458 \text{ m/s})^2 (3.57 \times 10^{-6} \text{ s})^2 + (230 \text{ m})^2}} \\ &= 0.210c. \quad (\text{Answer}) \end{aligned}$$

Thus, the relative speed between Sally and the ship is 21% of the speed of light.

Note that only the relative motion of Sally and Sam matters here; whether either is stationary relative to, say, a space station is irrelevant. In Figs. 37-8a and b we took Sally to be stationary, but we could instead have taken the ship to

be stationary, with Sally moving to the left past it. Event 1 is again when Sally and point B are aligned (Fig. 37-8c), and event 2 is again when Sally and point C are aligned (Fig. 37-8d). However, we are now using Sam's measurements. So the length between the two events in *his* frame is the proper length L_0 of the ship and the time interval between them is not Sally's measurement Δt but a dilated time interval $\gamma \Delta t$.

Substituting Sam's measurements into Eq. 37-17, we have

$$v = \frac{L_0}{\gamma \Delta t},$$

which is exactly what we found using Sally's measurements. Thus, we get the same result of $v = 0.210c$ with either set of measurements, *but we must be careful not to mix the measurements from the two frames.*

Sample Problem

Time dilation and length contraction in outrunning a supernova

Caught by surprise near a supernova, you race away from the explosion in your spaceship, hoping to outrun the high-speed material ejected toward you. Your Lorentz factor γ relative to the inertial reference frame of the local stars is 22.4.

(a) To reach a safe distance, you figure you need to cover $9.00 \times 10^{16} \text{ m}$ as measured in the reference frame of the local stars. How long will the flight take, as measured in that frame?

KEY IDEAS

From Chapter 2, for constant speed, we know that

$$\text{speed} = \frac{\text{distance}}{\text{time interval}}. \quad (37-19)$$

From Fig. 37-6, we see that because your Lorentz factor γ relative to the stars is 22.4 (large), your relative speed v is almost c —so close that we can approximate it as c . Then for speed $v \approx c$, we must be careful that the distance and the time interval in Eq. 37-19 are measured in the *same* reference frame.

Calculations: The given distance ($9.00 \times 10^{16} \text{ m}$) for the length of your travel path is measured in the reference frame of the stars, and the requested time interval Δt is to be measured in that same frame. Thus, we can write

$$\left(\text{time interval} \right)_{\text{relative to stars}} = \frac{\text{distance relative to stars}}{c}.$$

Then substituting the given distance, we find that

$$\begin{aligned} \left(\text{time interval} \right)_{\text{relative to stars}} &= \frac{9.00 \times 10^{16} \text{ m}}{299\,792\,458 \text{ m/s}} \\ &= 3.00 \times 10^8 \text{ s} = 9.51 \text{ y}. \quad (\text{Answer}) \end{aligned}$$

(b) How long does that trip take according to you (in your reference frame)?

KEY IDEAS

1. We now want the time interval measured in a different reference frame—namely, yours. Thus, we need to transform the data given in the reference frame of the stars to your frame.
2. The given path length of $9.00 \times 10^{16} \text{ m}$, measured in the reference frame of the stars, is a proper length L_0 , because the two ends of the path are at rest in that frame. As observed from your reference frame, the stars' reference frame and those two ends of the path race past you at a relative speed of $v \approx c$.
3. You measure a contracted length L_0/γ for the path, not the proper length L_0 .

Calculations: We can now rewrite Eq. 37-19 as

$$\left(\text{time interval} \right)_{\text{relative to you}} = \frac{\text{distance relative to you}}{c} = \frac{L_0/\gamma}{c}.$$

Substituting known data, we find

$$\begin{aligned} \left(\text{time interval} \right)_{\text{relative to you}} &= \frac{(9.00 \times 10^{16} \text{ m})/22.4}{299\,792\,458 \text{ m/s}} \\ &= 1.340 \times 10^7 \text{ s} = 0.425 \text{ y}. \quad (\text{Answer}) \end{aligned}$$

In part (a) we found that the flight takes 9.51 y in the reference frame of the stars. However, here we find that it takes only 0.425 y in your frame, due to the relative motion and the resulting contracted length of the path.



Additional examples, video, and practice available at WileyPLUS

37-7 The Lorentz Transformation

Figure 37-9 shows inertial reference frame S' moving with speed v relative to frame S , in the common positive direction of their horizontal axes (marked x and x'). An observer in S reports spacetime coordinates x, y, z, t for an event, and an observer in S' reports x', y', z', t' for the same event. How are these sets of numbers related?

We claim at once (although it requires proof) that the y and z coordinates, which are perpendicular to the motion, are not affected by the motion; that is, $y = y'$ and $z = z'$. Our interest then reduces to the relation between x and x' and that between t and t' .

The Galilean Transformation Equations

Prior to Einstein's publication of his special theory of relativity, the four coordinates of interest were assumed to be related by the *Galilean transformation equations*:

$$\begin{aligned}x' &= x - vt \\ t' &= t\end{aligned}\quad \begin{array}{l} \text{(Galilean transformation equations;} \\ \text{approximately valid at low speeds).} \end{array} \quad (37-20)$$

(These equations are written with the assumption that $t = t' = 0$ when the origins of S and S' coincide.) You can verify the first equation with Fig. 37-9. The second equation effectively claims that time passes at the same rate for observers in both reference frames. That would have been so obviously true to a scientist prior to Einstein that it would not even have been mentioned. When speed v is small compared to c , Eqs. 37-20 generally work well.




Fig. 37-9 Two inertial reference frames: frame S' has velocity $\vec{v}$ relative to frame S .

The Lorentz Transformation Equations

We state without proof that the correct transformation equations, which remain valid for all speeds up to the speed of light, can be derived from the postulates of relativity. The results, called the **Lorentz transformation equations*** or sometimes (more loosely) just the Lorentz transformations, are

$$\begin{aligned}x' &= \gamma(x - vt), \\ y' &= y, \\ z' &= z, \\ t' &= \gamma(t - vx/c^2)\end{aligned}\quad \begin{array}{l} \text{(Lorentz transformation equations;} \\ \text{valid at all physically possible speeds).} \end{array} \quad (37-21)$$

(The equations are written with the assumption that $t = t' = 0$ when the origins of S and S' coincide.) Note that the spatial values x and the temporal values t are

* You may wonder why we do not call these the *Einstein transformation equations* (and why not the *Einstein factor* for γ). H. A. Lorentz actually derived these equations before Einstein did, but as the great Dutch physicist graciously conceded, he did not take the further bold step of interpreting these equations as describing the true nature of space and time. It is this interpretation, first made by Einstein, that is at the heart of relativity.

bound together in the first and last equations. This entanglement of space and time was a prime message of Einstein's theory, a message that was long rejected by many of his contemporaries.

It is a formal requirement of relativistic equations that they should reduce to familiar classical equations if we let c approach infinity. That is, if the speed of light were infinitely great, *all* finite speeds would be “low” and classical equations would never fail. If we let $c \rightarrow \infty$ in Eqs. 37-21, $\gamma \rightarrow 1$ and these equations reduce—as we expect—to the Galilean equations (Eqs. 37-20). You should check this.

Equations 37-21 are written in a form that is useful if we are given x and t and wish to find x' and t' . We may wish to go the other way, however. In that case we simply solve Eqs. 37-21 for x and t , obtaining

$$x = \gamma(x' + vt') \quad \text{and} \quad t = \gamma(t' + vx'/c^2). \quad (37-22)$$

Comparison shows that, starting from either Eqs. 37-21 or Eqs. 37-22, you can find the other set by interchanging primed and unprimed quantities and reversing the sign of the relative velocity v . (For example, if the S' frame has a positive velocity relative to an observer in the S frame as in Fig. 37-9, then the S frame has a *negative* velocity relative to an observer in the S' frame.)

Equations 37-21 and 37-22 relate the coordinates of a single event as seen by two observers. Sometimes we want to know not the coordinates of a single event but the differences between coordinates for a pair of events. That is, if we label our events 1 and 2, we may want to relate

$$\Delta x = x_2 - x_1 \quad \text{and} \quad \Delta t = t_2 - t_1,$$

as measured by an observer in S , and

$$\Delta x' = x'_2 - x'_1 \quad \text{and} \quad \Delta t' = t'_2 - t'_1,$$

as measured by an observer in S' .

Table 37-2 displays the Lorentz equations in difference form, suitable for analyzing pairs of events. The equations in the table were derived by simply substituting differences (such as Δx and $\Delta x'$) for the four variables in Eqs. 37-21 and 37-22.

Table 37-2

The Lorentz Transformation Equations for Pairs of Events

1. $\Delta x = \gamma(\Delta x' + v \Delta t')$	1'. $\Delta x' = \gamma(\Delta x - v \Delta t)$
2. $\Delta t = \gamma(\Delta t' + v \Delta x'/c^2)$	2'. $\Delta t' = \gamma(\Delta t - v \Delta x/c^2)$

$$\gamma = \frac{1}{\sqrt{1 - (v/c)^2}} = \frac{1}{\sqrt{1 - \beta^2}}$$

Frame S' moves at velocity v relative to frame S .

Be careful: When substituting values for these differences, you must be consistent and not mix the values for the first event with those for the second event. Also, if, say, Δx is a negative quantity, you must be certain to include the minus sign in a substitution.



CHECKPOINT 2

In Fig. 37-9, frame S' has velocity $0.90c$ relative to frame S . An observer in frame S' measures two events as occurring at the following spacetime coordinates: event Yellow at (5.0 m, 20 ns) and event Green at (−2.0 m, 45 ns). An observer in frame S wants to find the temporal separation $\Delta t_{GY} = t_G - t_Y$ between the events. (a) Which equation in Table 37-2 should be used? (b) Should $+0.90c$ or $-0.90c$ be substituted for v in the parentheses on the equation's right side and in the Lorentz factor γ ? What value should be substituted into the (c) first and (d) second term in the parentheses?

37-8 Some Consequences of the Lorentz Equations

Here we use the equations of Table 37-2 to affirm some of the conclusions that we reached earlier by arguments based directly on the postulates.

Simultaneity

Consider Eq. 2 of Table 37-2,

$$\Delta t = \gamma \left(\Delta t' + \frac{v \Delta x'}{c^2} \right). \quad (37-23)$$

If two events occur at different places in reference frame S' of Fig. 37-9, then $\Delta x'$ in this equation is not zero. It follows that even if the events are simultaneous in S' (thus $\Delta t' = 0$), they will not be simultaneous in frame S . (This is in accord with our conclusion in Section 37-4.) The time interval between the events in S will be

$$\Delta t = \gamma \frac{v \Delta x'}{c^2} \quad (\text{simultaneous events in } S').$$

Thus, the spatial separation $\Delta x'$ guarantees a temporal separation Δt .

Time Dilation

Suppose now that two events occur at the same place in S' (thus $\Delta x' = 0$) but at different times (thus $\Delta t' \neq 0$). Equation 37-23 then reduces to

$$\Delta t = \gamma \Delta t' \quad (\text{events in same place in } S'). \quad (37-24)$$

This confirms time dilation between frames S and S' . Moreover, because the two events occur at the same place in S' , the time interval $\Delta t'$ between them can be measured with a single clock, located at that place. Under these conditions, the measured interval is a proper time interval, and we can label it Δt_0 as we have previously labeled proper times. Thus, with that label Eq. 37-24 becomes

$$\Delta t = \gamma \Delta t_0 \quad (\text{time dilation}),$$

which is exactly Eq. 37-9, the time dilation equation. Thus, time dilation is a special case of the more general Lorentz equations.

Length Contraction

Consider Eq. 1' of Table 37-2,

$$\Delta x' = \gamma(\Delta x - v \Delta t). \quad (37-25)$$

If a rod lies parallel to the x and x' axes of Fig. 37-9 and is at rest in reference frame S' , an observer in S' can measure its length at leisure. One way to do so is by subtracting the coordinates of the end points of the rod. The value of $\Delta x'$ that is obtained will be the proper length L_0 of the rod because the measurements are made in a frame where the rod is at rest.

Suppose the rod is moving in frame S . This means that Δx can be identified as the length L of the rod in frame S only if the coordinates of the rod's end points are measured *simultaneously*—that is, if $\Delta t = 0$. If we put $\Delta x' = L_0$, $\Delta x = L$, and $\Delta t = 0$ in Eq. 37-25, we find

$$L = \frac{L_0}{\gamma} \quad (\text{length contraction}), \quad (37-26)$$

which is exactly Eq. 37-13, the length contraction equation. Thus, length contraction is a special case of the more general Lorentz equations.

Sample Problem

Lorentz transformations and reversing the sequence of events

An Earth starship has been sent to check an Earth outpost on the planet P1407, whose moon houses a battle group of the often hostile Reptulians. As the ship follows a straight-line course first past the planet and then past the moon, it detects a high-energy microwave burst at the Reptulian moon base and then, 1.10 s later, an explosion at the Earth outpost, which is 4.00×10^8 m from the Reptulian base as measured from the ship's reference frame. The Reptulians have obviously attacked the Earth outpost, and so the starship begins to prepare for a confrontation with them.

(a) The speed of the ship relative to the planet and its moon is $0.980c$. What are the distance and time interval between the burst and the explosion as measured in the planet–moon frame (and thus according to the occupants of the stations)?

KEY IDEAS

1. This problem involves measurements made from two reference frames, the planet–moon frame and the starship frame.
2. We have two events: the burst and the explosion.
3. We need to transform the given data as measured in the starship frame to the corresponding data as measured in the planet–moon frame.

Starship frame: Before we get to the transformation, we need to carefully choose our notation. We begin with a sketch of the situation as shown in Fig. 37-10. There, we have chosen the ship's frame S to be stationary and the planet–moon frame S' to be moving with positive velocity (rightward). (This is an arbitrary choice; we could, instead, have chosen the planet–moon frame to be stationary. Then we would redraw $\vec{v}$ in Fig. 37-10 as being attached to the S frame and indicating leftward motion; v would then be a negative quantity. The results would be the same.) Let subscripts e and b represent the explosion and burst, respectively. Then the given data, all in the unprimed (starship) reference frame, are

$$\Delta x = x_e - x_b = +4.00 \times 10^8 \text{ m}$$

and $\Delta t = t_e - t_b = +1.10 \text{ s}$.

Here, Δx is a positive quantity because in Fig. 37-10, the coordinate x_e for the explosion is greater than the coordinate x_b for the burst; Δt is also a positive quantity because the time t_e of the explosion is greater (later) than the time t_b of the burst.

Planet–moon frame: We seek $\Delta x'$ and $\Delta t'$, which we shall get by transforming the given S -frame data to the planet–moon frame S' . Because we are considering a pair of events, we choose transformation equations from Table 37-2—namely, Eqs. 1' and 2':

$$\Delta x' = \gamma(\Delta x - v \Delta t) \quad (37-27)$$

The relative motion alters the time intervals between events and maybe even their sequence.




Fig. 37-10 A planet and its moon in reference frame S' move rightward with speed v relative to a starship in reference frame S .

$$\text{and} \quad \Delta t' = \gamma \left(\Delta t - \frac{v \Delta x}{c^2} \right). \quad (37-28)$$

Here, $v = +0.980c$ and the Lorentz factor is

$$\gamma = \frac{1}{\sqrt{1 - (v/c)^2}} = \frac{1}{\sqrt{1 - (+0.980c/c)^2}} = 5.0252.$$

Equation 37-27 then becomes

$$\begin{aligned} \Delta x' &= (5.0252)[4.00 \times 10^8 \text{ m} - (+0.980c)(1.10 \text{ s})] \\ &= 3.86 \times 10^8 \text{ m}, \end{aligned} \quad (\text{Answer})$$

and Eq. 37-28 becomes

$$\begin{aligned} \Delta t' &= (5.0252) \left[(1.10 \text{ s}) - \frac{(+0.980c)(4.00 \times 10^8 \text{ m})}{c^2} \right] \\ &= -1.04 \text{ s}. \end{aligned} \quad (\text{Answer})$$

(b) What is the meaning of the minus sign in the value for $\Delta t'$?

Reasoning: We must be consistent with the notation we set up in part (a). Recall how we originally defined the time interval between burst and explosion: $\Delta t = t_e - t_b = +1.10 \text{ s}$. To be consistent with that choice of notation, our definition of $\Delta t'$ must be $t'_e - t'_b$; thus, we have found that

$$\Delta t' = t'_e - t'_b = -1.04 \text{ s}.$$

The minus sign here tells us that $t'_b > t'_e$; that is, in the planet–moon reference frame, the burst occurred 1.04 s *after* the explosion, not 1.10 s *before* the explosion as detected in the ship frame.

(c) Did the burst cause the explosion, or vice versa?

KEY IDEA

The sequence of events measured in the planet–moon reference frame is the reverse of that measured in the ship frame. In either situation, if there is a causal relationship between the two events, information must travel from the location of one event to the location of the other to cause it.

Checking the speed: Let us check the required speed of the information. In the ship frame, this speed is

$$v_{\text{info}} = \frac{\Delta x}{\Delta t} = \frac{4.00 \times 10^8 \text{ m}}{1.10 \text{ s}} = 3.64 \times 10^8 \text{ m/s,}$$

but that speed is impossible because it exceeds c . In the planet–moon frame, the speed comes out to be $3.70 \times 10^8 \text{ m/s}$, also impossible. Therefore, neither event could possibly have caused the other event; that is, they are *unrelated* events. Thus, the starship should not confront the Reptulians.



Additional examples, video, and practice available at WileyPLUS

37-9 The Relativity of Velocities

Here we wish to use the Lorentz transformation equations to compare the velocities that two observers in different inertial reference frames S and S' would measure for the same moving particle. Let S' move with velocity v relative to S .

Suppose that the particle, moving with constant velocity parallel to the x and x' axes in Fig. 37-11, sends out two signals as it moves. Each observer measures the space interval and the time interval between these two events. These four measurements are related by Eqs. 1 and 2 of Table 37-2,

$$\Delta x = \gamma(\Delta x' + v \Delta t')$$

and

$$\Delta t = \gamma \left(\Delta t' + \frac{v \Delta x'}{c^2} \right).$$

If we divide the first of these equations by the second, we find

$$\frac{\Delta x}{\Delta t} = \frac{\Delta x' + v \Delta t'}{\Delta t' + v \Delta x'/c^2}.$$

Dividing the numerator and denominator of the right side by $\Delta t'$, we find

$$\frac{\Delta x}{\Delta t} = \frac{\Delta x'/\Delta t' + v}{1 + v(\Delta x'/\Delta t')/c^2}.$$


However, in the differential limit, $\Delta x/\Delta t$ is u , the velocity of the particle as measured in S , and $\Delta x'/\Delta t'$ is u' , the velocity of the particle as measured in S' . Then we have, finally,

$$u = \frac{u' + v}{1 + u'v/c^2} \quad (\text{relativistic velocity transformation}) \quad (37-29)$$

as the relativistic velocity transformation equation. This equation reduces to the classical, or Galilean, velocity transformation equation,

$$u = u' + v \quad (\text{classical velocity transformation}), \quad (37-30)$$

when we apply the formal test of letting $c \rightarrow \infty$. In other words, Eq. 37-29 is correct for all physically possible speeds, but Eq. 37-30 is approximately correct for speeds much less than c .



The speed of the moving particle depends on the frame.

Fig. 37-11 Reference frame S' moves with velocity $\vec{v}$ relative to frame S . A particle has velocity $\vec{u}'$ relative to reference frame S' and velocity $\vec{u}$ relative to reference frame S .

37-10 Doppler Effect for Light

In Section 17-9 we discussed the Doppler effect (a shift in detected frequency) for sound waves traveling in air. For such waves, the Doppler effect depends on two velocities—namely, the velocities of the source and detector with respect to the air. Air is the medium that transmits the waves.

That is not the situation with light waves, for they (and other electromagnetic waves) require no medium, being able to travel even through vacuum. The Doppler effect for light waves depends on only one velocity, the relative velocity $\vec{v}$ between source and detector, as measured from the reference frame of either. Let f_0 represent the **proper frequency** of the source—that is, the frequency that is measured by an observer in the rest frame of the source. Let f represent the frequency detected by an observer moving with velocity $\vec{v}$ relative to that rest frame. Then, when the direction of $\vec{v}$ is directly away from the source,

$$f = f_0 \sqrt{\frac{1 - \beta}{1 + \beta}} \quad (\text{source and detector separating}), \quad (37-31)$$

where $\beta = v/c$. When the direction of $\vec{v}$ is directly toward the source, we must change the signs in front of both β symbols in Eq. 37-31.

Low-Speed Doppler Effect

For low speeds ($\beta \ll 1$), Eq. 37-31 can be expanded in a power series in β and approximated as

$$f = f_0(1 - \beta + \tfrac{1}{2}\beta^2) \quad (\text{source and detector separating, } \beta \ll 1). \quad (37-32)$$

The corresponding low-speed equation for the Doppler effect with sound waves (or any waves except light waves) has the same first two terms but a different coefficient in the third term. Thus, the relativistic effect for low-speed light sources and detectors shows up only with the β^2 term.

A police radar unit employs the Doppler effect with microwaves to measure the speed v of a car. A source in the radar unit emits a microwave beam at a certain (proper) frequency f_0 along the road. A car that is moving toward the unit intercepts that beam but at a frequency that is shifted upward by the Doppler effect due to the car's motion toward the radar unit. The car reflects the beam back toward the radar unit. Because the car is moving toward the radar unit, the detector in the unit intercepts a reflected beam that is further shifted up in frequency. The unit compares that detected frequency with f_0 and computes the speed v of the car.

Astronomical Doppler Effect

In astronomical observations of stars, galaxies, and other sources of light, we can determine how fast the sources are moving, either directly away from us or directly toward us, by measuring the *Doppler shift* of the light that reaches us. If a certain star were at rest relative to us, we would detect light from it with a certain proper frequency f_0 . However, if the star is moving either directly away from us or directly toward us, the light we detect has a frequency f that is shifted from f_0 by the Doppler effect. This Doppler shift is due only to the *radial* motion of the star (its motion directly toward us or away from us), and the speed we can determine by measuring this Doppler shift is only the *radial speed* v of the star—that is, only the radial component of the star's velocity relative to us.

Suppose a star (or any other light source) moves away from us with a radial speed v that is low enough (β is small enough) for us to neglect the β^2 term in Eq. 37-32. Then we have

$$f = f_0(1 - \beta). \quad (37-33)$$

Because astronomical measurements involving light are usually done in wavelengths rather than frequencies, let us replace f with c/λ and f_0 with c/λ_0 , where λ is the measured wavelength and λ_0 is the **proper wavelength** (the wavelength associated with f_0). We then have

$$\frac{c}{\lambda} = \frac{c}{\lambda_0} (1 - \beta),$$

or
$$\lambda = \lambda_0 (1 - \beta)^{-1}. \quad (37-34)$$

Because we assume β is small, we can expand $(1 - \beta)^{-1}$ in a power series. Doing so and retaining only the first power of β , we have

$$\lambda = \lambda_0 (1 + \beta),$$

or
$$\beta = \frac{\lambda - \lambda_0}{\lambda_0}. \quad (37-35)$$

Replacing β with v/c and $\lambda - \lambda_0$ with $|\Delta\lambda|$ leads to


$$v = \frac{|\Delta\lambda|}{\lambda_0} c \quad (\text{radial speed of light source, } v \ll c). \quad (37-36)$$

The difference $\Delta\lambda$ is the *wavelength Doppler shift* of the light source. We enclose it with an absolute sign so that we always have a magnitude of the shift.

Equation 37-36 is an approximation that can be applied only when $v \ll c$. Under that condition, Eq. 37-36 can be applied whether the light source is moving toward or away from us. If it is moving away from us, then λ is longer than λ_0 , $\Delta\lambda$ is positive, and the Doppler shift is called a *red shift*. (The term *red* does not mean the detected light is red or even visible. It merely serves as a memory device because red is at the *long* wavelength end of the visible spectrum. Thus λ is longer than λ_0 .) If the light source is moving toward us, then λ is shorter than λ_0 , $\Delta\lambda$ is negative, and the Doppler shift is called a *blue shift*.

CHECKPOINT 3

The figure shows a source that emits light of proper frequency f_0 while moving directly toward the right with speed $c/4$ as measured from reference frame S . The figure also shows a light detector, which measures a frequency $f > f_0$ for the emitted light. (a) Is the detector moving toward the left or the right? (b) Is the speed of the detector as measured from reference frame S more than $c/4$, less than $c/4$, or equal to $c/4$?



Transverse Doppler Effect

So far, we have discussed the Doppler effect, here and in Chapter 17, only for situations in which the source and the detector move either directly toward or directly away from each other. Figure 37-12 shows a different arrangement, in which a source S moves past a detector D . When S reaches point P , the velocity of S is perpendicular to the line joining P and D , and at that instant S is moving neither toward nor away from D . If the source is emitting sound waves of frequency f_0 , D detects that frequency (with no Doppler effect) when it intercepts the waves that were emitted at point P . However, if the source is emitting light waves, there is still a Doppler effect, called the **transverse Doppler effect**. In this situation, the detected frequency of the light emitted when the source is at point P is

$$f = f_0 \sqrt{1 - \beta^2} \quad (\text{transverse Doppler effect}). \quad (37-37)$$




Fig. 37-12 A light source S travels with velocity $\vec{v}$ past a detector at D . The special theory of relativity predicts a transverse Doppler effect as the source passes through point P , where the direction of travel is perpendicular to the line extending through D . Classical theory predicts no such effect.

For low speeds ($\beta \ll 1$), Eq. 37-37 can be expanded in a power series in β and approximated as

$$f = f_0(1 - \frac{1}{2}\beta^2) \quad (\text{low speeds}). \quad (37-38)$$

Here the first term is what we would expect for sound waves, and again the relativistic effect for low-speed light sources and detectors appears with the β^2 term.

In principle, a police radar unit can determine the speed of a car even when the path of the radar beam is perpendicular (transverse) to the path of the car. However, Eq. 37-38 tells us that because β is small even for a fast car, the relativistic term $\beta^2/2$ in the transverse Doppler effect is extremely small. Thus, $f \approx f_0$ and the radar unit computes a speed of zero.

The transverse Doppler effect is really another test of time dilation. If we rewrite Eq. 37-37 in terms of the period T of oscillation of the emitted light wave instead of the frequency, we have, because $T = 1/f$,

$$T = \frac{T_0}{\sqrt{1 - \beta^2}} = \gamma T_0, \quad (37-39)$$

in which T_0 ($= 1/f_0$) is the **proper period** of the source. As comparison with Eq. 37-9 shows, Eq. 37-39 is simply the time dilation formula because a period is a time interval.

37-11 A New Look at Momentum

Suppose that a number of observers, each in a different inertial reference frame, watch an isolated collision between two particles. In classical mechanics, we have seen that—even though the observers measure different velocities for the colliding particles—they all find that the law of conservation of momentum holds. That is, they find that the total momentum of the system of particles after the collision is the same as it was before the collision.

How is this situation affected by relativity? We find that if we continue to define the momentum $\vec{p}$ of a particle as $m\vec{v}$, the product of its mass and its velocity, total momentum is *not* conserved for the observers in different inertial frames. We have two choices: (1) Give up the law of conservation of momentum or (2) see whether we can refine our definition of momentum in some new way so that the law of conservation of momentum still holds. The correct choice is the second one.

Consider a particle moving with constant speed v in the positive direction of an x axis. Classically, its momentum has magnitude

$$p = mv = m \frac{\Delta x}{\Delta t} \quad (\text{classical momentum}), \quad (37-40)$$

in which Δx is the distance it travels in time Δt . To find a relativistic expression for momentum, we start with the new definition

$$p = m \frac{\Delta x}{\Delta t_0}.$$

Here, as before, Δx is the distance traveled by a moving particle as viewed by an observer watching that particle. However, Δt_0 is the time required to travel that distance, measured not by the observer watching the moving particle but by an observer moving with the particle. The particle is at rest with respect to this second observer; thus that measured time is a proper time.

Using the time dilation formula, $\Delta t = \gamma \Delta t_0$ (Eq. 37-9), we can then write

$$p = m \frac{\Delta x}{\Delta t_0} = m \frac{\Delta x}{\Delta t} \frac{\Delta t}{\Delta t_0} = m \frac{\Delta x}{\Delta t} \gamma.$$

However, since $\Delta x/\Delta t$ is just the particle velocity v , we have

$$p = \gamma m v \quad (\text{momentum}). \quad (37-41)$$

Note that this differs from the classical definition of Eq. 37-40 only by the Lorentz factor γ . However, that difference is important: Unlike classical momentum, relativistic momentum approaches an infinite value as v approaches c .

We can generalize the definition of Eq. 37-41 to vector form as

$$\vec{p} = \gamma m \vec{v} \quad (\text{momentum}). \quad (37-42)$$

This equation gives the correct definition of momentum for all physically possible speeds. For a speed much less than c , it reduces to the classical definition of momentum ($\vec{p} = m\vec{v}$).

37-12 A New Look at Energy

Mass Energy

The science of chemistry was initially developed with the assumption that in chemical reactions, energy and mass are conserved separately. In 1905, Einstein showed that as a consequence of his theory of special relativity, mass can be considered to be another form of energy. Thus, the law of conservation of energy is really the law of conservation of mass–energy.

In a *chemical reaction* (a process in which atoms or molecules interact), the amount of mass that is transferred into other forms of energy (or vice versa) is such a tiny fraction of the total mass involved that there is no hope of measuring the mass change with even the best laboratory balances. Mass and energy truly *seem* to be separately conserved. However, in a *nuclear reaction* (in which nuclei or fundamental particles interact), the energy released is often about a million times greater than in a chemical reaction, and the change in mass can easily be measured.

An object's mass m and the equivalent energy E_0 are related by

$$E_0 = mc^2, \quad (37-43)$$

which, without the subscript 0, is the best-known science equation of all time. This energy that is associated with the mass of an object is called **mass energy** or **rest energy**. The second name suggests that E_0 is an energy that the object has even when it is at rest, simply because it has mass. (If you continue your study of physics beyond this book, you will see more refined discussions of the relation between mass and energy. You might even encounter disagreements about just what that relation is and means.)

Table 37-3 shows the (approximate) mass energy, or rest energy, of a few objects. The mass energy of, say, a U.S. penny is enormous; the equivalent amount

Table 37-3

The Energy Equivalents of a Few Objects

Object	Mass (kg)	Energy Equivalent	
Electron	$\approx 9.11 \times 10^{-31}$	$\approx 8.19 \times 10^{-14} \text{ J}$	($\approx 511 \text{ keV}$)
Proton	$\approx 1.67 \times 10^{-27}$	$\approx 1.50 \times 10^{-10} \text{ J}$	($\approx 938 \text{ MeV}$)
Uranium atom	$\approx 3.95 \times 10^{-25}$	$\approx 3.55 \times 10^{-8} \text{ J}$	($\approx 225 \text{ GeV}$)
Dust particle	$\approx 1 \times 10^{-13}$	$\approx 1 \times 10^4 \text{ J}$	($\approx 2 \text{ kcal}$)
U.S. penny	$\approx 3.1 \times 10^{-3}$	$\approx 2.8 \times 10^{14} \text{ J}$	($\approx 78 \text{ GW} \cdot \text{h}$)

of electrical energy would cost well over a million dollars. On the other hand, the entire annual U.S. electrical energy production corresponds to a mass of only a few hundred kilograms of matter (stones, burritos, or anything else).

In practice, SI units are rarely used with Eq. 37-43 because they are too large to be convenient. Masses are usually measured in atomic mass units, where

$$1 \text{ u} = 1.660\,538\,86 \times 10^{-27} \text{ kg}, \quad (37-44)$$

and energies are usually measured in electron-volts or multiples of it, where

$$1 \text{ eV} = 1.602\,176\,462 \times 10^{-19} \text{ J}. \quad (37-45)$$

In the units of Eqs. 37-44 and 37-45, the multiplying constant c^2 has the values

$$\begin{aligned} c^2 &= 9.314\,940\,13 \times 10^8 \text{ eV/u} = 9.314\,940\,13 \times 10^5 \text{ keV/u} \\ &= 931.494\,013 \text{ MeV/u}. \end{aligned} \quad (37-46)$$

Total Energy

Equation 37-43 gives, for any object, the mass energy E_0 that is associated with the object's mass m , regardless of whether the object is at rest or moving. If the object is moving, it has additional energy in the form of kinetic energy K . If we assume that the object's potential energy is zero, then its total energy E is the sum of its mass energy and its kinetic energy:

$$E = E_0 + K = mc^2 + K. \quad (37-47)$$

Although we shall not prove it, the total energy E can also be written as

$$E = \gamma mc^2, \quad (37-48)$$

where γ is the Lorentz factor for the object's motion.

Since Chapter 7, we have discussed many examples involving changes in the total energy of a particle or a system of particles. However, we did not include mass energy in the discussions because the changes in mass energy were either zero or small enough to be neglected. The law of conservation of total energy still applies when changes in mass energy are significant. Thus, regardless of what happens to the mass energy, the following statement from Section 8-8 is still true:



The total energy E of an *isolated system* cannot change.

For example, if the total mass energy of two interacting particles in an isolated system decreases, some other type of energy in the system must increase because the total energy cannot change.

In a system undergoing a chemical or nuclear reaction, a change in the total mass energy of the system due to the reaction is often given as a Q value. The Q value for a reaction is obtained from the relation

$$\left(\begin{array}{c} \text{system's initial} \\ \text{total mass energy} \end{array} \right) = \left(\begin{array}{c} \text{system's final} \\ \text{total mass energy} \end{array} \right) + Q$$

$$\text{or} \quad E_{0i} = E_{0f} + Q. \quad (37-49)$$

Using Eq. 37-43 ($E_0 = mc^2$), we can rewrite this in terms of the initial *total* mass M_i and the final *total* mass M_f as

$$M_i c^2 = M_f c^2 + Q$$

$$\text{or} \quad Q = M_i c^2 - M_f c^2 = -\Delta M c^2, \quad (37-50)$$

where the change in mass due to the reaction is $\Delta M = M_f - M_i$.

If a reaction results in the transfer of energy from mass energy to, say, kinetic energy of the reaction products, the system's total mass energy E_0 (and total mass M) decreases and Q is positive. If, instead, a reaction requires that energy be transferred to mass energy, the system's total mass energy E_0 (and its total mass M) increases and Q is negative.

For example, suppose two hydrogen nuclei undergo a *fusion reaction* in which they join together to form a single nucleus and release two particles in the process. The total mass energy (and total mass) of the resultant single nucleus and two released particles is less than the total mass energy (and total mass) of the initial hydrogen nuclei. Thus, the Q of the fusion reaction is positive, and energy is said to be *released* (transferred from mass energy) by the reaction. This release is important to you because the fusion of hydrogen nuclei in the Sun is one part of the process that results in sunshine on Earth and makes life here possible.

Kinetic Energy

In Chapter 7 we defined the kinetic energy K of an object of mass m moving at speed v well below c to be

$$K = \frac{1}{2}mv^2. \quad (37-51)$$

However, this classical equation is only an approximation that is good enough when the speed is well below the speed of light.

Let us now find an expression for kinetic energy that is correct for *all* physically possible speeds, including speeds close to c . Solving Eq. 37-47 for K and then substituting for E from Eq. 37-48 lead to

$$\begin{aligned} K &= E - mc^2 = \gamma mc^2 - mc^2 \\ &= mc^2(\gamma - 1) \quad (\text{kinetic energy}), \end{aligned} \quad (37-52)$$

where $\gamma (= 1/\sqrt{1 - (v/c)^2})$ is the Lorentz factor for the object's motion.

Figure 37-13 shows plots of the kinetic energy of an electron as calculated with the correct definition (Eq. 37-52) and the classical approximation (Eq. 37-51), both as functions of v/c . Note that on the left side of the graph the two plots coincide; this is the part of the graph—at lower speeds—where we have calculated kinetic energies so far in this book. This part of the graph tells us that we have been justified in calculating kinetic energy with the classical expression of Eq. 37-51. However, on the right side of the graph—at speeds near c —the two plots differ significantly. As v/c approaches 1.0, the plot for the classical definition of kinetic energy increases only moderately while the plot for the correct definition of kinetic energy increases dramatically, approaching an infinite value as v/c approaches 1.0. Thus, when an object's speed v is near c , we *must* use Eq. 37-52 to calculate its kinetic energy.

Figure 37-13 also tells us something about the work we must do on an object to increase its speed by, say, 1%. The required work W is equal to the resulting change ΔK in the object's kinetic energy. If the change is to occur on the low-speed, left side of Fig. 37-13, the required work might be modest. However, if the change is to occur on the high-speed, right side of Fig. 37-13, the required work could be enormous because the kinetic energy K increases so rapidly there with an increase in speed v . To increase an object's speed to c would require, in principle, an infinite amount of energy; thus, doing so is impossible.

The kinetic energies of electrons, protons, and other particles are often stated with the unit electron-volt or one of its multiples used as an adjective. For example, an electron with a kinetic energy of 20 MeV may be described as a 20 MeV electron.

As v/c approaches 1.0, the actual kinetic energy approaches infinity.




Fig. 37-13 The relativistic (Eq. 37-52) and classical (Eq. 37-51) equations for the kinetic energy of an electron, plotted as a function of v/c , where v is the speed of the electron and c is the speed of light. Note that the two curves blend together at low speeds and diverge widely at high speeds. Experimental data (at the $\times$ marks) show that at high speeds the relativistic curve agrees with experiment but the classical curve does not.

This might help you to remember the relations.




Fig. 37-14 A useful memory diagram for the relativistic relations among the total energy E , the rest energy or mass energy mc^2 , the kinetic energy K , and the momentum magnitude p .

Momentum and Kinetic Energy

In classical mechanics, the momentum p of a particle is mv and its kinetic energy K is $\frac{1}{2}mv^2$. If we eliminate v between these two expressions, we find a direct relation between momentum and kinetic energy:

$$p^2 = 2Km \quad (\text{classical}). \quad (37-53)$$

We can find a similar connection in relativity by eliminating v between the relativistic definition of momentum (Eq. 37-41) and the relativistic definition of kinetic energy (Eq. 37-52). Doing so leads, after some algebra, to

$$(pc)^2 = K^2 + 2Kmc^2. \quad (37-54)$$

With the aid of Eq. 37-47, we can transform Eq. 37-54 into a relation between the momentum p and the total energy E of a particle:

$$E^2 = (pc)^2 + (mc^2)^2. \quad (37-55)$$

The right triangle of Fig. 37-14 can help you keep these useful relations in mind. You can also show that, in that triangle,

$$\sin \theta = \beta \quad \text{and} \quad \cos \theta = 1/\gamma. \quad (37-56)$$

With Eq. 37-55 we can see that the product pc must have the same unit as energy E ; thus, we can express the unit of momentum p as an energy unit divided by c , usually as MeV/ c or GeV/ c in fundamental particle physics.



CHECKPOINT 4

Are (a) the kinetic energy and (b) the total energy of a 1 GeV electron more than, less than, or equal to those of a 1 GeV proton?

Sample Problem

Energy and momentum of a relativistic electron

(a) What is the total energy E of a 2.53 MeV electron?

KEY IDEA

From Eq. 37-47, the total energy E is the sum of the electron's mass energy (or rest energy) mc^2 and its kinetic energy:

$$E = mc^2 + K. \quad (37-57)$$

Calculations: The adjective “2.53 MeV” in the problem statement means that the electron's kinetic energy is 2.53 MeV. To evaluate the electron's mass energy mc^2 , we substitute the electron's mass m from Appendix B, obtaining

$$\begin{aligned} mc^2 &= (9.109 \times 10^{-31} \text{ kg})(299\,792\,458 \text{ m/s})^2 \\ &= 8.187 \times 10^{-14} \text{ J}. \end{aligned}$$

Then dividing this result by $1.602 \times 10^{-13} \text{ J/MeV}$ gives us 0.511 MeV as the electron's mass energy (confirming the value in Table 37-3). Equation 37-57 then yields

$$E = 0.511 \text{ MeV} + 2.53 \text{ MeV} = 3.04 \text{ MeV}. \quad (\text{Answer})$$

(b) What is the magnitude p of the electron's momentum, in the unit MeV/ c ? (Note that c is the symbol for the speed of light and not itself a unit.)

KEY IDEA

We can find p from the total energy E and the mass energy mc^2 via Eq. 37-55,

$$E^2 = (pc)^2 + (mc^2)^2.$$

Calculations: Solving for pc gives us

$$\begin{aligned} pc &= \sqrt{E^2 - (mc^2)^2} \\ &= \sqrt{(3.04 \text{ MeV})^2 - (0.511 \text{ MeV})^2} = 3.00 \text{ MeV}. \end{aligned}$$

Finally, dividing both sides by c we find

$$p = 3.00 \text{ MeV}/c. \quad (\text{Answer})$$



Additional examples, video, and practice available at WileyPLUS

Sample Problem

Energy and an astounding discrepancy in travel time

The most energetic proton ever detected in the cosmic rays coming to Earth from space had an astounding kinetic energy of 3.0×10^{20} eV (enough energy to warm a teaspoon of water by a few degrees).

(a) What were the proton's Lorentz factor γ and speed v (both relative to the ground-based detector)?

KEY IDEAS

(1) The proton's Lorentz factor γ relates its total energy E to its mass energy mc^2 via Eq. 37-48 ($E = \gamma mc^2$). (2) The proton's total energy is the sum of its mass energy mc^2 and its (given) kinetic energy K .

Calculations: Putting these ideas together we have

$$\gamma = \frac{E}{mc^2} = \frac{mc^2 + K}{mc^2} = 1 + \frac{K}{mc^2}. \quad (37-58)$$

From Table 37-3, the proton's mass energy mc^2 is 938 MeV. Substituting this and the given kinetic energy into Eq. 37-58, we obtain

$$\begin{aligned} \gamma &= 1 + \frac{3.0 \times 10^{20} \text{ eV}}{938 \times 10^6 \text{ eV}} \\ &= 3.198 \times 10^{11} \approx 3.2 \times 10^{11}. \quad (\text{Answer}) \end{aligned}$$

This computed value for γ is so large that we cannot use the definition of γ (Eq. 37-8) to find v . Try it; your calculator will tell you that β is effectively equal to 1 and thus that v is effectively equal to c . Actually, v is almost c , but we want a more accurate answer, which we can obtain by first solving Eq. 37-8 for $1 - \beta$. To begin we write

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}} = \frac{1}{\sqrt{(1 - \beta)(1 + \beta)}} \approx \frac{1}{\sqrt{2(1 - \beta)}},$$

where we have used the fact that β is so close to unity that $1 + \beta$ is very close to 2. (We can round off the sum of two very close numbers but not their difference.) The velocity we seek is contained in the $1 - \beta$ term. Solving for $1 - \beta$ then yields

$$\begin{aligned} 1 - \beta &= \frac{1}{2\gamma^2} = \frac{1}{(2)(3.198 \times 10^{11})^2} \\ &= 4.9 \times 10^{-24} \approx 5 \times 10^{-24}. \end{aligned}$$

Thus,
and, since $v = \beta c$,

$$v \approx 0.999\,999\,999\,999\,999\,999\,995c. \quad (\text{Answer})$$

(b) Suppose that the proton travels along a diameter of the Milky Way galaxy (9.8×10^4 ly). Approximately how long does the proton take to travel that diameter as measured from the common reference frame of Earth and the Galaxy?

Reasoning: We just saw that this *ultrarelativistic* proton is traveling at a speed barely less than c . By the definition of light-year, light takes 1 y to travel a distance of 1 ly, and so light should take 9.8×10^4 y to travel 9.8×10^4 ly, and this proton should take almost the same time. Thus, from our Earth–Milky Way reference frame, the proton's trip takes

$$\Delta t = 9.8 \times 10^4 \text{ y}. \quad (\text{Answer})$$

(c) How long does the trip take as measured in the reference frame of the proton?

KEY IDEAS

1. This problem involves measurements made from two (inertial) reference frames: one is the Earth–Milky Way frame and the other is attached to the proton.
2. This problem also involves two events: the first is when the proton passes one end of the diameter along the Galaxy, and the second is when it passes the opposite end.
3. The time interval between those two events as measured in the proton's reference frame is the proper time interval Δt_0 because the events occur at the same location in that frame—namely, at the proton itself.
4. We can find the proper time interval Δt_0 from the time interval Δt measured in the Earth–Milky Way frame by using Eq. 37-9 ($\Delta t = \gamma \Delta t_0$) for time dilation. (Note that we can use that equation because one of the time measures is a proper time. However, we get the same relation if we use a Lorentz transformation.)

Calculation: Solving Eq. 37-9 for Δt_0 and substituting γ from (a) and Δt from (b), we find

$$\begin{aligned} \Delta t_0 &= \frac{\Delta t}{\gamma} = \frac{9.8 \times 10^4 \text{ y}}{3.198 \times 10^{11}} \\ &= 3.06 \times 10^{-7} \text{ y} = 9.7 \text{ s}. \quad (\text{Answer}) \end{aligned}$$

In our frame, the trip takes 98 000 y. In the proton's frame, it takes 9.7 s! As promised at the start of this chapter, relative motion can alter the rate at which time passes, and we have here an extreme example.



Additional examples, video, and practice available at WileyPLUS

REVIEW & SUMMARY

The Postulates Einstein's **special theory of relativity** is based on two postulates:

1. The laws of physics are the same for observers in all inertial reference frames. No one frame is preferred over any other.
2. The speed of light in vacuum has the same value c in all directions and in all inertial reference frames.

The speed of light c in vacuum is an ultimate speed that cannot be exceeded by any entity carrying energy or information.

Coordinates of an Event Three space coordinates and one time coordinate specify an **event**. One task of special relativity is to relate these coordinates as assigned by two observers who are in uniform motion with respect to each other.

Simultaneous Events If two observers are in relative motion, they will not, in general, agree as to whether two events are simultaneous. If one of the observers finds two events at different locations to be simultaneous, the other will not, and conversely. Simultaneity is *not* an absolute concept but a relative one, depending on the motion of the observer. The relativity of simultaneity is a direct consequence of the finite ultimate speed c .

Time Dilation If two successive events occur at the same place in an inertial reference frame, the time interval Δt_0 between them, measured on a single clock where they occur, is the **proper time** between the events. *Observers in frames moving relative to that frame will measure a larger value for this interval.* For an observer moving with relative speed v , the measured time interval is

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - (v/c)^2}} = \frac{\Delta t_0}{\sqrt{1 - \beta^2}} = \gamma \Delta t_0 \quad (\text{time dilation}). \quad (37-7 \text{ to } 37-9)$$

Here $\beta = v/c$ is the **speed parameter** and $\gamma = 1/\sqrt{1 - \beta^2}$ is the **Lorentz factor**. An important result of time dilation is that moving clocks run slow as measured by an observer at rest.

Length Contraction The length L_0 of an object measured by an observer in an inertial reference frame in which the object is at rest is called its **proper length**. *Observers in frames moving relative to that frame and parallel to that length will measure a shorter length.* For an observer moving with relative speed v , the measured length is

$$L = L_0 \sqrt{1 - \beta^2} = \frac{L_0}{\gamma} \quad (\text{length contraction}). \quad (37-13)$$

The Lorentz Transformation The *Lorentz transformation* equations relate the spacetime coordinates of a single event as seen by observers in two inertial frames, S and S' , where S' is moving relative to S with velocity v in the positive x and x' direction. The four coordinates are related by

$$\begin{aligned} x' &= \gamma(x - vt), \\ y' &= y, \\ z' &= z, \\ t' &= \gamma(t - vx/c^2) \end{aligned} \quad (37-21)$$

Relativity of Velocities When a particle is moving with speed u' in the positive x' direction in an inertial reference frame S' that itself is moving with speed v parallel to the x direction of a second inertial frame S , the speed u of the particle as measured in S is

$$u = \frac{u' + v}{1 + u'v/c^2} \quad (\text{relativistic velocity}). \quad (37-29)$$

Relativistic Doppler Effect If a source emitting light waves of frequency f_0 moves directly away from a detector with relative radial speed v (and speed parameter $\beta = v/c$), the frequency f measured by the detector is

$$f = f_0 \sqrt{\frac{1 - \beta}{1 + \beta}}. \quad (37-31)$$

The motion results in a decrease in the detected frequency and (as more commonly measured) an increase in the wavelength (red shift). If the source moves directly toward the detector, the signs in Eq. 37-31 are reversed. Now the motion results in an increase in frequency and a decrease in wavelength (blue shift).

For astronomical observations, the Doppler effect is measured in wavelengths. For speeds much less than c , Eq. 37-31 leads to

$$v = \frac{|\Delta\lambda|}{\lambda_0} c, \quad (37-36)$$

where $\Delta\lambda (= \lambda - \lambda_0)$ is the *Doppler shift* in wavelength due to the motion.

Transverse Doppler Effect If the relative motion of the light source is perpendicular to a line joining the source and detector, the Doppler frequency formula is

$$f = f_0 \sqrt{1 - \beta^2}. \quad (37-37)$$

This **transverse Doppler effect** is due to time dilation.

Momentum and Energy The following definitions of linear momentum $\vec{p}$, kinetic energy K , and total energy E for a particle of mass m are valid at any physically possible speed:

$$\vec{p} = \gamma m \vec{v} \quad (\text{momentum}), \quad (37-42)$$

$$E = mc^2 + K = \gamma mc^2 \quad (\text{total energy}), \quad (37-47, 37-48)$$

$$K = mc^2(\gamma - 1) \quad (\text{kinetic energy}). \quad (37-52)$$

Here γ is the Lorentz factor for the particle's motion, and mc^2 is the *mass energy*, or *rest energy*, associated with the mass of the particle. These equations lead to the relationships

$$(pc)^2 = K^2 + 2Kmc^2, \quad (37-54)$$

$$\text{and} \quad E^2 = (pc)^2 + (mc^2)^2. \quad (37-55)$$

When a system of particles undergoes a chemical or nuclear reaction, the Q of the reaction is the negative of the change in the system's total mass energy:

$$Q = M_i c^2 - M_f c^2 = -\Delta M c^2, \quad (37-50)$$

where M_i is the system's total mass before the reaction and M_f is its total mass after the reaction.

QUESTIONS

1 A rod is to move at constant speed v along the x axis of reference frame S , with the rod's length parallel to that axis. An observer in frame S is to measure the length L of the rod. Which of the curves in Fig. 37-15 best gives length L (vertical axis of the graph) versus speed parameter β ?




Fig. 37-15
Questions 1 and 3.

2 Figure 37-16 shows a ship (attached to reference frame S') passing us (standing in reference frame S). A proton is fired at nearly the speed of light along the length of the ship, from the front to the rear. (a) Is the spatial separation $\Delta x'$ between the point at which the proton is fired and the point at which it hits the ship's rear wall a positive or negative quantity? (b) Is the temporal separation $\Delta t'$ between those events a positive or negative quantity?




Fig. 37-16 Question 2 and Problem 68.

3 Reference frame S' is to pass reference frame S at speed v along the common direction of the x' and x axes, as in Fig. 37-9. An observer who rides along with frame S' is to count off 25 s on his wristwatch. The corresponding time interval Δt is to be measured by an observer in frame S . Which of the curves in Fig. 37-15 best gives Δt (vertical axis of the graph) versus speed parameter β ?

4 Figure 37-17 shows two clocks in stationary frame S' (they are synchronized in that frame) and one clock in moving frame S . Clocks C_1 and C'_1 read zero when they pass each other. When clocks C_1 and C'_2 pass each other, (a) which clock has the smaller reading and (b) which clock measures a proper time?




Fig. 37-17 Question 4.

5 Figure 37-18 shows two clocks in stationary frame S (they are synchronized in that frame) and one clock in moving frame S' . Clocks C_1 and C'_1 read zero when they pass each other. When clocks C'_1 and C_2 pass each other, (a) which clock has the smaller reading and (b) which clock measures a proper time?




Fig. 37-18
Question 5.

6 Sam leaves Venus in a spaceship headed to Mars and passes Sally, who is on Earth, with a relative speed of $0.5c$. (a) Each measures the Venus–Mars voyage time. Who measures a proper time: Sam, Sally, or neither? (b) On the way, Sam sends a pulse of light to Mars. Each measures the travel time of the pulse. Who measures a proper time: Sam, Sally, or neither?

7 The plane of clocks and measuring rods in Fig. 37-19 is like that in Fig. 37-3. The clocks along the x axis are separated (center to cen-

ter) by 1 light-second, as are the clocks along the y axis, and all the clocks are synchronized via the procedure described in Section 37-3. When the initial synchronizing signal of $t = 0$ from the origin reaches (a) clock A, (b) clock B, and (c) clock C, what initial time is then set on those clocks? An event occurs at clock A when it reads 10 s. (d) How long does the signal of that event take to travel to an observer stationed at the origin? (e) What time does that observer assign to the event?




Fig. 37-19 Question 7.

8 The rest energy and total energy, respectively, of three particles, expressed in terms of a basic amount A are (1) A , $2A$; (2) A , $3A$; (3) $3A$, $4A$. Without written calculation, rank the particles according to their (a) mass, (b) kinetic energy, (c) Lorentz factor, and (d) speed, greatest first.




Fig. 37-20
Question 9.

9 Figure 37-20 shows the triangle of Fig 37-14 for six particles; the slanted lines 2 and 4 have the same length. Rank the particles according to (a) mass, (b) momentum magnitude, and (c) Lorentz factor, greatest first. (d) Identify which two particles have the same total energy. (e) Rank the three lowest-mass particles according to kinetic energy, greatest first.

10 While on board a starship, you intercept signals from four shuttle craft that are moving either directly toward or directly away from you. The signals have the same proper frequency f_0 . The speed and direction (both relative to you) of the shuttle craft are (a) $0.3c$ toward, (b) $0.6c$ toward, (c) $0.3c$ away, and (d) $0.6c$ away. Rank the shuttle craft according to the frequency you receive, greatest first.

11 Figure 37-21 shows one of four star cruisers that are in a race. As each cruiser passes the starting line, a shuttle craft leaves the cruiser and races toward the finish line. You, judging the race, are stationary relative to the starting and finish lines. The speeds v_c of the cruisers relative to you and the speeds v_s of the shuttle craft relative to their respective starships are, in that order, (1) $0.70c$, $0.40c$; (2) $0.40c$, $0.70c$; (3) $0.20c$, $0.90c$; (4) $0.50c$, $0.60c$. (a) Rank the shuttle craft according to their speeds relative to you, greatest first. (b) Rank the shuttle craft according to the distances their pilots measure from the starting line to the finish line, greatest first. (c) Each starship sends a signal to its shuttle craft at a certain frequency f_0 as measured on board the starship. Rank the shuttle craft according to the frequencies they detect, greatest first.




Fig. 37-21 Question 11.



Tutoring problem available (at instructor's discretion) in WileyPLUS and WebAssign



Worked-out solution available in Student Solutions Manual

WWW Worked-out solution is at



Number of dots indicates level of problem difficulty



Interactive solution is at

<http://www.wiley.com/college/halliday>Additional information available in *The Flying Circus of Physics* and at flyingcircusofphysics.com**sec. 37-5 The Relativity of Time**

•1 The mean lifetime of stationary muons is measured to be $2.2000 \mu\text{s}$. The mean lifetime of high-speed muons in a burst of cosmic rays observed from Earth is measured to be $16.000 \mu\text{s}$. To five significant figures, what is the speed parameter β of these cosmic-ray muons relative to Earth?


•2 To eight significant figures, what is speed parameter β if the Lorentz factor γ is (a) 1.010 000 0, (b) 10.000 000, (c) 100.000 00, and (d) 1000.000 0?

•3 You wish to make a round trip from Earth in a spaceship, traveling at constant speed in a straight line for exactly 6 months (as you measure the time interval) and then returning at the same constant speed. You wish further, on your return, to find Earth as it will be exactly 1000 years in the future. (a) To eight significant figures, at what speed parameter β must you travel? (b) Does it matter whether you travel in a straight line on your journey?

•4 (Come) back to the future. Suppose that a father is 20.00 y older than his daughter. He wants to travel outward from Earth for 2.000 y and then back to Earth for another 2.000 y (both intervals as he measures them) such that he is then 20.00 y younger than his daughter. What constant speed parameter β (relative to Earth) is required for the trip?

•5 ILW An unstable high-energy particle enters a detector and leaves a track of length 1.05 mm before it decays. Its speed relative to the detector was $0.992c$. What is its proper lifetime? That is, how long would the particle have lasted before decay had it been at rest with respect to the detector?

•6 Reference frame S' is to pass reference frame S at speed v along the common direction of the x' and x axes, as in Fig. 37-9. An observer who rides along with frame S' is to count off a certain time interval on his wristwatch. The corresponding time interval Δt is to be measured by an observer in frame S . Figure 37-22 gives Δt versus speed parameter β for a range of values for β . The vertical axis scale is set by $\Delta t_a = 14.0$ s. What is interval Δt if $v = 0.98c$?

**Fig. 37-22** Problem 6.

•7 The premise of the *Planet of the Apes* movies and book is that hibernating astronauts travel far into Earth's future, to a time when human civilization has been replaced by an ape civilization. Considering only special relativity, determine how far into Earth's

future the astronauts would travel if they slept for 120 y while traveling relative to Earth with a speed of $0.9990c$, first outward from Earth and then back again.

sec. 37-6 The Relativity of Length

•8 An electron of $\beta = 0.999\,987$ moves along the axis of an evacuated tube that has a length of 3.00 m as measured by a laboratory observer S at rest relative to the tube. An observer S' who is at rest relative to the electron, however, would see this tube moving with speed $v (= \beta c)$. What length would observer S' measure for the tube?

•9 SSM A spaceship of rest length 130 m races past a timing station at a speed of $0.740c$. (a) What is the length of the spaceship as measured by the timing station? (b) What time interval will the station clock record between the passage of the front and back ends of the ship?


•10 A meter stick in frame S' makes an angle of 30° with the x' axis. If that frame moves parallel to the x axis of frame S with speed $0.90c$ relative to frame S , what is the length of the stick as measured from S ?

•11 A rod lies parallel to the x axis of reference frame S , moving along this axis at a speed of $0.630c$. Its rest length is 1.70 m. What will be its measured length in frame S ?

•12 The length of a spaceship is measured to be exactly half its rest length. (a) To three significant figures, what is the speed parameter β of the spaceship relative to the observer's frame? (b) By what factor do the spaceship's clocks run slow relative to clocks in the observer's frame?

•13 GO A space traveler takes off from Earth and moves at speed $0.9900c$ toward the star Vega, which is 26.00 ly distant. How much time will have elapsed by Earth clocks (a) when the traveler reaches Vega and (b) when Earth observers receive word from the traveler that she has arrived? (c) How much older will Earth observers calculate the traveler to be (measured from her frame) when she reaches Vega than she was when she started the trip?

•14 A rod is to move at constant speed v along the x axis of reference frame S , with the rod's length parallel to that axis. An observer in frame S is to measure the length L of the rod. Figure 37-23 gives length L versus speed parameter β for a range of values for β . The vertical axis scale is set by $L_a = 1.00$ m. What is L if $v = 0.95c$?

**Fig. 37-23** Problem 14.

- 15  The center of our Milky Way galaxy is about 23 000 ly away. (a) To eight significant figures, at what constant speed parameter would you need to travel exactly 23 000 ly (measured in the Galaxy frame) in exactly 30 y (measured in your frame)? (b) Measured in your frame and in light-years, what length of the Galaxy would pass by you during the trip?

sec. 37-8 Some Consequences of the Lorentz Equations

- 16 Observer S reports that an event occurred on the x axis of his reference frame at $x = 3.00 \times 10^8$ m at time $t = 2.50$ s. Observer S' and her frame are moving in the positive direction of the x axis at a speed of $0.400c$. Further, $x = x' = 0$ at $t = t' = 0$. What are the (a) spatial and (b) temporal coordinate of the event according to S' ? If S' were, instead, moving in the *negative* direction of the x axis, what would be the (c) spatial and (d) temporal coordinate of the event according to S' ?

- 17 **SSM** **WWW** In Fig. 37-9, the origins of the two frames coincide at $t = t' = 0$ and the relative speed is $0.950c$. Two micrometeorites collide at coordinates $x = 100$ km and $t = 200$ μ s according to an observer in frame S . What are the (a) spatial and (b) temporal coordinate of the collision according to an observer in frame S' ?

- 18 Inertial frame S' moves at a speed of $0.60c$ with respect to frame S (Fig. 37-9). Further, $x = x' = 0$ at $t = t' = 0$. Two events are recorded. In frame S , event 1 occurs at the origin at $t = 0$ and event 2 occurs on the x axis at $x = 3.0$ km at $t = 4.0$ μ s. According to observer S' , what is the time of (a) event 1 and (b) event 2? (c) Do the two observers see the two events in the same sequence or the reverse sequence?

- 19 An experimenter arranges to trigger two flashbulbs simultaneously, producing a big flash located at the origin of his reference frame and a small flash at $x = 30.0$ km. An observer moving at a speed of $0.250c$ in the positive direction of x also views the flashes. (a) What is the time interval between them according to her? (b) Which flash does she say occurs first?

- 20  As in Fig. 37-9, reference frame S' passes reference frame S with a certain velocity. Events 1 and 2 are to have a certain temporal separation $\Delta t'$ according to the S' observer. However, their spatial separation $\Delta x'$ according to that observer has not been set yet. Figure 37-24 gives their temporal separation Δt according to the S observer as a function of $\Delta x'$ for a range of $\Delta x'$ values. The vertical axis scale is set by $\Delta t_a = 6.00$ μ s. What is $\Delta t'$?




Fig. 37-24 Problem 20.

- 21 **Relativistic reversal of events.** Figures 37-25a and b show the (usual) situation in which a primed reference frame passes an unprimed reference frame, in the common positive direction of the x

and x' axes, at a constant relative velocity of magnitude v . We are at rest in the unprimed frame; Bullwinkle, an astute student of relativity in spite of his cartoon upbringing, is at rest in the primed frame. The figures also indicate events A and B that occur at the following spacetime coordinates as measured in our unprimed frame and in Bullwinkle's primed frame:

Event	Unprimed	Primed
A	(x_A, t_A)	(x'_A, t'_A)
B	(x_B, t_B)	(x'_B, t'_B)

In our frame, event A occurs before event B , with temporal separation $\Delta t = t_B - t_A = 1.00$ μ s and spatial separation $\Delta x = x_B - x_A = 400$ m. Let $\Delta t'$ be the temporal separation of the events according to Bullwinkle. (a) Find an expression for $\Delta t'$ in terms of the speed parameter $\beta (= v/c)$ and the given data. Graph $\Delta t'$ versus β for the following two ranges of β :

- (b) 0 to 0.01 (v is low, from 0 to $0.01c$)
(c) 0.1 to 1 (v is high, from $0.1c$ to the limit c)

- (d) At what value of β is $\Delta t' = 0$? For what range of β is the sequence of events A and B according to Bullwinkle (e) the same as ours and (f) the reverse of ours? (g) Can event A cause event B , or vice versa? Explain.




Fig. 37-25 Problems 21, 22, 60, and 61.

- 22 For the passing reference frames in Fig. 37-25, events A and B occur at the following spacetime coordinates: according to the unprimed frame, (x_A, t_A) and (x_B, t_B) ; according to the primed frame, (x'_A, t'_A) and (x'_B, t'_B) . In the unprimed frame, $\Delta t = t_B - t_A = 1.00$ μ s and $\Delta x = x_B - x_A = 400$ m. (a) Find an expression for $\Delta x'$ in terms of the speed parameter β and the given data. Graph $\Delta x'$ versus β for two ranges of β : (b) 0 to 0.01 and (c) 0.1 to 1. (d) At what value of β is $\Delta x'$ minimum, and (e) what is that minimum?

- 23 **ILW** A clock moves along an x axis at a speed of $0.600c$ and reads zero as it passes the origin. (a) Calculate the clock's Lorentz factor. (b) What time does the clock read as it passes $x = 180$ m?

- 24 Bullwinkle in reference frame S' passes you in reference frame S along the common direction of the x' and x axes, as in Fig. 37-9. He carries three meter sticks: meter stick 1 is parallel to the x' axis, meter stick 2 is parallel to the y' axis, and meter stick 3 is parallel to the z' axis. On his wristwatch he counts off 15.0 s, which takes 30.0 s according to you. Two events occur during his passage. According to you, event 1 occurs at $x_1 = 33.0$ m and $t_1 = 22.0$ ns, and event 2 occurs at $x_2 = 53.0$ m and $t_2 = 62.0$ ns. According to your measurements, what is the length of (a) meter stick 1, (b) meter stick 2, and (c) meter stick 3? According to Bullwinkle, what are (d) the spatial separation and (e) the temporal separation between events 1 and 2, and (f) which event occurs first?

••25 In Fig. 37-9, observer S detects two flashes of light. A big flash occurs at $x_1 = 1200$ m and, $5.00 \mu\text{s}$ later, a small flash occurs at $x_2 = 480$ m. As detected by observer S' , the two flashes occur at a single coordinate x' . (a) What is the speed parameter of S' , and (b) is S' moving in the positive or negative direction of the x axis? To S' , (c) which flash occurs first and (d) what is the time interval between the flashes?

••26 In Fig. 37-9, observer S detects two flashes of light. A big flash occurs at $x_1 = 1200$ m and, slightly later, a small flash occurs at $x_2 = 480$ m. The time interval between the flashes is $\Delta t = t_2 - t_1$. What is the smallest value of Δt for which observer S' will determine that the two flashes occur at the same x' coordinate?

sec. 37-9 The Relativity of Velocities

•27 SSM A particle moves along the x' axis of frame S' with velocity $0.40c$. Frame S' moves with velocity $0.60c$ with respect to frame S . What is the velocity of the particle with respect to frame S ?

•28 In Fig. 37-11, frame S' moves relative to frame S with velocity $0.62c$ while a particle moves parallel to the common x and x' axes. An observer attached to frame S' measures the particle's velocity to be $0.47c$. In terms of c , what is the particle's velocity as measured by an observer attached to frame S according to the (a) relativistic and (b) classical velocity transformation? Suppose, instead, that the S' measure of the particle's velocity is $-0.47c$. What velocity does the observer in S now measure according to the (c) relativistic and (d) classical velocity transformation?

•29 Galaxy A is reported to be receding from us with a speed of $0.35c$. Galaxy B, located in precisely the opposite direction, is also found to be receding from us at this same speed. What multiple of c gives the recessional speed an observer on Galaxy A would find for (a) our galaxy and (b) Galaxy B?

•30 Stellar system Q_1 moves away from us at a speed of $0.800c$. Stellar system Q_2 , which lies in the same direction in space but is closer to us, moves away from us at speed $0.400c$. What multiple of c gives the speed of Q_2 as measured by an observer in the reference frame of Q_1 ?

••31 SSM ILW WWW A spaceship whose rest length is 350 m has a speed of $0.82c$ with respect to a certain reference frame. A micrometeorite, also with a speed of $0.82c$ in this frame, passes the spaceship on an antiparallel track. How long does it take this object to pass the ship as measured on the ship?

••32 GO In Fig. 37-26a, particle P is to move parallel to the x and x' axes of reference frames S and S' , at a certain velocity relative to




Fig. 37-26 Problem 32.

frame S . Frame S' is to move parallel to the x axis of frame S at velocity v . Figure 37-26b gives the velocity u' of the particle relative to frame S' for a range of values for v . The vertical axis scale is set by $u'_a = 0.800c$. What value will u' have if (a) $v = 0.90c$ and (b) $v \rightarrow c$?

••33 GO An armada of spaceships that is 1.00 ly long (in its rest frame) moves with speed $0.800c$ relative to a ground station in frame S . A messenger travels from the rear of the armada to the front with a speed of $0.950c$ relative to S . How long does the trip take as measured (a) in the messenger's rest frame, (b) in the armada's rest frame, and (c) by an observer in frame S ?

sec. 37-10 Doppler Effect for Light

•34 A sodium light source moves in a horizontal circle at a constant speed of $0.100c$ while emitting light at the proper wavelength of $\lambda_0 = 589.00$ nm. Wavelength λ is measured for that light by a detector fixed at the center of the circle. What is the wavelength shift $\lambda - \lambda_0$?

•35 SSM A spaceship, moving away from Earth at a speed of $0.900c$, reports back by transmitting at a frequency (measured in the spaceship frame) of 100 MHz. To what frequency must Earth receivers be tuned to receive the report?

•36 Certain wavelengths in the light from a galaxy in the constellation Virgo are observed to be 0.4% longer than the corresponding light from Earth sources. (a) What is the radial speed of this galaxy with respect to Earth? (b) Is the galaxy approaching or receding from Earth?

•37 Assuming that Eq. 37-36 holds, find how fast you would have to go through a red light to have it appear green. Take 620 nm as the wavelength of red light and 540 nm as the wavelength of green light.

•38 Figure 37-27 is a graph of intensity versus wavelength for light reaching Earth from galaxy NGC 7319, which is about 3×10^8 light-years away. The most intense light is emitted by the oxygen in NGC 7319. In a laboratory that emission is at wavelength $\lambda = 513$ nm, but in the light from NGC 7319 it has been shifted to 525 nm due to the Doppler effect (all the emissions from NGC 7319 have been shifted). (a) What is the radial speed of NGC 7319 relative to Earth? (b) Is the relative motion toward or away from our planet?




Fig. 37-27 Problem 38.

••39 SSM A spaceship is moving away from Earth at speed $0.20c$. A source on the rear of the ship emits light at wavelength 450 nm according to someone on the ship. What (a) wavelength and (b) color (blue, green, yellow, or red) are detected by someone on Earth watching the ship?

sec. 37-12 A New Look at Energy

•40 How much work must be done to increase the speed of an electron from rest to (a) $0.500c$, (b) $0.990c$, and (c) $0.9990c$?

•41 **SSM WWW** The mass of an electron is $9.109\,381\,88 \times 10^{-31}$ kg. To six significant figures, find (a) γ and (b) β for an electron with kinetic energy $K = 100.000$ MeV.

•42 What is the minimum energy that is required to break a nucleus of ^{12}C (of mass $11.996\,71$ u) into three nuclei of ^4He (of mass $4.001\,51$ u each)?

•43 How much work must be done to increase the speed of an electron (a) from $0.18c$ to $0.19c$ and (b) from $0.98c$ to $0.99c$? Note that the speed increase is $0.01c$ in both cases.

•44 In the reaction $p + {}^{19}\text{F} \rightarrow \alpha + {}^{16}\text{O}$, the masses are

$$\begin{aligned} m(p) &= 1.007825 \text{ u}, & m(\alpha) &= 4.002603 \text{ u}, \\ m(\text{F}) &= 18.998405 \text{ u}, & m(\text{O}) &= 15.994915 \text{ u}. \end{aligned}$$

Calculate the Q of the reaction from these data.

•45 In a high-energy collision between a cosmic-ray particle and a particle near the top of Earth's atmosphere, 120 km above sea level, a pion is created. The pion has a total energy E of 1.35×10^5 MeV and is traveling vertically downward. In the pion's rest frame, the pion decays 35.0 ns after its creation. At what altitude above sea level, as measured from Earth's reference frame, does the decay occur? The rest energy of a pion is 139.6 MeV.

•46 (a) If m is a particle's mass, p is its momentum magnitude, and K is its kinetic energy, show that

$$m = \frac{(pc)^2 - K^2}{2Kc^2}.$$

(b) For low particle speeds, show that the right side of the equation reduces to m . (c) If a particle has $K = 55.0$ MeV when $p = 121$ MeV/ c , what is the ratio m/m_e of its mass to the electron mass?

•47 **SSM** A 5.00-grain aspirin tablet has a mass of 320 mg. For how many kilometers would the energy equivalent of this mass power an automobile? Assume 12.75 km/L and a heat of combustion of 3.65×10^7 J/L for the gasoline used in the automobile.

•48 **GO** The mass of a muon is 207 times the electron mass; the average lifetime of muons at rest is $2.20 \mu\text{s}$. In a certain experiment, muons moving through a laboratory are measured to have an average lifetime of $6.90 \mu\text{s}$. For the moving muons, what are (a) β , (b) K , and (c) p (in MeV/ c)?

•49 **GO** As you read this page (on paper or monitor screen), a cosmic ray proton passes along the left-right width of the page with relative speed v and a total energy of 14.24 nJ. According to your measurements, that left-right width is 21.0 cm. (a) What is the width according to the proton's reference frame? How much time did the passage take according to (b) your frame and (c) the proton's frame?

•50 To four significant figures, find the following when the kinetic energy is 10.00 MeV: (a) γ and (b) β for an electron ($E_0 = 0.510\,998$ MeV), (c) γ and (d) β for a proton ($E_0 = 938.272$ MeV), and (e) γ and (f) β for an α particle ($E_0 = 3727.40$ MeV).

•51 **ILW** What must be the momentum of a particle with mass m so that the total energy of the particle is 3.00 times its rest energy?

•52 Apply the binomial theorem (Appendix E) to the last part of Eq. 37-52 for the kinetic energy of a particle. (a) Retain the first two terms of the expansion to show the kinetic energy in the form

$$K = (\text{first term}) + (\text{second term}).$$

The first term is the classical expression for kinetic energy. The second term is the first-order correction to the classical expression. Assume the particle is an electron. If its speed v is $c/20$, what is the value of (b) the classical expression and (c) the first-order correction? If the electron's speed is $0.80c$, what is the value of (d) the classical expression and (e) the first-order correction? (f) At what speed parameter β does the first-order correction become 10% or greater of the classical expression?

•53 In Section 28-6, we showed that a particle of charge q and mass m will move in a circle of radius $r = mv/|q|B$ when its velocity $\vec{v}$ is perpendicular to a uniform magnetic field $\vec{B}$. We also found that the period T of the motion is independent of speed v . These two results are approximately correct if $v \ll c$. For relativistic speeds, we must use the correct equation for the radius:

$$r = \frac{p}{|q|B} = \frac{\gamma mv}{|q|B}.$$

(a) Using this equation and the definition of period ($T = 2\pi r/v$), find the correct expression for the period. (b) Is T independent of v ? If a 10.0 MeV electron moves in a circular path in a uniform magnetic field of magnitude 2.20 T, what are (c) the radius according to Chapter 28, (d) the correct radius, (e) the period according to Chapter 28, and (f) the correct period?

•54 What is β for a particle with (a) $K = 2.00E_0$ and (b) $E = 2.00E_0$?

•55 A certain particle of mass m has momentum of magnitude mc . What are (a) β , (b) γ , and (c) the ratio K/E_0 ?

•56 (a) The energy released in the explosion of 1.00 mol of TNT is 3.40 MJ. The molar mass of TNT is 0.227 kg/mol. What weight of TNT is needed for an explosive release of 1.80×10^{14} J? (b) Can you carry that weight in a backpack, or is a truck or train required? (c) Suppose that in an explosion of a fission bomb, 0.080% of the fissionable mass is converted to released energy. What weight of fissionable material is needed for an explosive release of 1.80×10^{14} J? (d) Can you carry that weight in a backpack, or is a truck or train required?

•57 Quasars are thought to be the nuclei of active galaxies in the early stages of their formation. A typical quasar radiates energy at the rate of 10^{41} W. At what rate is the mass of this quasar being reduced to supply this energy? Express your answer in solar mass units per year, where one solar mass unit (1 smu = 2.0×10^{30} kg) is the mass of our Sun.

•58 The mass of an electron is $9.109\,381\,88 \times 10^{-31}$ kg. To eight significant figures, find the following for the given electron kinetic energy: (a) γ and (b) β for $K = 1.000\,000\,0$ keV, (c) γ and (d) β for $K = 1.000\,000\,0$ MeV, and then (e) γ and (f) β for $K = 1.000\,000\,0$ GeV.

•59 An alpha particle with kinetic energy 7.70 MeV collides with an ^{14}N nucleus at rest, and the two transform into an ^{17}O nucleus and a proton. The proton is emitted at 90° to the direction of the incident alpha particle and has a kinetic energy of 4.44 MeV. The masses of the various particles are alpha particle, 4.00260 u; ^{14}N , 14.00307 u; proton, 1.007825 u; and ^{17}O , 16.99914 u.

In MeV, what are (a) the kinetic energy of the oxygen nucleus and (b) the Q of the reaction? (*Hint:* The speeds of the particles are much less than c .)

Additional Problems

60 Temporal separation between two events. Events A and B occur with the following spacetime coordinates in the reference frames of Fig. 37-25: according to the unprimed frame, (x_A, t_A) and (x_B, t_B) ; according to the primed frame, (x'_A, t'_A) and (x'_B, t'_B) . In the unprimed frame, $\Delta t = t_B - t_A = 1.00 \mu\text{s}$ and $\Delta x = x_B - x_A = 240 \text{ m}$. (a) Find an expression for $\Delta t'$ in terms of the speed parameter β and the given data. Graph $\Delta t'$ versus β for the following two ranges of β : (b) 0 to 0.01 and (c) 0.1 to 1. (d) At what value of β is $\Delta t'$ minimum and (e) what is that minimum? (f) Can one of these events cause the other? Explain.

61 Spatial separation between two events. For the passing reference frames of Fig. 37-25, events A and B occur with the following spacetime coordinates: according to the unprimed frame, (x_A, t_A) and (x_B, t_B) ; according to the primed frame, (x'_A, t'_A) and (x'_B, t'_B) . In the unprimed frame, $\Delta t = t_B - t_A = 1.00 \mu\text{s}$ and $\Delta x = x_B - x_A = 240 \text{ m}$. (a) Find an expression for $\Delta x'$ in terms of the speed parameter β and the given data. Graph $\Delta x'$ versus β for two ranges of β : (b) 0 to 0.01 and (c) 0.1 to 1. (d) At what value of β is $\Delta x' = 0$?

62 In Fig. 37-28a, particle P is to move parallel to the x and x' axes of reference frames S and S' , at a certain velocity relative to frame S . Frame S' is to move parallel to the x axis of frame S at velocity v . Figure 37-28b gives the velocity u' of the particle relative to frame S' for a range of values for v . The vertical axis scale is set by $u'_a = -0.800c$. What value will u' have if (a) $v = 0.80c$ and (b) $v \rightarrow c$?




Fig. 37-28 Problem 62.

63 Superluminal jets. Figure 37-29a shows the path taken by a knot in a jet of ionized gas that has been expelled from a galaxy. The knot travels at constant velocity $\vec{v}$ at angle θ from the direction of Earth. The knot occasionally emits a burst of light, which is eventually detected on Earth. Two bursts are indicated in Fig. 37-29a, separated by time t as measured in a stationary frame near the bursts. The bursts are shown in Fig. 37-29b as if they were photographed on the same piece of film, first when light from burst 1 arrived on Earth and then later when light from burst 2 arrived. The apparent distance D_{app} traveled by the knot between the two bursts is the distance across an Earth-observer's view of the knot's path. The apparent time T_{app} between the bursts is the difference in the arrival times of the light from them. The apparent speed of the knot is then $V_{\text{app}} = D_{\text{app}}/T_{\text{app}}$. In terms of v , t , and θ , what are (a) D_{app} and (b) T_{app} ? (c) Evaluate V_{app} for $v = 0.980c$ and $\theta = 30.0^\circ$.

When superluminal (faster than light) jets were first observed, they seemed to defy special relativity—at least until the correct geometry (Fig. 37-29a) was understood.




Fig. 37-29 Problem 63.

64 Reference frame S' passes reference frame S with a certain velocity as in Fig. 37-9. Events 1 and 2 are to have a certain spatial separation $\Delta x'$ according to the S' observer. However, their temporal separation $\Delta t'$ according to that observer has not been set yet. Figure 37-30 gives their spatial separation Δx according to the S observer as a function of $\Delta t'$ for a range of $\Delta t'$ values. The vertical axis scale is set by $\Delta x_a = 10.0 \text{ m}$. What is $\Delta x'$?




Fig. 37-30 Problem 64.

65 Another approach to velocity transformations. In Fig. 37-31, reference frames B and C move past reference frame A in the com-




Fig. 37-31 Problems 65, 66, and 67.

mon direction of their x axes. Represent the x components of the velocities of one frame relative to another with a two-letter subscript. For example, v_{AB} is the x component of the velocity of A relative to B . Similarly, represent the corresponding speed parameters with two-letter subscripts. For example, β_{AB} ($= v_{AB}/c$) is the speed parameter corresponding to v_{AB} . (a) Show that

$$\beta_{AC} = \frac{\beta_{AB} + \beta_{BC}}{1 + \beta_{AB}\beta_{BC}}.$$

Let M_{AB} represent the ratio $(1 - \beta_{AB})/(1 + \beta_{AB})$, and let M_{BC} and M_{AC} represent similar ratios. (b) Show that the relation

$$M_{AC} = M_{AB}M_{BC}$$

is true by deriving the equation of part (a) from it.

66 Continuation of Problem 65. Use the result of part (b) in Problem 65 for the motion along a single axis in the following situation. Frame A in Fig. 37-31 is attached to a particle that moves with velocity $+0.500c$ past frame B , which moves past frame C with a velocity of $+0.500c$. What are (a) M_{AC} , (b) β_{AC} , and (c) the velocity of the particle relative to frame C ?

67 Continuation of Problem 65. Let reference frame C in Fig. 37-31 move past reference frame D (not shown). (a) Show that

$$M_{AD} = M_{AB}M_{BC}M_{CD}.$$

(b) Now put this general result to work: Three particles move parallel to a single axis on which an observer is stationed. Let plus and minus signs indicate the directions of motion along that axis. Particle A moves past particle B at $\beta_{AB} = +0.20$. Particle B moves past particle C at $\beta_{BC} = -0.40$. Particle C moves past observer D at $\beta_{CD} = +0.60$. What is the velocity of particle A relative to observer D ? (The solution technique here is *much* faster than using Eq. 37-29.)

68 Figure 37-16 shows a ship (attached to reference frame S') passing us (standing in reference frame S) with velocity $\vec{v} = 0.950c\hat{i}$. A proton is fired at speed $0.980c$ relative to the ship from the front of the ship to the rear. The proper length of the ship is 760 m. What is the temporal separation between the time the proton is fired and the time it hits the rear wall of the ship according to (a) a passenger in the ship and (b) us? Suppose that, instead, the proton is fired from the rear to the front. What then is the temporal separation between the time it is fired and the time it hits the front wall according to (c) the passenger and (d) us?

69 The car-in-the-garage problem. Carman has just purchased the world's longest stretch limo, which has a proper length of $L_c = 30.5$ m. In Fig. 37-32a, it is shown parked in front of a garage with a proper length of $L_g = 6.00$ m. The garage has a front door (shown open) and a back door (shown closed). The limo is obviously longer than the garage. Still, Garageman, who owns the garage and knows something about relativistic length contraction, makes a bet with Carman that the limo can fit in the garage with both doors closed. Carman, who dropped his physics course before reaching special relativity, says such a thing, even in principle, is impossible.

To analyze Garageman's scheme, an x_c axis is attached to the limo, with $x_c = 0$ at the rear bumper, and an x_g axis is attached to the garage, with $x_g = 0$ at the (now open) front door. Then Carman is to drive the limo directly toward the front door at a velocity of $0.9980c$ (which is, of course, both technically and financially impossible). Carman is stationary in the x_c reference frame; Garageman is stationary in the x_g reference frame.

There are two events to consider. *Event 1*: When the rear bumper clears the front door, the front door is closed. Let the time of this event be zero to both Carman and Garageman: $t_{g1} = t_{c1} = 0$. The event occurs at $x_c = x_g = 0$. Figure 37-32b shows event 1 according to the x_g reference frame. *Event 2*: When the front bumper reaches the back door, that door opens. Figure 37-32c shows event 2 according to the x_g reference frame.

According to Garageman, (a) what is the length of the limo, and what are the spacetime coordinates (b) x_{g2} and (c) t_{g2} of event 2? (d) For how long is the limo temporarily "trapped" inside the garage with both doors shut? Now consider the situation from the x_c reference frame, in which the garage comes racing past the limo at a velocity of $-0.9980c$. According to Carman, (e) what is the length of the passing garage, what are the spacetime coordinates (f) x_{c2} and (g) t_{c2} of event 2, (h) is the limo ever in the garage with both doors shut, and (i) which event occurs first? (j) Sketch events 1 and 2 as seen by Carman. (k) Are the events causally related; that is, does one of them cause the other? (l) Finally, who wins the bet?




Fig. 37-32 Problem 69.

70 An airplane whose rest length is 40.0 m is moving at uniform velocity with respect to Earth, at a speed of 630 m/s. (a) By what fraction of its rest length is it shortened to an observer on Earth? (b) How long would it take, according to Earth clocks, for the airplane's clock to fall behind by $1.00 \mu\text{s}$?

71 **SSM** To circle Earth in low orbit, a satellite must have a speed of about 2.7×10^4 km/h. Suppose that two such satellites orbit Earth in opposite directions. (a) What is their relative speed as they pass, according to the classical Galilean velocity transformation equation? (b) What fractional error do you make in (a) by not using the (correct) relativistic transformation equation?

72 Find the speed parameter of a particle that takes 2.0 y longer than light to travel a distance of 6.0 ly.

73 **SSM** How much work is needed to accelerate a proton from a speed of 0.9850c to a speed of 0.9860c?

74 A pion is created in the higher reaches of Earth's atmosphere when an incoming high-energy cosmic-ray particle collides with an atomic nucleus. A pion so formed descends toward Earth with a speed of $0.99c$. In a reference frame in which they are at rest, pions decay with an average life of 26 ns. As measured in a frame fixed with respect to Earth, how far (on the average) will such a pion move through the atmosphere before it decays?

75 SSM If we intercept an electron having total energy 1533 MeV that came from Vega, which is 26 ly from us, how far in light-years was the trip in the rest frame of the electron?

76 The total energy of a proton passing through a laboratory apparatus is 10.611 nJ. What is its speed parameter β ? Use the proton mass given in Appendix B under “Best Value,” not the commonly remembered rounded number.

77 A spaceship at rest in a certain reference frame S is given a speed increment of $0.50c$. Relative to its new rest frame, it is then given a further $0.50c$ increment. This process is continued until its speed with respect to its original frame S exceeds $0.999c$. How many increments does this process require?

78 In the red shift of radiation from a distant galaxy, a certain radiation, known to have a wavelength of 434 nm when observed in the laboratory, has a wavelength of 462 nm. (a) What is the radial speed of the galaxy relative to Earth? (b) Is the galaxy approaching or receding from Earth?

79 SSM What is the momentum in MeV/ c of an electron with a kinetic energy of 2.00 MeV?

80 The radius of Earth is 6370 km, and its orbital speed about the Sun is 30 km/s. Suppose Earth moves past an observer at this speed. To the observer, by how much does Earth’s diameter contract along the direction of motion?

81 A particle with mass m has speed $c/2$ relative to inertial frame S . The particle collides with an identical particle at rest relative to frame S . Relative to S , what is the speed of a frame S' in which the total momentum of these particles is zero? This frame is called the *center of momentum frame*.

82 An elementary particle produced in a laboratory experiment travels 0.230 mm through the lab at a relative speed of $0.960c$ before it decays (becomes another particle). (a) What is the proper lifetime of the particle? (b) What is the distance the particle travels as measured from its rest frame?

83 What are (a) K , (b) E , and (c) p (in GeV/ c) for a proton moving at speed $0.990c$? What are (d) K , (e) E , and (f) p (in MeV/ c) for an electron moving at speed $0.990c$?

84 A radar transmitter T is fixed to a reference frame S' that is moving to the right with speed v relative to reference frame S (Fig. 37-33). A mechanical timer (essentially a clock) in frame S' , having a period τ_0 (measured in S'), causes transmitter T to emit timed radar pulses, which travel at the speed of light and are received by R , a receiver fixed in frame S . (a) What is the period τ of the timer as detected by observer A , who is fixed in frame S ? (b) Show that at receiver R the time interval between pulses arriving from T is not τ or τ_0 , but

$$\tau_R = \tau_0 \sqrt{\frac{c+v}{c-v}}.$$

(c) Explain why receiver R and observer A , who are in the same

reference frame, measure a different period for the transmitter. (*Hint:* A clock and a radar pulse are not the same thing.)




Fig. 37-33 Problem 84.

85 One cosmic-ray particle approaches Earth along Earth’s north–south axis with a speed of $0.80c$ toward the geographic north pole, and another approaches with a speed of $0.60c$ toward the geographic south pole (Fig. 37-34). What is the relative speed of approach of one particle with respect to the other?




Fig. 37-34 Problem 85.

86 (a) How much energy is released in the explosion of a fission bomb containing 3.0 kg of fissionable material? Assume that 0.10% of the mass is converted to released energy. (b) What mass of TNT would have to explode to provide the same energy release? Assume that each mole of TNT liberates 3.4 MJ of energy on exploding. The molecular mass of TNT is 0.227 kg/mol. (c) For the same mass of explosive, what is the ratio of the energy released in a nuclear explosion to that released in a TNT explosion?

87 (a) What potential difference would accelerate an electron to speed c according to classical physics? (b) With this potential difference, what speed would the electron actually attain?

88 A Foron cruiser moving directly toward a Reptulian scout ship fires a decoy toward the scout ship. Relative to the scout ship, the speed of the decoy is $0.980c$ and the speed of the Foron cruiser is $0.900c$. What is the speed of the decoy relative to the cruiser?

The International System of Units (SI)*

TABLE 1

The SI Base Units

Quantity	Name	Symbol	Definition
length	meter	m	"... the length of the path traveled by light in vacuum in $1/299,792,458$ of a second." (1983)
mass	kilogram	kg	"... this prototype [a certain platinum-iridium cylinder] shall henceforth be considered to be the unit of mass." (1889)
time	second	s	"... the duration of 9,192,631,770 periods of the radiation corresponding to the transition between the two hyperfine levels of the ground state of the cesium-133 atom." (1967)
electric current	ampere	A	"... that constant current which, if maintained in two straight parallel conductors of infinite length, of negligible circular cross section, and placed 1 meter apart in vacuum, would produce between these conductors a force equal to 2×10^{-7} newton per meter of length." (1946)
thermodynamic temperature	kelvin	K	"... the fraction $1/273.16$ of the thermodynamic temperature of the triple point of water." (1967)
amount of substance	mole	mol	"... the amount of substance of a system which contains as many elementary entities as there are atoms in 0.012 kilogram of carbon-12." (1971)
luminous intensity	candela	cd	"... the luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency 540×10^{12} hertz and that has a radiant intensity in that direction of $1/683$ watt per steradian." (1979)

*Adapted from "The International System of Units (SI)," National Bureau of Standards Special Publication 330, 1972 edition. The definitions above were adopted by the General Conference of Weights and Measures, an international body, on the dates shown. In this book we do not use the candela.

TABLE 2

Some SI Derived Units

Quantity	Name of Unit	Symbol	
area	square meter	m^2	
volume	cubic meter	m^3	
frequency	hertz	Hz	s^{-1}
mass density (density)	kilogram per cubic meter	kg/m^3	
speed, velocity	meter per second	m/s	
angular velocity	radian per second	rad/s	
acceleration	meter per second per second	m/s^2	
angular acceleration	radian per second per second	rad/s^2	
force	newton	N	$\text{kg} \cdot \text{m}/\text{s}^2$
pressure	pascal	Pa	N/m^2
work, energy, quantity of heat	joule	J	$\text{N} \cdot \text{m}$
power	watt	W	J/s
quantity of electric charge	coulomb	C	$\text{A} \cdot \text{s}$
potential difference, electromotive force	volt	V	W/A
electric field strength	volt per meter (or newton per coulomb)	V/m	N/C
electric resistance	ohm	Ω	V/A
capacitance	farad	F	$\text{A} \cdot \text{s}/\text{V}$
magnetic flux	weber	Wb	$\text{V} \cdot \text{s}$
inductance	henry	H	$\text{V} \cdot \text{s}/\text{A}$
magnetic flux density	tesla	T	Wb/m^2
magnetic field strength	ampere per meter	A/m	
entropy	joule per kelvin	J/K	
specific heat	joule per kilogram kelvin	$\text{J}/(\text{kg} \cdot \text{K})$	
thermal conductivity	watt per meter kelvin	$\text{W}/(\text{m} \cdot \text{K})$	
radiant intensity	watt per steradian	W/sr	

TABLE 3

The SI Supplementary Units

Quantity	Name of Unit	Symbol
plane angle	radian	rad
solid angle	steradian	sr

Some Fundamental Constants of Physics*

Constant	Symbol	Computational Value	Best (1998) Value	
			Value ^a	Uncertainty ^b
Speed of light in a vacuum	c	3.00×10^8 m/s	2.997 924 58	exact
Elementary charge	e	1.60×10^{-19} C	1.602 176 487	0.025
Gravitational constant	G	6.67×10^{-11} m ³ /s ² · kg	6.674 28	100
Universal gas constant	R	8.31 J/mol · K	8.314 472	1.7
Avogadro constant	N_A	6.02×10^{23} mol ⁻¹	6.022 141 79	0.050
Boltzmann constant	k	1.38×10^{-23} J/K	1.380 650 4	1.7
Stefan–Boltzmann constant	σ	5.67×10^{-8} W/m ² · K ⁴	5.670 400	7.0
Molar volume of ideal gas at STP ^d	V_m	2.27×10^{-2} m ³ /mol	2.271 098 1	1.7
Permittivity constant	ϵ_0	8.85×10^{-12} F/m	8.854 187 817 62	exact
Permeability constant	μ_0	1.26×10^{-6} H/m	1.256 637 061 43	exact
Planck constant	h	6.63×10^{-34} J · s	6.626 068 96	0.050
Electron mass ^c	m_e	9.11×10^{-31} kg	9.109 382 15	0.050
		5.49×10^{-4} u	5.485 799 094 3	4.2×10^{-4}
Proton mass ^c	m_p	1.67×10^{-27} kg	1.672 621 637	0.050
		1.0073 u	1.007 276 466 77	1.0×10^{-4}
Ratio of proton mass to electron mass	m_p/m_e	1840	1836.152 672 47	4.3×10^{-4}
Electron charge-to-mass ratio	e/m_e	1.76×10^{11} C/kg	1.758 820 150	0.025
Neutron mass ^c	m_n	1.68×10^{-27} kg	1.674 927 211	0.050
		1.0087 u	1.008 664 915 97	4.3×10^{-4}
Hydrogen atom mass ^c	m_{1H}	1.0078 u	1.007 825 031 6	0.0005
Deuterium atom mass ^c	m_{2H}	2.0136 u	2.013 553 212 724	3.9×10^{-5}
Helium atom mass ^c	m_{4He}	4.0026 u	4.002 603 2	0.067
Muon mass	m_μ	1.88×10^{-28} kg	1.883 531 30	0.056
Electron magnetic moment	μ_e	9.28×10^{-24} J/T	9.284 763 77	0.025
Proton magnetic moment	μ_p	1.41×10^{-26} J/T	1.410 606 662	0.026
Bohr magneton	μ_B	9.27×10^{-24} J/T	9.274 009 15	0.025
Nuclear magneton	μ_N	5.05×10^{-27} J/T	5.050 783 24	0.025
Bohr radius	a	5.29×10^{-11} m	5.291 772 085 9	6.8×10^{-4}
Rydberg constant	R	1.10×10^7 m ⁻¹	1.097 373 156 852 7	6.6×10^{-6}
Electron Compton wavelength	λ_C	2.43×10^{-12} m	2.426 310 217 5	0.0014

^aValues given in this column should be given the same unit and power of 10 as the computational value.

^bParts per million.

^cMasses given in u are in unified atomic mass units, where $1 \text{ u} = 1.660\,538\,782 \times 10^{-27} \text{ kg}$.

^dSTP means standard temperature and pressure: 0°C and 1.0 atm (0.1 MPa).

*The values in this table were selected from the 1998 CODATA recommended values (www.physics.nist.gov).

Some Astronomical Data

Some Distances from Earth

To the Moon*	3.82×10^8 m	To the center of our galaxy	2.2×10^{20} m
To the Sun*	1.50×10^{11} m	To the Andromeda Galaxy	2.1×10^{22} m
To the nearest star (Proxima Centauri)	4.04×10^{16} m	To the edge of the observable universe	$\sim 10^{26}$ m

*Mean distance.

The Sun, Earth, and the Moon

Property	Unit	Sun	Earth	Moon
Mass	kg	1.99×10^{30}	5.98×10^{24}	7.36×10^{22}
Mean radius	m	6.96×10^8	6.37×10^6	1.74×10^6
Mean density	kg/m ³	1410	5520	3340
Free-fall acceleration at the surface	m/s ²	274	9.81	1.67
Escape velocity	km/s	618	11.2	2.38
Period of rotation ^a	—	37 d at poles ^b 26 d at equator ^b	23 h 56 min	27.3 d
Radiation power ^c	W	3.90×10^{26}		

^aMeasured with respect to the distant stars.

^bThe Sun, a ball of gas, does not rotate as a rigid body.

^cJust outside Earth's atmosphere solar energy is received, assuming normal incidence, at the rate of 1340 W/m².

Some Properties of the Planets

	Mercury	Venus	Earth	Mars	Jupiter	Saturn	Uranus	Neptune	Pluto
Mean distance from Sun, 10 ⁶ km	57.9	108	150	228	778	1430	2870	4500	5900
Period of revolution, y	0.241	0.615	1.00	1.88	11.9	29.5	84.0	165	248
Period of rotation, ^a d	58.7	−243 ^b	0.997	1.03	0.409	0.426	−0.451 ^b	0.658	6.39
Orbital speed, km/s	47.9	35.0	29.8	24.1	13.1	9.64	6.81	5.43	4.74
Inclination of axis to orbit	<28°	≈3°	23.4°	25.0°	3.08°	26.7°	97.9°	29.6°	57.5°
Inclination of orbit to Earth's orbit	7.00°	3.39°		1.85°	1.30°	2.49°	0.77°	1.77°	17.2°
Eccentricity of orbit	0.206	0.0068	0.0167	0.0934	0.0485	0.0556	0.0472	0.0086	0.250
Equatorial diameter, km	4880	12 100	12 800	6790	143 000	120 000	51 800	49 500	2300
Mass (Earth = 1)	0.0558	0.815	1.000	0.107	318	95.1	14.5	17.2	0.002
Density (water = 1)	5.60	5.20	5.52	3.95	1.31	0.704	1.21	1.67	2.03
Surface value of g , ^c m/s ²	3.78	8.60	9.78	3.72	22.9	9.05	7.77	11.0	0.5
Escape velocity, ^c km/s	4.3	10.3	11.2	5.0	59.5	35.6	21.2	23.6	1.3
Known satellites	0	0	1	2	63 + ring	60 + rings	27 + rings	13 + rings	3

^aMeasured with respect to the distant stars.

^bVenus and Uranus rotate opposite their orbital motion.

^cGravitational acceleration measured at the planet's equator.

Conversion Factors

Conversion factors may be read directly from these tables. For example, 1 degree = 2.778×10^{-3} revolutions, so $16.7^\circ = 16.7 \times 2.778 \times 10^{-3}$ rev. The SI units are fully capitalized. Adapted in part from G. Shortley and D. Williams, *Elements of Physics*, 1971, Prentice-Hall, Englewood Cliffs, NJ.

Plane Angle

	°	'	"	RADIAN	rev
1 degree = 1		60	3600	1.745×10^{-2}	2.778×10^{-3}
1 minute = 1.667×10^{-2}		1	60	2.909×10^{-4}	4.630×10^{-5}
1 second = 2.778×10^{-4}		1.667×10^{-2}	1	4.848×10^{-6}	7.716×10^{-7}
1 RADIAN = 57.30		3438	2.063×10^5	1	0.1592
1 revolution = 360		2.16×10^4	1.296×10^6	6.283	1

Solid Angle

$$1 \text{ sphere} = 4\pi \text{ steradians} = 12.57 \text{ steradians}$$

Length

	cm	METER	km	in.	ft	mi
1 centimeter = 1		10^{-2}	10^{-5}	0.3937	3.281×10^{-2}	6.214×10^{-6}
1 METER = 100		1	10^{-3}	39.37	3.281	6.214×10^{-4}
1 kilometer = 10^5		1000	1	3.937×10^4	3281	0.6214
1 inch = 2.540		2.540×10^{-2}	2.540×10^{-5}	1	8.333×10^{-2}	1.578×10^{-5}
1 foot = 30.48		0.3048	3.048×10^{-4}	12	1	1.894×10^{-4}
1 mile = 1.609×10^5		1609	1.609	6.336×10^4	5280	1
1 angström = 10^{-10} m		1 fermi = 10^{-15} m		1 fathom = 6 ft		1 rod = 16.5 ft
1 nautical mile = 1852 m		1 light-year = 9.461×10^{12} km		1 Bohr radius = 5.292×10^{-11} m		1 mil = 10^{-3} in.
= 1.151 miles = 6076 ft		1 parsec = 3.084×10^{13} km		1 yard = 3 ft		1 nm = 10^{-9} m

Area

	METER ²	cm ²	ft ²	in. ²
1 SQUARE METER = 1		10^4	10.76	1550
1 square centimeter = 10^{-4}		1	1.076×10^{-3}	0.1550
1 square foot = 9.290×10^{-2}		929.0	1	144
1 square inch = 6.452×10^{-4}		6.452	6.944×10^{-3}	1
1 square mile = 2.788×10^7 ft ² = 640 acres			1 acre = 43 560 ft ²	
1 barn = 10^{-28} m ²			1 hectare = 10^4 m ² = 2.471 acres	

A-6 APPENDIX D CONVERSION FACTORS

Volume

	METER ³	cm ³	L	ft ³	in. ³
1 CUBIC METER = 1		10 ⁶	1000	35.31	6.102 × 10 ⁴
1 cubic centimeter = 10 ⁻⁶		1	1.000 × 10 ⁻³	3.531 × 10 ⁻⁵	6.102 × 10 ⁻²
1 liter = 1.000 × 10 ⁻³		1000	1	3.531 × 10 ⁻²	61.02
1 cubic foot = 2.832 × 10 ⁻²		2.832 × 10 ⁴	28.32	1	1728
1 cubic inch = 1.639 × 10 ⁻⁵		16.39	1.639 × 10 ⁻²	5.787 × 10 ⁻⁴	1

1 U.S. fluid gallon = 4 U.S. fluid quarts = 8 U.S. pints = 128 U.S. fluid ounces = 231 in.³

1 British imperial gallon = 277.4 in.³ = 1.201 U.S. fluid gallons

Mass

Quantities in the colored areas are not mass units but are often used as such. For example, when we write 1 kg “=” 2.205 lb, this means that a kilogram is a *mass* that *weighs* 2.205 pounds at a location where *g* has the standard value of 9.80665 m/s².

	g	KILOGRAM	slug	u	oz	lb	ton
1 gram = 1		0.001	6.852 × 10 ⁻⁵	6.022 × 10 ²³	3.527 × 10 ⁻²	2.205 × 10 ⁻³	1.102 × 10 ⁻⁶
1 KILOGRAM = 1000		1	6.852 × 10 ⁻²	6.022 × 10 ²⁶	35.27	2.205	1.102 × 10 ⁻³
1 slug = 1.459 × 10 ⁴		14.59	1	8.786 × 10 ²⁷	514.8	32.17	1.609 × 10 ⁻²
1 atomic mass unit = 1.661 × 10 ⁻²⁴		1.661 × 10 ⁻²⁷	1.138 × 10 ⁻²⁸	1	5.857 × 10 ⁻²⁶	3.662 × 10 ⁻²⁷	1.830 × 10 ⁻³⁰
1 ounce = 28.35		2.835 × 10 ⁻²	1.943 × 10 ⁻³	1.718 × 10 ²⁵	1	6.250 × 10 ⁻²	3.125 × 10 ⁻⁵
1 pound = 453.6		0.4536	3.108 × 10 ⁻²	2.732 × 10 ²⁶	16	1	0.0005
1 ton = 9.072 × 10 ⁵		907.2	62.16	5.463 × 10 ²⁹	3.2 × 10 ⁴	2000	1

1 metric ton = 1000 kg

Density

Quantities in the colored areas are weight densities and, as such, are dimensionally different from mass densities. See the note for the mass table.

	slug/ft ³	KILOGRAM/ METER ³	g/cm ³	lb/ft ³	lb/in. ³
1 slug per foot ³ = 1		515.4	0.5154	32.17	1.862 × 10 ⁻²
1 KILOGRAM per METER ³ = 1.940 × 10 ⁻³		1	0.001	6.243 × 10 ⁻²	3.613 × 10 ⁻⁵
1 gram per centimeter ³ = 1.940		1000	1	62.43	3.613 × 10 ⁻²
1 pound per foot ³ = 3.108 × 10 ⁻²		16.02	16.02 × 10 ⁻²	1	5.787 × 10 ⁻⁴
1 pound per inch ³ = 53.71		2.768 × 10 ⁴	27.68	1728	1

Time

	y	d	h	min	SECOND
1 year = 1		365.25	8.766 × 10 ³	5.259 × 10 ⁵	3.156 × 10 ⁷
1 day = 2.738 × 10 ⁻³		1	24	1440	8.640 × 10 ⁴
1 hour = 1.141 × 10 ⁻⁴		4.167 × 10 ⁻²	1	60	3600
1 minute = 1.901 × 10 ⁻⁶		6.944 × 10 ⁻⁴	1.667 × 10 ⁻²	1	60
1 SECOND = 3.169 × 10 ⁻⁸		1.157 × 10 ⁻⁵	2.778 × 10 ⁻⁴	1.667 × 10 ⁻²	1

Speed

	ft/s	km/h	METER/SECOND	mi/h	cm/s
1 foot per second = 1		1.097	0.3048	0.6818	30.48
1 kilometer per hour = 0.9113		1	0.2778	0.6214	27.78
1 METER per SECOND = 3.281		3.6	1	2.237	100
1 mile per hour = 1.467		1.609	0.4470	1	44.70
1 centimeter per second = 3.281×10^{-2}		3.6×10^{-2}	0.01	2.237×10^{-2}	1

1 knot = 1 nautical mi/h = 1.688 ft/s 1 mi/min = 88.00 ft/s = 60.00 mi/h

Force

Force units in the colored areas are now little used. To clarify: 1 gram-force (= 1 gf) is the force of gravity that would act on an object whose mass is 1 gram at a location where g has the standard value of 9.80665 m/s^2 .

	dyne	NEWTON	lb	pdl	gf	kgf
1 dyne = 1		10^{-5}	2.248×10^{-6}	7.233×10^{-5}	1.020×10^{-3}	1.020×10^{-6}
1 NEWTON = 10^5		1	0.2248	7.233	102.0	0.1020
1 pound = 4.448×10^5		4.448	1	32.17	453.6	0.4536
1 poundal = 1.383×10^4		0.1383	3.108×10^{-2}	1	14.10	1.410×10^2
1 gram-force = 980.7		9.807×10^{-3}	2.205×10^{-3}	7.093×10^{-2}	1	0.001
1 kilogram-force = 9.807×10^5		9.807	2.205	70.93	1000	1

1 ton = 2000 lb

Pressure

	atm	dyne/cm ²	inch of water	cm Hg	PASCAL	lb/in. ²	lb/ft ²
1 atmosphere = 1		1.013×10^6	406.8	76	1.013×10^5	14.70	2116
1 dyne per centimeter ² = 9.869×10^{-7}		1	4.015×10^{-4}	7.501×10^{-5}	0.1	1.405×10^{-5}	2.089×10^{-3}
1 inch of water ^a at 4°C = 2.458×10^{-3}		2491	1	0.1868	249.1	3.613×10^{-2}	5.202
1 centimeter of mercury ^a at 0°C = 1.316×10^{-2}		1.333×10^4	5.353	1	1333	0.1934	27.85
1 PASCAL = 9.869×10^{-6}		10	4.015×10^{-3}	7.501×10^{-4}	1	1.450×10^{-4}	2.089×10^{-2}
1 pound per inch ² = 6.805×10^{-2}		6.895×10^4	27.68	5.171	6.895×10^3	1	144
1 pound per foot ² = 4.725×10^{-4}		478.8	0.1922	3.591×10^{-2}	47.88	6.944×10^{-3}	1

^aWhere the acceleration of gravity has the standard value of 9.80665 m/s^2 .

1 bar = 10^6 dyne/cm^2 = 0.1 MPa

1 millibar = 10^3 dyne/cm^2 = 10^2 Pa

1 torr = 1 mm Hg

A-8 APPENDIX D CONVERSION FACTORS

Energy, Work, Heat

Quantities in the colored areas are not energy units but are included for convenience. They arise from the relativistic mass-energy equivalence formula $E = mc^2$ and represent the energy released if a kilogram or unified atomic mass unit (u) is completely converted to energy (bottom two rows) or the mass that would be completely converted to one unit of energy (rightmost two columns).

	Btu	erg	ft · lb	hp · h	JOULE	cal	kW · h	eV	MeV	kg	u
1 British thermal unit = 1		1.055 $\times 10^{10}$	777.9	3.929 $\times 10^{-4}$	1055	252.0	2.930 $\times 10^{-4}$	6.585 $\times 10^{21}$	6.585 $\times 10^{15}$	1.174 $\times 10^{-14}$	7.070 $\times 10^{12}$
1 erg =	9.481 $\times 10^{-11}$	1	7.376 $\times 10^{-8}$	3.725 $\times 10^{-14}$	10 ⁻⁷	2.389 $\times 10^{-8}$	2.778 $\times 10^{-14}$	6.242 $\times 10^{11}$	6.242 $\times 10^5$	1.113 $\times 10^{-24}$	670.2
1 foot-pound =	1.285 $\times 10^{-3}$	1.356 $\times 10^7$	1	5.051 $\times 10^{-7}$	1.356	0.3238	3.766 $\times 10^{-7}$	8.464 $\times 10^{18}$	8.464 $\times 10^{12}$	1.509 $\times 10^{-17}$	9.037 $\times 10^9$
1 horsepower-hour =	2545	2.685 $\times 10^{13}$	1.980 $\times 10^6$	1	2.685 $\times 10^6$	6.413 $\times 10^5$	0.7457	1.676 $\times 10^{25}$	1.676 $\times 10^{19}$	2.988 $\times 10^{-11}$	1.799 $\times 10^{16}$
1 JOULE =	9.481 $\times 10^{-4}$	10 ⁷	0.7376	3.725 $\times 10^{-7}$	1	0.2389	2.778 $\times 10^{-7}$	6.242 $\times 10^{18}$	6.242 $\times 10^{12}$	1.113 $\times 10^{-17}$	6.702 $\times 10^9$
1 calorie =	3.968 $\times 10^{-3}$	4.1868 $\times 10^7$	3.088	1.560 $\times 10^{-6}$	4.1868	1	1.163 $\times 10^{-6}$	2.613 $\times 10^{19}$	2.613 $\times 10^{13}$	4.660 $\times 10^{-17}$	2.806 $\times 10^{10}$
1 kilowatt-hour =	3413	3.600 $\times 10^{13}$	2.655 $\times 10^6$	1.341 $\times 10^6$	3.600 $\times 10^6$	8.600 $\times 10^5$	1	2.247 $\times 10^{25}$	2.247 $\times 10^{19}$	4.007 $\times 10^{-11}$	2.413 $\times 10^{16}$
1 electron-volt =	1.519 $\times 10^{-22}$	1.602 $\times 10^{-12}$	1.182 $\times 10^{-19}$	5.967 $\times 10^{-26}$	1.602 $\times 10^{-19}$	3.827 $\times 10^{-20}$	4.450 $\times 10^{-26}$	1	10 ⁻⁶	1.783 $\times 10^{-36}$	1.074 $\times 10^{-9}$
1 million electron-volts =	1.519 $\times 10^{-16}$	1.602 $\times 10^{-6}$	1.182 $\times 10^{-13}$	5.967 $\times 10^{-20}$	1.602 $\times 10^{-13}$	3.827 $\times 10^{-14}$	4.450 $\times 10^{-20}$	10 ⁻⁶	1	1.783 $\times 10^{-30}$	1.074 $\times 10^{-3}$
1 kilogram =	8.521 $\times 10^{13}$	8.987 $\times 10^{23}$	6.629 $\times 10^{16}$	3.348 $\times 10^{10}$	8.987 $\times 10^{16}$	2.146 $\times 10^{16}$	2.497 $\times 10^{10}$	5.610 $\times 10^{35}$	5.610 $\times 10^{29}$	1	6.022 $\times 10^{26}$
1 unified atomic mass unit =	1.415 $\times 10^{-13}$	1.492 $\times 10^{-3}$	1.101 $\times 10^{-10}$	5.559 $\times 10^{-17}$	1.492 $\times 10^{-10}$	3.564 $\times 10^{-11}$	4.146 $\times 10^{-17}$	9.320 $\times 10^8$	932.0	1.661 $\times 10^{-27}$	1

Power

	Btu/h	ft · lb/s	hp	cal/s	kW	WATT
1 British thermal unit per hour = 1		0.2161	3.929 $\times 10^{-4}$	6.998 $\times 10^{-2}$	2.930 $\times 10^{-4}$	0.2930
1 foot-pound per second = 4.628		1	1.818 $\times 10^{-3}$	0.3239	1.356 $\times 10^{-3}$	1.356
1 horsepower = 2545		550	1	178.1	0.7457	745.7
1 calorie per second = 14.29		3.088	5.615 $\times 10^{-3}$	1	4.186 $\times 10^{-3}$	4.186
1 kilowatt = 3413		737.6	1.341	238.9	1	1000
1 WATT = 3.413		0.7376	1.341 $\times 10^{-3}$	0.2389	0.001	1

Magnetic Field

	gauss	TESLA	milligauss
1 gauss = 1		10 ⁻⁴	1000
1 TESLA = 10 ⁴		1	10 ⁷
1 milligauss = 0.001		10 ⁻⁷	1

1 tesla = 1 weber/meter²

Magnetic Flux

	maxwell	WEBER
1 maxwell = 1		10 ⁻⁸
1 WEBER = 10 ⁸		1

Mathematical Formulas

Geometry

Circle of radius r : circumference = $2\pi r$; area = πr^2 .

Sphere of radius r : area = $4\pi r^2$; volume = $\frac{4}{3}\pi r^3$.

Right circular cylinder of radius r and height h :
area = $2\pi r^2 + 2\pi rh$; volume = $\pi r^2 h$.

Triangle of base a and altitude h : area = $\frac{1}{2}ah$.

Quadratic Formula


If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Trigonometric Functions of Angle θ

$$\sin \theta = \frac{y}{r} \quad \cos \theta = \frac{x}{r}$$


$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

$$\sec \theta = \frac{r}{x} \quad \csc \theta = \frac{r}{y}$$



Pythagorean Theorem

In this right triangle,
 $a^2 + b^2 = c^2$



Triangles

Angles are A, B, C


Opposite sides are a, b, c

Angles $A + B + C = 180^\circ$

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Exterior angle $D = A + C$



Mathematical Signs and Symbols

= equals

≈ equals approximately

~ is the order of magnitude of

≠ is not equal to

≡ is identical to, is defined as

> is greater than (≫ is much greater than)

< is less than (≪ is much less than)

≥ is greater than or equal to (or, is no less than)

≤ is less than or equal to (or, is no more than)

± plus or minus

∝ is proportional to

Σ the sum of

x_{avg} the average value of x

Trigonometric Identities

$$\sin(90^\circ - \theta) = \cos \theta$$

$$\cos(90^\circ - \theta) = \sin \theta$$

$$\sin \theta / \cos \theta = \tan \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\sec^2 \theta - \tan^2 \theta = 1$$

$$\csc^2 \theta - \cot^2 \theta = 1$$

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$$

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

$$\sin \alpha \pm \sin \beta = 2 \sin \frac{1}{2}(\alpha \pm \beta) \cos \frac{1}{2}(\alpha \mp \beta)$$

$$\cos \alpha + \cos \beta = 2 \cos \frac{1}{2}(\alpha + \beta) \cos \frac{1}{2}(\alpha - \beta)$$

$$\cos \alpha - \cos \beta = -2 \sin \frac{1}{2}(\alpha + \beta) \sin \frac{1}{2}(\alpha - \beta)$$

Binomial Theorem

$$(1 + x)^n = 1 + \frac{nx}{1!} + \frac{n(n-1)x^2}{2!} + \cdots \quad (x^2 < 1)$$

Exponential Expansion

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

Logarithmic Expansion

$$\ln(1 + x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \cdots \quad (|x| < 1)$$

Trigonometric Expansions (θ in radians)

$$\sin \theta = \theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \cdots$$

$$\cos \theta = 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \cdots$$

$$\tan \theta = \theta + \frac{\theta^3}{3} + \frac{2\theta^5}{15} + \cdots$$

Cramer's Rule

Two simultaneous equations in unknowns x and y ,

$$a_1x + b_1y = c_1 \quad \text{and} \quad a_2x + b_2y = c_2,$$

have the solutions

$$x = \frac{\begin{vmatrix} c_1 & b_1 \\ c_2 & b_2 \end{vmatrix}}{\begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}} = \frac{c_1b_2 - c_2b_1}{a_1b_2 - a_2b_1}$$

and

$$y = \frac{\begin{vmatrix} a_1 & c_1 \\ a_2 & c_2 \end{vmatrix}}{\begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}} = \frac{a_1c_2 - a_2c_1}{a_1b_2 - a_2b_1}.$$

Products of Vectors

Let $\hat{i}$, $\hat{j}$, and $\hat{k}$ be unit vectors in the x , y , and z directions. Then

$$\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1, \quad \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0,$$

$$\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = 0,$$

$$\hat{i} \times \hat{j} = \hat{k}, \quad \hat{j} \times \hat{k} = \hat{i}, \quad \hat{k} \times \hat{i} = \hat{j}$$

Any vector $\vec{a}$ with components a_x , a_y , and a_z along the x , y , and z axes can be written as

$$\vec{a} = a_x\hat{i} + a_y\hat{j} + a_z\hat{k}.$$

Let $\vec{a}$, $\vec{b}$, and $\vec{c}$ be arbitrary vectors with magnitudes a , b , and c . Then

$$\vec{a} \times (\vec{b} + \vec{c}) = (\vec{a} \times \vec{b}) + (\vec{a} \times \vec{c})$$

$$(s\vec{a}) \times \vec{b} = \vec{a} \times (s\vec{b}) = s(\vec{a} \times \vec{b}) \quad (s = \text{a scalar}).$$

Let θ be the smaller of the two angles between $\vec{a}$ and $\vec{b}$. Then

$$\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a} = a_xb_x + a_yb_y + a_zb_z = ab \cos \theta$$

$$\vec{a} \times \vec{b} = -\vec{b} \times \vec{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix}$$

$$= \hat{i} \begin{vmatrix} a_y & a_z \\ b_y & b_z \end{vmatrix} - \hat{j} \begin{vmatrix} a_x & a_z \\ b_x & b_z \end{vmatrix} + \hat{k} \begin{vmatrix} a_x & a_y \\ b_x & b_y \end{vmatrix}$$

$$= (a_yb_z - b_ya_z)\hat{i} + (a_zb_x - b_za_x)\hat{j} + (a_xb_y - b_xa_y)\hat{k}$$

$$|\vec{a} \times \vec{b}| = ab \sin \theta$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{b} \cdot (\vec{c} \times \vec{a}) = \vec{c} \cdot (\vec{a} \times \vec{b})$$

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c}$$

Derivatives and Integrals

In what follows, the letters u and v stand for any functions of x , and a and m are constants. To each of the indefinite integrals should be added an arbitrary constant of integration. The *Handbook of Chemistry and Physics* (CRC Press Inc.) gives a more extensive tabulation.

$$1. \frac{dx}{dx} = 1$$

$$2. \frac{d}{dx}(au) = a \frac{du}{dx}$$

$$3. \frac{d}{dx}(u + v) = \frac{du}{dx} + \frac{dv}{dx}$$

$$4. \frac{d}{dx} x^m = mx^{m-1}$$

$$5. \frac{d}{dx} \ln x = \frac{1}{x}$$

$$6. \frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$7. \frac{d}{dx} e^x = e^x$$

$$8. \frac{d}{dx} \sin x = \cos x$$

$$9. \frac{d}{dx} \cos x = -\sin x$$

$$10. \frac{d}{dx} \tan x = \sec^2 x$$

$$11. \frac{d}{dx} \cot x = -\csc^2 x$$

$$12. \frac{d}{dx} \sec x = \tan x \sec x$$

$$13. \frac{d}{dx} \csc x = -\cot x \csc x$$

$$14. \frac{d}{dx} e^u = e^u \frac{du}{dx}$$

$$15. \frac{d}{dx} \sin u = \cos u \frac{du}{dx}$$

$$16. \frac{d}{dx} \cos u = -\sin u \frac{du}{dx}$$

$$1. \int dx = x$$

$$2. \int au \, dx = a \int u \, dx$$

$$3. \int (u + v) \, dx = \int u \, dx + \int v \, dx$$

$$4. \int x^m \, dx = \frac{x^{m+1}}{m+1} \quad (m \neq -1)$$

$$5. \int \frac{dx}{x} = \ln |x|$$

$$6. \int u \frac{dv}{dx} \, dx = uv - \int v \frac{du}{dx} \, dx$$

$$7. \int e^x \, dx = e^x$$

$$8. \int \sin x \, dx = -\cos x$$

$$9. \int \cos x \, dx = \sin x$$

$$10. \int \tan x \, dx = \ln |\sec x|$$

$$11. \int \sin^2 x \, dx = \frac{1}{2}x - \frac{1}{4}\sin 2x$$

$$12. \int e^{-ax} \, dx = -\frac{1}{a} e^{-ax}$$

$$13. \int x e^{-ax} \, dx = -\frac{1}{a^2} (ax + 1) e^{-ax}$$

$$14. \int x^2 e^{-ax} \, dx = -\frac{1}{a^3} (a^2 x^2 + 2ax + 2) e^{-ax}$$

$$15. \int_0^\infty x^n e^{-ax} \, dx = \frac{n!}{a^{n+1}}$$

$$16. \int_0^\infty x^{2n} e^{-ax^2} \, dx = \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^{n+1} a^n} \sqrt{\frac{\pi}{a}}$$

$$17. \int \frac{dx}{\sqrt{x^2 + a^2}} = \ln(x + \sqrt{x^2 + a^2})$$

$$18. \int \frac{x \, dx}{(x^2 + a^2)^{3/2}} = -\frac{1}{(x^2 + a^2)^{1/2}}$$

$$19. \int \frac{dx}{(x^2 + a^2)^{3/2}} = \frac{x}{a^2(x^2 + a^2)^{1/2}}$$

$$20. \int_0^\infty x^{2n+1} e^{-ax^2} \, dx = \frac{n!}{2a^{n+1}} \quad (a > 0)$$

$$21. \int \frac{x \, dx}{x + d} = x - d \ln(x + d)$$

Properties of the Elements

All physical properties are for a pressure of 1 atm unless otherwise specified.

Element	Symbol	Atomic Number Z	Molar Mass, g/mol	Density, g/cm ³ at 20°C	Melting Point, °C	Boiling Point, °C	Specific Heat, J/(g · °C) at 25°C
Actinium	Ac	89	(227)	10.06	1323	(3473)	0.092
Aluminum	Al	13	26.9815	2.699	660	2450	0.900
Americium	Am	95	(243)	13.67	1541	—	—
Antimony	Sb	51	121.75	6.691	630.5	1380	0.205
Argon	Ar	18	39.948	1.6626×10^{-3}	-189.4	-185.8	0.523
Arsenic	As	33	74.9216	5.78	817 (28 atm)	613	0.331
Astatine	At	85	(210)	—	(302)	—	—
Barium	Ba	56	137.34	3.594	729	1640	0.205
Berkelium	Bk	97	(247)	14.79	—	—	—
Beryllium	Be	4	9.0122	1.848	1287	2770	1.83
Bismuth	Bi	83	208.980	9.747	271.37	1560	0.122
Bohrium	Bh	107	262.12	—	—	—	—
Boron	B	5	10.811	2.34	2030	—	1.11
Bromine	Br	35	79.909	3.12 (liquid)	-7.2	58	0.293
Cadmium	Cd	48	112.40	8.65	321.03	765	0.226
Calcium	Ca	20	40.08	1.55	838	1440	0.624
Californium	Cf	98	(251)	—	—	—	—
Carbon	C	6	12.01115	2.26	3727	4830	0.691
Cerium	Ce	58	140.12	6.768	804	3470	0.188
Cesium	Cs	55	132.905	1.873	28.40	690	0.243
Chlorine	Cl	17	35.453	3.214×10^{-3} (0°C)	-101	-34.7	0.486
Chromium	Cr	24	51.996	7.19	1857	2665	0.448
Cobalt	Co	27	58.9332	8.85	1495	2900	0.423
Copernicium	Cp	112	(285)	—	—	—	—
Copper	Cu	29	63.54	8.96	1083.40	2595	0.385
Curium	Cm	96	(247)	13.3	—	—	—
Darmstadtium	Ds	110	(271)	—	—	—	—
Dubnium	Db	105	262.114	—	—	—	—
Dysprosium	Dy	66	162.50	8.55	1409	2330	0.172
Einsteinium	Es	99	(254)	—	—	—	—
Erbium	Er	68	167.26	9.15	1522	2630	0.167
Europium	Eu	63	151.96	5.243	817	1490	0.163
Fermium	Fm	100	(237)	—	—	—	—
Fluorine	F	9	18.9984	1.696×10^{-3} (0°C)	-219.6	-188.2	0.753
Francium	Fr	87	(223)	—	(27)	—	—
Gadolinium	Gd	64	157.25	7.90	1312	2730	0.234
Gallium	Ga	31	69.72	5.907	29.75	2237	0.377
Germanium	Ge	32	72.59	5.323	937.25	2830	0.322
Gold	Au	79	196.967	19.32	1064.43	2970	0.131

Element	Symbol	Atomic Number Z	Molar Mass, g/mol	Density, g/cm ³ at 20°C	Melting Point, °C	Boiling Point, °C	Specific Heat, J/(g · °C) at 25°C
Hafnium	Hf	72	178.49	13.31	2227	5400	0.144
Hassium	Hs	108	(265)	—	—	—	—
Helium	He	2	4.0026	0.1664×10^{-3}	-269.7	-268.9	5.23
Holmium	Ho	67	164.930	8.79	1470	2330	0.165
Hydrogen	H	1	1.00797	0.08375×10^{-3}	-259.19	-252.7	14.4
Indium	In	49	114.82	7.31	156.634	2000	0.233
Iodine	I	53	126.9044	4.93	113.7	183	0.218
Iridium	Ir	77	192.2	22.5	2447	(5300)	0.130
Iron	Fe	26	55.847	7.874	1536.5	3000	0.447
Krypton	Kr	36	83.80	3.488×10^{-3}	-157.37	-152	0.247
Lanthanum	La	57	138.91	6.189	920	3470	0.195
Lawrencium	Lr	103	(257)	—	—	—	—
Lead	Pb	82	207.19	11.35	327.45	1725	0.129
Lithium	Li	3	6.939	0.534	180.55	1300	3.58
Lutetium	Lu	71	174.97	9.849	1663	1930	0.155
Magnesium	Mg	12	24.312	1.738	650	1107	1.03
Manganese	Mn	25	54.9380	7.44	1244	2150	0.481
Meitnerium	Mt	109	(266)	—	—	—	—
Mendelevium	Md	101	(256)	—	—	—	—
Mercury	Hg	80	200.59	13.55	-38.87	357	0.138
Molybdenum	Mo	42	95.94	10.22	2617	5560	0.251
Neodymium	Nd	60	144.24	7.007	1016	3180	0.188
Neon	Ne	10	20.183	0.8387×10^{-3}	-248.597	-246.0	1.03
Neptunium	Np	93	(237)	20.25	637	—	1.26
Nickel	Ni	28	58.71	8.902	1453	2730	0.444
Niobium	Nb	41	92.906	8.57	2468	4927	0.264
Nitrogen	N	7	14.0067	1.1649×10^{-3}	-210	-195.8	1.03
Nobelium	No	102	(255)	—	—	—	—
Osmium	Os	76	190.2	22.59	3027	5500	0.130
Oxygen	O	8	15.9994	1.3318×10^{-3}	-218.80	-183.0	0.913
Palladium	Pd	46	106.4	12.02	1552	3980	0.243
Phosphorus	P	15	30.9738	1.83	44.25	280	0.741
Platinum	Pt	78	195.09	21.45	1769	4530	0.134
Plutonium	Pu	94	(244)	19.8	640	3235	0.130
Polonium	Po	84	(210)	9.32	254	—	—
Potassium	K	19	39.102	0.862	63.20	760	0.758
Praseodymium	Pr	59	140.907	6.773	931	3020	0.197
Promethium	Pm	61	(145)	7.22	(1027)	—	—
Protactinium	Pa	91	(231)	15.37 (estimated)	(1230)	—	—
Radium	Ra	88	(226)	5.0	700	—	—
Radon	Rn	86	(222)	9.96×10^{-3} (0°C)	(-71)	-61.8	0.092
Rhenium	Re	75	186.2	21.02	3180	5900	0.134
Rhodium	Rh	45	102.905	12.41	1963	4500	0.243
Roentgenium	Rg	111	(280)	—	—	—	—
Rubidium	Rb	37	85.47	1.532	39.49	688	0.364
Ruthenium	Ru	44	101.107	12.37	2250	4900	0.239
Rutherfordium	Rf	104	261.11	—	—	—	—

A-14 APPENDIX F PROPERTIES OF THE ELEMENTS

Element	Symbol	Atomic Number Z	Molar Mass, g/mol	Density, g/cm ³ at 20°C	Melting Point, °C	Boiling Point, °C	Specific Heat, J/(g · °C) at 25°C
Samarium	Sm	62	150.35	7.52	1072	1630	0.197
Scandium	Sc	21	44.956	2.99	1539	2730	0.569
Seaborgium	Sg	106	263.118	—	—	—	—
Selenium	Se	34	78.96	4.79	221	685	0.318
Silicon	Si	14	28.086	2.33	1412	2680	0.712
Silver	Ag	47	107.870	10.49	960.8	2210	0.234
Sodium	Na	11	22.9898	0.9712	97.85	892	1.23
Strontium	Sr	38	87.62	2.54	768	1380	0.737
Sulfur	S	16	32.064	2.07	119.0	444.6	0.707
Tantalum	Ta	73	180.948	16.6	3014	5425	0.138
Technetium	Tc	43	(99)	11.46	2200	—	0.209
Tellurium	Te	52	127.60	6.24	449.5	990	0.201
Terbium	Tb	65	158.924	8.229	1357	2530	0.180
Thallium	Tl	81	204.37	11.85	304	1457	0.130
Thorium	Th	90	(232)	11.72	1755	(3850)	0.117
Thulium	Tm	69	168.934	9.32	1545	1720	0.159
Tin	Sn	50	118.69	7.2984	231.868	2270	0.226
Titanium	Ti	22	47.90	4.54	1670	3260	0.523
Tungsten	W	74	183.85	19.3	3380	5930	0.134
Unnamed	Uut	113	(284)	—	—	—	—
Unnamed	Unq	114	(289)	—	—	—	—
Unnamed	Uup	115	(288)	—	—	—	—
Unnamed	Uuh	116	(293)	—	—	—	—
Unnamed	Uus	117	—	—	—	—	—
Unnamed	Uuo	118	(294)	—	—	—	—
Uranium	U	92	(238)	18.95	1132	3818	0.117
Vanadium	V	23	50.942	6.11	1902	3400	0.490
Xenon	Xe	54	131.30	5.495×10^{-3}	−111.79	−108	0.159
Ytterbium	Yb	70	173.04	6.965	824	1530	0.155
Yttrium	Y	39	88.905	4.469	1526	3030	0.297
Zinc	Zn	30	65.37	7.133	419.58	906	0.389
Zirconium	Zr	40	91.22	6.506	1852	3580	0.276

The values in parentheses in the column of molar masses are the mass numbers of the longest-lived isotopes of those elements that are radioactive. Melting points and boiling points in parentheses are uncertain.

The data for gases are valid only when these are in their usual molecular state, such as H₂, He, O₂, Ne, etc. The specific heats of the gases are the values at constant pressure.

Source: Adapted from J. Emsley, *The Elements*, 3rd ed., 1998, Clarendon Press, Oxford. See also www.webelements.com for the latest values and newest elements.

Periodic Table of the Elements

CHAPTER 1

P 1. (a) 4.00×10^4 km; (b) 5.10×10^8 km²; (c) 1.08×10^{12} km³
 3. (a) 10^9 μ m; (b) 10^{-4} ; (c) 9.1×10^5 μ m 5. (a) 160 rods; (b) 40 chains
 7. 1.1×10^3 acre-feet 9. 1.9×10^{22} cm³ 11. (a) 1.43; (b) 0.864 13. (a) 495 s; (b) 141 s; (c) 198 s; (d) -245 s 15. 1.21×10^{12} μ s
 17. C, D, A, B, E; the important criterion is the consistency of the daily variation, not its magnitude 19. 5.2×10^6 m 21. 9.0×10^{49} atoms
 23. (a) 1×10^3 kg; (b) 158 kg/s 25. 1.9×10^5 kg 27. (a) 1.18×10^{-29} m³; (b) 0.282 nm 29. 1.75×10^3 kg 31. 1.43 kg/min
 33. (a) 293 U.S. bushels; (b) 3.81×10^3 U.S. bushels 35. (a) 22 pecks; (b) 5.5 Imperial bushels; (c) 200 L 37. 8×10^2 km
 39. (a) 18.8 gallons; (b) 22.5 gallons 41. 0.3 cord 43. 3.8 mg/s 45. (a) yes; (b) 8.6 universe seconds 47. 0.12 AU/min 49. (a) 3.88; (b) 7.65; (c) 156 ken³; (d) 1.19×10^3 m³ 51. (a) 3.9 m, 4.8 m; (b) 3.9×10^3 mm, 4.8×10^3 mm; (c) 2.2 m³, 4.2 m³ 53. (a) 4.9×10^{-6} pc; (b) 1.6×10^{-5} ly

CHAPTER 2

CP 1. b and c 2. (check the derivative dx/dt) (a) 1 and 4; (b) 2 and 3 3. (a) plus; (b) minus; (c) minus; (d) plus 4. 1 and 4 ($a = d^2x/dt^2$ must be constant) 5. (a) plus (upward displacement on y axis); (b) minus (downward displacement on y axis); (c) $a = -g = -9.8$ m/s²
Q 1. (a) negative; (b) positive; (c) yes; (d) positive; (e) constant 3. (a) all tie; (b) 4, tie of 1 and 2, then 3 5. (a) positive direction; (b) negative direction; (c) 3 and 5; (d) 2 and 6 tie, then 3 and 5 tie, then 1 and 4 tie (zero) 7. (a) D; (b) E 9. (a) 3, 2, 1; (b) 1, 2, 3; (c) all tie; (d) 1, 2, 3
P 1. 13 m 3. (a) +40 km/h; (b) 40 km/h 5. (a) 0; (b) -2 m; (c) 0; (d) 12 m; (e) +12 m; (f) +7 m/s 7. 60 km 9. 1.4 m 11. 128 km/h
 13. (a) 73 km/h; (b) 68 km/h; (c) 70 km/h; (d) 0 15. (a) -6 m/s; (b) -x direction; (c) 6 m/s; (d) decreasing; (e) 2 s; (f) no 17. (a) 28.5 cm/s; (b) 18.0 cm/s; (c) 40.5 cm/s; (d) 28.1 cm/s; (e) 30.3 cm/s 19. -20 m/s²
 21. (a) 1.10 m/s; (b) 6.11 mm/s²; (c) 1.47 m/s; (d) 6.11 mm/s² 23. 1.62×10^{15} m/s² 25. (a) 30 s; (b) 300 m 27. (a) +1.6 m/s; (b) +18 m/s 29. (a) 10.6 m; (b) 41.5 s 31. (a) 3.1×10^6 s; (b) 4.6×10^{13} m 33. (a) 3.56 m/s²; (b) 8.43 m/s 35. 0.90 m/s² 37. (a) 4.0 m/s²; (b) +x 39. (a) -2.5 m/s²; (b) 1; (d) 0; (e) 2 41. 40 m 43. (a) 0.994 m/s² 45. (a) 31 m/s; (b) 6.4 s 47. (a) 29.4 m; (b) 2.45 s 49. (a) 5.4 s; (b) 41 m/s 51. (a) 20 m; (b) 59 m 53. 4.0 m/s 55. (a) 857 m/s²; (b) up 57. (a) 1.26×10^3 m/s²; (b) up 59. (a) 89 cm; (b) 22 cm 61. 20.4 m 63. 2.34 m 65. (a) 2.25 m/s; (b) 3.90 m/s 67. 0.56 m/s 69. 100 m 71. (a) 2.00 s; (b) 12 cm; (c) -9.00 cm/s²; (d) right; (e) left; (f) 3.46 s 73. (a) 82 m; (b) 19 m/s 75. (a) 0.74 s; (b) 6.2 m/s² 77. (a) 3.1 m/s²; (b) 45 m; (c) 13 s 79. 17 m/s 81. +47 m/s 83. (a) 1.23 cm; (b) 4 times; (c) 9 times; (d) 16 times; (e) 25 times 85. 25 km/h 87. 1.2 h 89. 4H 91. (a) 3.2 s; (b) 1.3 s 93. (a) 8.85 m/s; (b) 1.00 m 95. (a) 2.0 m/s²; (b) 12 m/s; (c) 45 m 97. (a) 48.5 m/s; (b) 4.95 s; (c) 34.3 m/s; (d) 3.50 s 99. 22.0 m/s 101. (a) $v = (v_0^2 + 2gh)^{0.5}$; (b) $t = [(v_0^2 + 2gh)^{0.5} - v_0] / g$; (c) same as (a); (d) $t = [(v_0^2 + 2gh)^{0.5} + v_0] / g$, greater

CHAPTER 3

CP 1. (a) 7 m ($\vec{a}$ and $\vec{b}$ are in same direction); (b) 1 m ($\vec{a}$ and $\vec{b}$ are in opposite directions) 2. c, d, f (components must be head to tail; $\vec{a}$ must extend from tail of one component to head of the other) 3. (a) +, +; (b) +, -; (c) +, + (draw vector from tail of $\vec{d}_1$ to head of $\vec{d}_2$)

4. (a) 90°; (b) 0° (vectors are parallel—same direction); (c) 180° (vectors are antiparallel—opposite directions) 5. (a) 0° or 180°; (b) 90°
Q 1. yes, when the vectors are in same direction 3. Either the sequence $\vec{d}_2, \vec{d}_1$ or the sequence $\vec{d}_2, \vec{d}_3, \vec{d}_3$ 5. all but (e) 7. (a) yes; (b) yes; (c) no 9. (a) +x for (1), +z for (2), +z for (3); (b) -x for (1), -z for (2), -z for (3)
P 1. (a) -2.5 m; (b) -6.9 m 3. (a) 47.2 m; (b) 122° 5. (a) 156 km; (b) 39.8° west of due north 7. (a) 6.42 m; (b) no; (c) yes; (d) yes; (e) a possible answer: $(4.30 \text{ m})\hat{i} + (3.70 \text{ m})\hat{j} + (3.00 \text{ m})\hat{k}$; (f) 7.96 m 9. (a) $(3.0 \text{ m})\hat{i} - (2.0 \text{ m})\hat{j} + (5.0 \text{ m})\hat{k}$; (b) $(5.0 \text{ m})\hat{i} - (4.0 \text{ m})\hat{j} - (3.0 \text{ m})\hat{k}$; (c) $(-5.0 \text{ m})\hat{i} + (4.0 \text{ m})\hat{j} + (3.0 \text{ m})\hat{k}$ 11. (a) $(-9.0 \text{ m})\hat{i} + (10 \text{ m})\hat{j}$; (b) 13 m; (c) 132° 13. 4.74 km 15. (a) 1.59 m; (b) 12.1 m; (c) 12.2 m; (d) 82.5° 17. (a) 38 m; (b) -37.5°; (c) 130 m; (d) 1.2°; (e) 62 m; (f) 130° 19. 5.39 m at 21.8° left of forward 21. (a) -70.0 cm; (b) 80.0 cm; (c) 141 cm; (d) -172° 23. 3.2 25. 2.6 km 27. (a) $8\hat{i} + 16\hat{j}$; (b) $2\hat{i} + 4\hat{j}$ 29. (a) 7.5 cm; (b) 90°; (c) 8.6 cm; (d) 48° 31. (a) $a\hat{i} + a\hat{j} + a\hat{k}$; (b) $-a\hat{i} + a\hat{j} + a\hat{k}$; (c) $a\hat{i} - a\hat{j} + a\hat{k}$; (d) $-a\hat{i} - a\hat{j} + a\hat{k}$; (e) 54.7°; (f) $3^{0.5}a$ 33. (a) 12; (b) +z; (c) 12; (d) -z; (e) 12; (f) +z 35. (a) -18.8 units; (b) 26.9 units, +z direction 37. (a) -21; (b) -9; (c) $5\hat{i} - 11\hat{j} - 9\hat{k}$ 39. 70.5° 41. 22° 43. (a) 3.00 m; (b) 0; (c) 3.46 m; (d) 2.00 m; (e) -5.00 m; (f) 8.66 m; (g) -6.67; (h) 4.33 45. (a) -83.4; (b) $(1.14 \times 10^3)\hat{k}$; (c) 1.14×10^3 , θ not defined, $\phi = 0^\circ$; (d) 90.0°; (e) $-5.14\hat{i} + 6.13\hat{j} + 3.00\hat{k}$; (f) 8.54, $\theta = 130^\circ$, $\phi = 69.4^\circ$ 47. (a) 140°; (b) 90.0°; (c) 99.1° 49. (a) 103 km; (b) 60.9° north of due west 51. (a) 27.8 m; (b) 13.4 m 53. (a) 30; (b) 52 55. (a) -2.83 m; (b) -2.83 m; (c) 5.00 m; (d) 0; (e) 3.00 m; (f) 5.20 m; (g) 5.17 m; (h) 2.37 m; (i) 5.69 m; (j) 25° north of due east; (k) 5.69 m; (l) 25° south of due west 57. 4.1 59. (a) $(9.19 \text{ m})\hat{i}' + (7.71 \text{ m})\hat{j}'$; (b) $(14.0 \text{ m})\hat{i}' + (3.41 \text{ m})\hat{j}'$ 61. (a) $11\hat{i} + 5.0\hat{j} - 7.0\hat{k}$; (b) 120°; (c) -4.9; (d) 7.3 63. (a) 3.0 m²; (b) 52 m³; (c) $(11 \text{ m}^2)\hat{i} + (9.0 \text{ m}^2)\hat{j} + (3.0 \text{ m}^2)\hat{k}$ 65. (a) $(-40\hat{i} - 20\hat{j} + 25\hat{k})$ m; (b) 45 m

CHAPTER 4

CP 1. (draw $\vec{v}$ tangent to path, tail on path) (a) first; (b) third 2. (take second derivative with respect to time) (1) and (3) a_x and a_y are both constant and thus $\vec{a}$ is constant; (2) and (4) a_y is constant but a_x is not, thus $\vec{a}$ is not 3. yes 4. (a) v_x constant; (b) v_y initially positive, decreases to zero, and then becomes progressively more negative; (c) $a_x = 0$ throughout; (d) $a_y = -g$ throughout 5. (a) $-(4 \text{ m/s})\hat{i}$; (b) $-(8 \text{ m/s}^2)\hat{j}$
Q 1. a and c tie, then b 3. decreases 5. a, b, c 7. (a) 0; (b) 350 km/h; (c) 350 km/h; (d) same (nothing changed about the vertical motion) 9. (a) all tie; (b) all tie; (c) 3, 2, 1; (d) 3, 2, 1 11. 2, then 1 and 4 tie, then 3 13. (a) yes; (b) no; (c) yes
P 1. (a) 6.2 m 3. $(-2.0 \text{ m})\hat{i} + (6.0 \text{ m})\hat{j} - (10 \text{ m})\hat{k}$ 5. (a) 7.59 km/h; (b) 22.5° east of due north 7. $(-0.70 \text{ m/s})\hat{i} + (1.4 \text{ m/s})\hat{j} - (0.40 \text{ m/s})\hat{k}$ 9. (a) 0.83 cm/s; (b) 0°; (c) 0.11 m/s; (d) -63° 11. (a) $(6.00 \text{ m})\hat{i} - (106 \text{ m})\hat{j}$; (b) $(19.0 \text{ m/s})\hat{i} - (224 \text{ m/s})\hat{j}$; (c) $(24.0 \text{ m/s}^2)\hat{i} - (336 \text{ m/s}^2)\hat{j}$; (d) -85.2° 13. (a) $(8 \text{ m/s}^2)\hat{i}\hat{j} + (1 \text{ m/s})\hat{k}$; (b) $(8 \text{ m/s}^2)\hat{j}$ 15. (a) $(-1.50 \text{ m/s})\hat{j}$; (b) $(4.50 \text{ m})\hat{i} - (2.25 \text{ m})\hat{j}$ 17. $(32 \text{ m/s})\hat{i}$ 19. (a) $(72.0 \text{ m})\hat{i} + (90.7 \text{ m})\hat{j}$; (b) 49.5° 21. (a) 18 cm; (b) 1.9 m 23. (a) 3.03 s; (b) 758 m; (c) 29.7 m/s 25. 43.1 m/s (155 km/h) 27. (a) 10.0 s; (b) 897 m 29. 78.5° 31. 3.35 m 33. (a) 202 m/s; (b) 806 m; (c) 161 m/s; (d) -171 m/s 35. 4.84 cm 37. (a) 1.60 m; (b) 6.86 m; (c) 2.86 m 39. (a) 32.3 m; (b) 21.9 m/s; (c) 40.4°; (d) below 41. 55.5° 43. (a) 11 m; (b) 23 m; (c) 17 m/s; (d) 63°

45. (a) ramp; (b) 5.82 m; (c) 31.0° 47. (a) yes; (b) 2.56 m 49. (a) 31° ; (b) 63° 51. (a) 2.3° ; (b) 1.4 m; (c) 18° 53. (a) 75.0 m; (b) 31.9 m/s; (c) 66.9° ; (d) 25.5 m 55. the third 57. (a) 7.32 m; (b) west; (c) north 59. (a) 12 s; (b) 4.1 m/s^2 ; (c) down; (d) 4.1 m/s^2 ; (e) up 61. (a) $1.3 \times 10^5 \text{ m/s}$; (b) $7.9 \times 10^5 \text{ m/s}^2$; (c) increase 63. 2.92 m 65. $(3.00 \text{ m/s}^2)\hat{i} + (6.00 \text{ m/s}^2)\hat{j}$ 67. 160 m/s^2 69. (a) 13 m/s^2 ; (b) eastward; (c) 13 m/s^2 ; (d) eastward 71. 1.67 73. (a) $(80 \text{ km/h})\hat{i} - (60 \text{ km/h})\hat{j}$; (b) 0° ; (c) answers do not change 75. 32 m/s 77. 60° 79. (a) 38 knots; (b) 1.5° east of due north; (c) 4.2 h; (d) 1.5° west of due south 81. (a) $(-32 \text{ km/h})\hat{i} - (46 \text{ km/h})\hat{j}$; (b) $[(2.5 \text{ km}) - (32 \text{ km/h})\hat{i}] + [(4.0 \text{ km}) - (46 \text{ km/h})\hat{j}]$; (c) 0.084 h; (d) $2 \times 10^2 \text{ m}$ 83. (a) -30° ; (b) 69 min; (c) 80 min; (d) 80 min; (e) 0° ; (f) 60 min 85. (a) 2.7 km; (b) 76° clockwise 87. (a) 44 m; (b) 13 m; (c) 8.9 m 89. (a) 45 m; (b) 22 m/s 91. (a) $2.6 \times 10^2 \text{ m/s}$; (b) 45 s; (c) increase 93. (a) 63 km; (b) 18° south of due east; (c) 0.70 km/h; (d) 18° south of due east; (e) 1.6 km/h; (f) 1.2 km/h; (g) 33° north of due east 95. (a) 1.5; (b) (36 m, 54 m) 97. (a) 62 ms; (b) $4.8 \times 10^2 \text{ m/s}$ 99. 2.64 m 101. (a) 2.5 m; (b) 0.82 m; (c) 9.8 m/s^2 ; (d) 9.8 m/s^2 103. (a) 6.79 km/h; (b) 6.96° 105. (a) 16 m/s; (b) 23° ; (c) above; (d) 27 m/s; (e) 57° ; (f) below 107. (a) 4.2 m, 45° ; (b) 5.5 m, 68° ; (c) 6.0 m, 90° ; (d) 4.2 m, 135° ; (e) 0.85 m/s, 135° ; (f) 0.94 m/s, 90° ; (g) 0.94 m/s, 180° ; (h) 0.30 m/s^2 , 180° ; (i) 0.30 m/s^2 , 270° 109. (a) $5.4 \times 10^{-13} \text{ m}$; (b) decrease 111. (a) 0.034 m/s^2 ; (b) 84 min 113. (a) 8.43 m; (b) -129° 115. (a) 2.00 ns; (b) 2.00 mm; (c) $1.00 \times 10^7 \text{ m/s}$; (d) $2.00 \times 10^6 \text{ m/s}$ 117. (a) 24 m/s; (b) 65° 119. 93° from the car's direction of motion

CHAPTER 5

CP 1. c, d, and e ($\vec{F}_1$ and $\vec{F}_2$ must be head to tail, $\vec{F}_{\text{net}}$ must be from tail of one of them to head of the other) 2. (a) and (b) 2 N, leftward (acceleration is zero in each situation) 3. (a) equal; (b) greater (acceleration is upward, thus net force on body must be upward) 4. (a) equal; (b) greater; (c) less 5. (a) increase; (b) yes; (c) same; (d) yes

Q 1. (a) 2, 3, 4; (b) 1, 3, 4; (c) 1, +y; 2, +x; 3, fourth quadrant; 4, third quadrant 3. increase 5. (a) 2 and 4; (b) 2 and 4 7. (a) M ; (b) M ; (c) M ; (d) $2M$; (e) $3M$ 9. (a) 20 kg; (b) 18 kg; (c) 10 kg; (d) all tie; (e) 3, 2, 1 11. (a) increases from initial value mg ; (b) decreases from mg to zero (after which the block moves up away from the floor)

P 1. 2.9 m/s^2 3. (a) 1.88 N; (b) 0.684 N; (c) $(1.88 \text{ N})\hat{i} + (0.684 \text{ N})\hat{j}$ 5. (a) $(0.86 \text{ m/s}^2)\hat{i} - (0.16 \text{ m/s}^2)\hat{j}$; (b) 0.88 m/s^2 ; (c) -11° 7. (a) $(-32.0 \text{ N})\hat{i} - (20.8 \text{ N})\hat{j}$; (b) 38.2 N; (c) -147° 9. (a) 8.37 N; (b) -133° ; (c) -125° 11. 9.0 m/s^2 13. (a) 4.0 kg; (b) 1.0 kg; (c) 4.0 kg; (d) 1.0 kg 15. (a) 108 N; (b) 108 N; (c) 108 N 17. (a) 42 N; (b) 72 N; (c) 4.9 m/s^2 19. $1.2 \times 10^5 \text{ N}$ 21. (a) 11.7 N; (b) -59.0° 23. (a) $(285 \text{ N})\hat{i} + (705 \text{ N})\hat{j}$; (b) $(285 \text{ N})\hat{i} - (115 \text{ N})\hat{j}$; (c) 307 N; (d) -22.0° ; (e) 3.67 m/s^2 ; (f) -22.0° 25. (a) 0.022 m/s^2 ; (b) $8.3 \times 10^4 \text{ km}$; (c) $1.9 \times 10^3 \text{ m/s}$ 27. 1.5 mm 29. (a) 494 N; (b) up; (c) 494 N; (d) down 31. (a) 1.18 m; (b) 0.674 s; (c) 3.50 m/s 33. $1.8 \times 10^4 \text{ N}$ 35. (a) 46.7° ; (b) 28.0° 37. (a) 0.62 m/s²; (b) 0.13 m/s²; (c) 2.6 m 39. (a) $2.2 \times 10^{-3} \text{ N}$; (b) $3.7 \times 10^{-3} \text{ N}$ 41. (a) 1.4 m/s^2 ; (b) 4.1 m/s 43. (a) 1.23 N; (b) 2.46 N; (c) 3.69 N; (d) 4.92 N; (e) 6.15 N; (f) 0.250 N 45. (a) 31.3 kN; (b) 24.3 kN 47. $6.4 \times 10^3 \text{ N}$ 49. (a) 2.18 m/s^2 ; (b) 116 N; (c) 21.0 m/s^2 51. (a) 3.6 m/s^2 ; (b) 17 N 53. (a) 0.970 m/s^2 ; (b) 11.6 N; (c) 34.9 N 55. (a) 1.1 N 57. (a) 0.735 m/s^2 ; (b) down; (c) 20.8 N 59. (a) 4.9 m/s^2 ; (b) 2.0 m/s^2 ; (c) up; (d) 120 N 61. $2M/(a + g)$ 63. (a) 8.0 m/s; (b) +x 65. (a) 0.653 m/s^3 ; (b) 0.896 m/s^3 ; (c) 6.50 s 67. 81.7 N 69. 2.4 N 71. 16 N 73. (a) 2.6 N; (b) 17° 75. (a) 0; (b) 0.83 m/s^2 ; (c) 0 77. (a) 0.74 m/s^2 ; (b) 7.3 m/s^2 79. (a) 11 N; (b) 2.2 kg; (c) 0; (d) 2.2 kg 81. 195 N 83. (a) 4.6 m/s^2 ; (b) 2.6 m/s^2 85. (a) rope breaks; (b) 1.6 m/s^2 87. (a) 65 N; (b) 49 N 89. (a) $4.6 \times 10^3 \text{ N}$; (b) $5.8 \times 10^3 \text{ N}$ 91. (a) $1.8 \times 10^2 \text{ N}$; (b) $6.4 \times$

10^2 N 93. (a) 44 N; (b) 78 N; (c) 54 N; (d) 152 N 95. (a) 4 kg; (b) 6.5 m/s^2 ; (c) 13 N

CHAPTER 6

CP 1. (a) zero (because there is no attempt at sliding); (b) 5 N; (c) no; (d) yes; (e) 8 N 2. ($\vec{a}$ is directed toward center of circular path) (a) $\vec{a}$ downward, $\vec{F}_N$ upward; (b) $\vec{a}$ and $\vec{F}_N$ upward Q 1. (a) decrease; (b) decrease; (c) increase; (d) increase; (e) increase 3. (a) same; (b) increases; (c) increases; (d) no 5. (a) upward; (b) horizontal, toward you; (c) no change; (d) increases; (e) increases 7. At first, $\vec{f}_s$ is directed up the ramp and its magnitude increases from $mg \sin \theta$ until it reaches $f_{s,\text{max}}$. Thereafter the force is kinetic friction directed up the ramp, with magnitude f_k (a constant value smaller than $f_{s,\text{max}}$). 9. 4, 3, then 1, 2, and 5 tie 11. (a) all tie; (b) all tie; (c) 2, 3, 1

P 1. 36 m 3. (a) $2.0 \times 10^2 \text{ N}$; (b) $1.2 \times 10^2 \text{ N}$ 5. (a) 6.0 N; (b) 3.6 N; (c) 3.1 N 7. (a) $1.9 \times 10^2 \text{ N}$; (b) 0.56 m/s^2 9. (a) 11 N; (b) 0.14 m/s^2 11. (a) $3.0 \times 10^2 \text{ N}$; (b) 1.3 m/s^2 13. (a) $1.3 \times 10^2 \text{ N}$; (b) no; (c) $1.1 \times 10^2 \text{ N}$; (d) 46 N; (e) 17 N 15. 2° 17. (a) $(17 \text{ N})\hat{i}$; (b) $(20 \text{ N})\hat{i}$; (c) $(15 \text{ N})\hat{i}$ 19. (a) no; (b) $(-12 \text{ N})\hat{i} + (5.0 \text{ N})\hat{j}$ 21. (a) 19° ; (b) 3.3 kN 23. 0.37 25. $1.0 \times 10^2 \text{ N}$ 27. (a) 0; (b) $(-3.9 \text{ m/s}^2)\hat{i}$; (c) $(-1.0 \text{ m/s}^2)\hat{i}$ 29. (a) 66 N; (b) 2.3 m/s^2 31. (a) 3.5 m/s^2 ; (b) 0.21 N 33. 9.9 s 35. $4.9 \times 10^2 \text{ N}$ 37. (a) $3.2 \times 10^2 \text{ km/h}$; (b) $6.5 \times 10^2 \text{ km/h}$; (c) no 39. 2.3 41. 0.60 43. 21 m 45. (a) light; (b) 778 N; (c) 223 N; (d) 1.11 kN 47. (a) 10 s; (b) $4.9 \times 10^2 \text{ N}$; (c) $1.1 \times 10^3 \text{ N}$ 49. $1.37 \times 10^3 \text{ N}$ 51. 2.2 km 53. 12° 55. $2.6 \times 10^3 \text{ N}$ 57. 1.81 m/s 59. (a) 8.74 N; (b) 37.9 N; (c) 6.45 m/s; (d) radially inward 61. (a) 27 N; (b) 3.0 m/s^2 63. (b) 240 N; (c) 0.60 65. (a) 69 km/h; (b) 139 km/h; (c) yes 67. $g(\sin \theta - 2^{0.5} \mu_k \cos \theta)$ 69. 3.4 m/s^2 71. (a) 35.3 N; (b) 39.7 N; (c) 320 N 73. (a) 7.5 m/s^2 ; (b) down; (c) 9.5 m/s^2 ; (d) down 75. (a) $3.0 \times 10^5 \text{ N}$; (b) 1.2° 77. 147 m/s 79. (a) 13 N; (b) 1.6 m/s^2 81. (a) 275 N; (b) 877 N 83. (a) 84.2 N; (b) 52.8 N; (c) 1.87 m/s^2 85. 3.4% 87. (a) $3.21 \times 10^3 \text{ N}$; (b) yes 89. (a) 222 N; (b) 334 N; (c) 311 N; (d) 311 N; (e) c, d 91. (a) $v_0^2/(4g \sin \theta)$; (b) no 93. (a) 0.34; (b) 0.24 95. (a) $\mu_k mg/(\sin \theta - \mu_k \cos \theta)$; (b) $\theta_0 = \tan^{-1} \mu_s$ 97. 0.18

CHAPTER 7

CP 1. (a) decrease; (b) same; (c) negative, zero 2. (a) positive; (b) negative; (c) zero 3. zero

Q 1. all tie 3. (a) positive; (b) negative; (c) negative 5. b (positive work), a (zero work), c (negative work), d (more negative work) 7. all tie 9. (a) A; (b) B

P 1. (a) $2.9 \times 10^7 \text{ m/s}$; (b) $2.1 \times 10^{-13} \text{ J}$ 3. (a) $5 \times 10^{14} \text{ J}$; (b) 0.1 megaton TNT; (c) 8 bombs 5. (a) 2.4 m/s; (b) 4.8 m/s 7. 0.96 J 9. 20 J 11. (a) 62.3° ; (b) 118° 13. (a) $1.7 \times 10^2 \text{ N}$; (b) $3.4 \times 10^2 \text{ m}$; (c) $-5.8 \times 10^4 \text{ J}$; (d) $3.4 \times 10^2 \text{ N}$; (e) $1.7 \times 10^2 \text{ m}$; (f) $-5.8 \times 10^4 \text{ J}$ 15. (a) 1.50 J; (b) increases 17. (a) 12 kJ; (b) -11 kJ ; (c) 1.1 kJ; (d) 5.4 m/s 19. 25 J 21. (a) $-3Mgd/4$; (b) Mgd ; (c) $Mgd/4$; (d) $(gd/2)^{0.5}$ 23. 4.41 J 25. (a) 25.9 kJ; (b) 2.45 N 27. (a) 7.2 J; (b) 7.2 J; (c) 0; (d) -25 J 29. (a) 0.90 J; (b) 2.1 J; (c) 0 31. (a) 6.6 m/s; (b) 4.7 m 33. (a) 0.12 m; (b) 0.36 J; (c) -0.36 J ; (d) 0.060 m; (e) 0.090 J 35. (a) 0; (b) 0 37. (a) 42 J; (b) 30 J; (c) 12 J; (d) 6.5 m/s, +x axis; (e) 5.5 m/s, +x axis; (f) 3.5 m/s, +x axis 39. 4.00 N/m 41. $5.3 \times 10^2 \text{ J}$ 43. (a) 0.83 J; (b) 2.5 J; (c) 4.2 J; (d) 5.0 W 45. $4.9 \times 10^2 \text{ W}$ 47. (a) $1.0 \times 10^2 \text{ J}$; (b) 8.4 W 49. $7.4 \times 10^2 \text{ W}$ 51. (a) 32.0 J; (b) 8.00 W; (c) 78.2° 53. (a) 1.20 J; (b) 1.10 m/s 55. (a) $1.8 \times 10^5 \text{ ft} \cdot \text{lb}$; (b) 0.55 hp 57. (a) 797 N; (b) 0; (c) -1.55 kJ ; (d) 0; (e) 1.55 kJ; (f) F varies during displacement 59. (a) 1×10^5 megatons TNT; (b) 1×10^7 bombs 61. -6 J 63. (a) 314 J; (b) -155 J ; (c) 0; (d) 158 J 65. (a) 98 N; (b) 4.0 cm; (c) 3.9 J; (d) -3.9 J 67. (a) 23 mm; (b) 45 N 69. 165 kW

71. -37 J 73. (a) 13 J ; (b) 13 J 75. 235 kW 77. (a) 6 J ; (b) 6.0 J
79. (a) 0.6 J ; (b) 0 ; (c) -0.6 J

CHAPTER 8

CP 1. no (consider round trip on the small loop) 2. 3, 1, 2 (see Eq. 8-6) 3. (a) all tie; (b) all tie 4. (a) CD, AB, BC (0) (check slope magnitudes); (b) positive direction of x 5. all tie
Q 1. (a) 3, 2, 1; (b) 1, 2, 3 3. (a) 12 J ; (b) -2 J 5. (a) increasing; (b) decreasing; (c) decreasing; (d) constant in AB and BC , decreasing in CD 7. $+30\text{ J}$ 9. 2, 1, 3
P 1. 89 N/cm 3. (a) 167 J ; (b) -167 J ; (c) 196 J ; (d) 29 J ; (e) 167 J ; (f) -167 J ; (g) 296 J ; (h) 129 J 5. (a) 4.31 mJ ; (b) -4.31 mJ ; (c) 4.31 mJ ; (d) -4.31 mJ ; (e) all increase 7. (a) 13.1 J ; (b) -13.1 J ; (c) 13.1 J ; (d) all increase 9. (a) 17.0 m/s ; (b) 26.5 m/s ; (c) 33.4 m/s ; (d) 56.7 m/s ; (e) all the same 11. (a) 2.08 m/s ; (b) 2.08 m/s ; (c) increase 13. (a) 0.98 J ; (b) -0.98 J ; (c) 3.1 N/cm 15. (a) $2.6 \times 10^2\text{ m}$; (b) same; (c) decrease 17. (a) 2.5 N ; (b) 0.31 N ; (c) 30 cm 19. (a) 784 N/m ; (b) 62.7 J ; (c) 62.7 J ; (d) 80.0 cm 21. (a) 8.35 m/s ; (b) 4.33 m/s ; (c) 7.45 m/s ; (d) both decrease 23. (a) 4.85 m/s ; (b) 2.42 m/s 25. $-3.2 \times 10^2\text{ J}$ 27. (a) no; (b) $9.3 \times 10^2\text{ N}$ 29. (a) 35 cm ; (b) 1.7 m/s 31. (a) 39.2 J ; (b) 39.2 J ; (c) 4.00 m 33. (a) 2.40 m/s ; (b) 4.19 m/s 35. (a) 39.6 cm ; (b) 3.64 cm 37. -18 mJ 39. (a) 2.1 m/s ; (b) 10 N ; (c) $+x$ direction; (d) 5.7 m ; (e) 30 N ; (f) $-x$ direction 41. (a) -3.7 J ; (c) 1.3 m ; (d) 9.1 m ; (e) 2.2 J ; (f) 4.0 m ; (g) $(4-x)e^{-x/4}$; (h) 4.0 m 43. (a) 5.6 J ; (b) 3.5 J 45. (a) 30.1 J ; (b) 30.1 J ; (c) 0.225 47. 0.53 J 49. (a) -2.9 kJ ; (b) $3.9 \times 10^2\text{ J}$; (c) $2.1 \times 10^2\text{ N}$ 51. (a) 1.5 MJ ; (b) 0.51 MJ ; (c) 1.0 MJ ; (d) 63 m/s 53. (a) 67 J ; (b) 67 J ; (c) 46 cm 55. (a) -0.90 J ; (b) 0.46 J ; (c) 1.0 m/s 57. 1.2 m 59. (a) 19.4 m ; (b) 19.0 m/s 61. (a) $1.5 \times 10^{-2}\text{ N}$; (b) $(3.8 \times 10^2)\text{g}$ 63. (a) 7.4 m/s ; (b) 90 cm ; (c) 2.8 m ; (d) 15 m 65. 20 cm 67. (a) 7.0 J ; (b) 22 J 69. 3.7 J 71. 4.33 m/s 73. 25 J 75. (a) 4.9 m/s ; (b) 4.5 N ; (c) 71° ; (d) same 77. (a) 4.8 N ; (b) $+x$ direction; (c) 1.5 m ; (d) 13.5 m ; (e) 3.5 m/s 79. (a) 24 kJ ; (b) $4.7 \times 10^2\text{ N}$ 81. (a) 5.00 J ; (b) 9.00 J ; (c) 11.0 J ; (d) 3.00 J ; (e) 12.0 J ; (f) 2.00 J ; (g) 13.0 J ; (h) 1.00 J ; (i) 13.0 J ; (j) 1.00 J ; (l) 11.0 J ; (m) 10.8 m ; (n) It returns to $x = 0$ and stops. 83. (a) 6.0 kJ ; (b) $6.0 \times 10^2\text{ W}$; (c) $3.0 \times 10^2\text{ W}$; (d) $9.0 \times 10^2\text{ W}$ 85. 880 MW 87. (a) $v_0 = (2gL)^{0.5}$; (b) $5mg$; (c) $-mgL$; (d) $-2mgL$ 89. (a) 109 J ; (b) 60.3 J ; (c) 68.2 J ; (d) 41.0 J 91. (a) 2.7 J ; (b) 1.8 J ; (c) 0.39 m 93. (a) 10 m ; (b) 49 N ; (c) 4.1 m ; (d) $1.2 \times 10^2\text{ N}$ 95. (a) 5.5 m/s ; (b) 5.4 m ; (c) same 97. 80 mJ 99. 24 W 101. -12 J 103. (a) 8.8 m/s ; (b) 2.6 kJ ; (c) 1.6 kW 105. (a) $7.4 \times 10^2\text{ J}$; (b) $2.4 \times 10^2\text{ J}$ 107. 15 J 109. (a) $2.35 \times 10^3\text{ J}$; (b) 352 J 111. 738 m 113. (a) -3.8 kJ ; (b) 31 kN 115. (a) 300 J ; (b) 93.8 J ; (c) 6.38 m 117. (a) 5.6 J ; (b) 12 J ; (c) 13 J 119. (a) 1.2 J ; (b) 11 m/s ; (c) no; (d) no 121. (a) $2.1 \times 10^6\text{ kg}$; (b) $(100 + 1.5t)^{0.5}\text{ m/s}$; (c) $(1.5 \times 10^6)/(100 + 1.5t)^{0.5}\text{ N}$; (d) 6.7 km

CHAPTER 9

CP 1. (a) origin; (b) fourth quadrant; (c) on y axis below origin; (d) origin; (e) third quadrant; (f) origin 2. (a)–(c) at the center of mass, still at the origin (their forces are internal to the system and cannot move the center of mass) 3. (Consider slopes and Eq. 9-23.) (a) 1, 3, and then 2 and 4 tie (zero force); (b) 3 4. (a) unchanged; (b) unchanged (see Eq. 9-32); (c) decrease (Eq. 9-35) 5. (a) zero; (b) positive (initial p_y down y ; final p_y up y); (c) positive direction of y 6. (No net external force; $\vec{P}$ conserved.) (a) 0; (b) no; (c) $-x$ 7. (a) $10\text{ kg} \cdot \text{m/s}$; (b) $14\text{ kg} \cdot \text{m/s}$; (c) $6\text{ kg} \cdot \text{m/s}$ 8. (a) $4\text{ kg} \cdot \text{m/s}$; (b) $8\text{ kg} \cdot \text{m/s}$; (c) 3 J 9. (a) $2\text{ kg} \cdot \text{m/s}$ (conserve momentum along x); (b) $3\text{ kg} \cdot \text{m/s}$ (conserve momentum along y)
Q 1. (a) 2 N , rightward; (b) 2 N , rightward; (c) greater than 2 N , rightward 3. b, c, a 5. (a) x yes, y no; (b) x yes, y no; (c) x no, y yes 7. (a) c , kinetic energy cannot be negative; d , total kinetic energy

cannot increase; (b) a ; (c) b 9. (a) one was stationary; (b) 2; (c) 5; (d) equal (pool player's result) 11. (a) C ; (b) B ; (c) 3
P 1. (a) -1.50 m ; (b) -1.43 m 3. (a) -6.5 cm ; (b) 8.3 cm ; (c) 1.4 cm 5. (a) -0.45 cm ; (b) -2.0 cm 7. (a) 0; (b) $3.13 \times 10^{-11}\text{ m}$ 9. (a) 28 cm ; (b) 2.3 m/s 11. $(-4.0\text{ m})\hat{i} + (4.0\text{ m})\hat{j}$ 13. 53 m 15. (a) $(2.35\hat{i} - 1.57\hat{j})\text{ m/s}^2$; (b) $(2.35\hat{i} - 1.57\hat{j})t\text{ m/s}$, with t in seconds; (d) straight, at downward angle 34° 17. 4.2 m 19. (a) $7.5 \times 10^4\text{ J}$; (b) $3.8 \times 10^4\text{ kg} \cdot \text{m/s}$; (c) 39° south of due east 21. (a) $5.0\text{ kg} \cdot \text{m/s}$; (b) $10\text{ kg} \cdot \text{m/s}$ 23. 1.0×10^3 to $1.2 \times 10^3\text{ kg} \cdot \text{m/s}$ 25. (a) $42\text{ N} \cdot \text{s}$; (b) 2.1 kN 27. (a) 67 m/s ; (b) $-x$; (c) 1.2 kN ; (d) $-x$ 29. 5 N 31. (a) $2.39 \times 10^3\text{ N} \cdot \text{s}$; (b) $4.78 \times 10^5\text{ N}$; (c) $1.76 \times 10^3\text{ N} \cdot \text{s}$; (d) $3.52 \times 10^5\text{ N}$ 33. (a) $5.86\text{ kg} \cdot \text{m/s}$; (b) 59.8° ; (c) 2.93 kN ; (d) 59.8° 35. $9.9 \times 10^2\text{ N}$ 37. (a) $9.0\text{ kg} \cdot \text{m/s}$; (b) 3.0 kN ; (c) 4.5 kN ; (d) 20 m/s 39. 3.0 mm/s 41. (a) $-(0.15\text{ m/s})\hat{i}$; (b) 0.18 m 43. 55 cm 45. (a) $(1.00\hat{i} - 0.167\hat{j})\text{ km/s}$; (b) 3.23 MJ 47. (a) 14 m/s ; (b) 45° 49. $3.1 \times 10^2\text{ m/s}$ 51. (a) 721 m/s ; (b) 937 m/s 53. (a) 33% ; (b) 23% ; (c) decreases 55. (a) $+2.0\text{ m/s}$; (b) -1.3 J ; (c) $+40\text{ J}$; (d) system got energy from some source, such as a small explosion 57. (a) 4.4 m/s ; (b) 0.80 59. 25 cm 61. (a) 99 g ; (b) 1.9 m/s ; (c) 0.93 m/s 63. (a) 3.00 m/s ; (b) 6.00 m/s 65. (a) 1.2 kg ; (b) 2.5 m/s 67. -28 cm 69. (a) 0.21 kg ; (b) 7.2 m 71. (a) $4.15 \times 10^5\text{ m/s}$; (b) $4.84 \times 10^5\text{ m/s}$ 73. 120° 75. (a) 433 m/s ; (b) 250 m/s 77. (a) 46 N ; (b) none 79. (a) $1.57 \times 10^6\text{ N}$; (b) $1.35 \times 10^5\text{ kg}$; (c) 2.08 km/s 81. (a) 7290 m/s ; (b) 8200 m/s ; (c) $1.271 \times 10^{10}\text{ J}$; (d) $1.275 \times 10^{10}\text{ J}$ 83. (a) 1.92 m ; (b) 0.640 m 85. (a) 1.78 m/s ; (b) less; (c) less; (d) greater 87. (a) 3.7 m/s ; (b) $1.3\text{ N} \cdot \text{s}$; (c) $1.8 \times 10^2\text{ N}$ 89. (a) $(7.4 \times 10^3\text{ N} \cdot \text{s})\hat{i} - (7.4 \times 10^3\text{ N} \cdot \text{s})\hat{j}$; (b) $(-7.4 \times 10^3\text{ N} \cdot \text{s})\hat{i}$; (c) $2.3 \times 10^3\text{ N}$; (d) $2.1 \times 10^4\text{ N}$; (e) -45° 91. $+4.4\text{ m/s}$ 93. $1.18 \times 10^4\text{ kg}$ 95. (a) 1.9 m/s ; (b) -30° ; (c) elastic 97. (a) 6.9 m/s ; (b) 30° ; (c) 6.9 m/s ; (d) -30° ; (e) 2.0 m/s ; (f) -180° 99. (a) 25 mm ; (b) 26 mm ; (c) down; (d) $1.6 \times 10^{-2}\text{ m/s}^2$ 101. 29 J 103. 2.2 kg 105. 5.0 kg 107. (a) 50 kg/s ; (b) $1.6 \times 10^2\text{ kg/s}$ 109. (a) $4.6 \times 10^3\text{ km}$; (b) 73% 111. 190 m/s 113. 28.8 N 115. (a) 0.745 mm ; (b) 153° ; (c) 1.67 mJ 117. (a) $(2.67\text{ m/s})\hat{i} + (-3.00\text{ m/s})\hat{j}$; (b) 4.01 m/s ; (c) 48.4° 119. (a) -0.50 m ; (b) -1.8 cm ; (c) 0.50 m 121. 0.22%

CHAPTER 10

CP 1. b and c 2. (a) and (d) ($\alpha = d^2\theta/dt^2$ must be a constant) 3. (a) yes; (b) no; (c) yes; (d) yes 4. all tie 5. 1, 2, 4, 3 (see Eq. 10-36) 6. (see Eq. 10-40) 1 and 3 tie, 4, then 2 and 5 tie (zero) 7. (a) downward in the figure ($\tau_{\text{net}} = 0$); (b) less (consider moment arms)
Q 1. (a) c, a , then b and d tie; (b) b , then a and c tie, then d 3. all tie 5. (a) decrease; (b) clockwise; (c) counterclockwise 7. larger 9. c, a, b
P 1. 14 rev 3. (a) 4.0 rad/s ; (b) 11.9 rad/s 5. 11 rad/s 7. (a) 4.0 m/s ; (b) no 9. (a) 3.00 s ; (b) 18.9 rad 11. (a) 30 s ; (b) $1.8 \times 10^3\text{ rad}$ 13. (a) $3.4 \times 10^2\text{ s}$; (b) $-4.5 \times 10^{-3}\text{ rad/s}^2$; (c) 98 s 15. 8.0 s 17. (a) 44 rad ; (b) 5.5 s ; (c) 32 s ; (d) -2.1 s ; (e) 40 s 19. (a) $2.50 \times 10^{-3}\text{ rad/s}$; (b) 20.2 m/s^2 ; (c) 0 21. $6.9 \times 10^{-13}\text{ rad/s}$ 23. (a) 20.9 rad/s ; (b) 12.5 m/s ; (c) 800 rev/min^2 ; (d) 600 rev 25. (a) $7.3 \times 10^{-5}\text{ rad/s}$; (b) $3.5 \times 10^2\text{ m/s}$; (c) $7.3 \times 10^{-5}\text{ rad/s}$; (d) $4.6 \times 10^2\text{ m/s}$ 27. (a) 73 cm/s^2 ; (b) 0.075 ; (c) 0.11 29. (a) $3.8 \times 10^3\text{ rad/s}$; (b) $1.9 \times 10^2\text{ m/s}$ 31. (a) 40 s ; (b) 2.0 rad/s^2 33. $12.3\text{ kg} \cdot \text{m}^2$ 35. (a) 1.1 kJ ; (b) 9.7 kJ 37. $0.097\text{ kg} \cdot \text{m}^2$ 39. (a) 49 MJ ; (b) $1.0 \times 10^2\text{ min}$ 41. (a) $0.023\text{ kg} \cdot \text{m}^2$; (b) 1.1 mJ 43. $4.7 \times 10^{-4}\text{ kg} \cdot \text{m}^2$ 45. $-3.85\text{ N} \cdot \text{m}$ 47. $4.6\text{ N} \cdot \text{m}$ 49. (a) 28.2 rad/s^2 ; (b) $338\text{ N} \cdot \text{m}$ 51. (a) 6.00 cm/s^2 ; (b) 4.81 N ; (c) 4.54 N ; (d) 1.20 rad/s^2 ; (e) $0.0138\text{ kg} \cdot \text{m}^2$ 53. 0.140 N 55. $2.57 \times 10^{-4}\text{ kg} \cdot \text{m}^2$ 57. (a) $4.2 \times 10^2\text{ rad/s}^2$; (b) $5.0 \times 10^2\text{ rad/s}$ 59. $396\text{ N} \cdot \text{m}$ 61. (a) -19.8 kJ ; (b) 1.32 kW 63. 5.42 m/s 65. (a) 5.32 m/s^2 ; (b) 8.43 m/s^2 ; (c) 41.8° 67. 9.82 rad/s 69. $6.16 \times 10^{-5}\text{ kg} \cdot \text{m}^2$ 71. (a) 31.4 rad/s^2 ; (b) 0.754 m/s^2 ; (c) 56.1 N ; (d) 55.1 N 73. (a) $4.81 \times 10^5\text{ N}$; (b) $1.12 \times 10^4\text{ N} \cdot \text{m}$; (c) $1.25 \times 10^6\text{ J}$ 75. (a) 2.3

rad/s²; (b) 1.4 rad/s² 77. (a) -67 rev/min²; (b) 8.3 rev 81. 3.1 rad/s
83. (a) 1.57 m/s²; (b) 4.55 N; (c) 4.94 N 85. 30 rev 87. 0.054 kg · m²
89. $1.4 \times 10^2 \text{ N} \cdot \text{m}$ 91. (a) 10 J; (b) 0.27 m 93. 4.6 rad/s² 95. 2.6 J
97. (a) $5.92 \times 10^4 \text{ m/s}^2$; (b) $4.39 \times 10^4 \text{ s}^{-2}$ 99. (a) 0.791 kg · m²; (b)
 $1.79 \times 10^{-2} \text{ N} \cdot \text{m}$ 101. (a) $1.5 \times 10^2 \text{ cm/s}$; (b) 15 rad/s; (c) 15 rad/s;
(d) 75 cm/s; (e) 3.0 rad/s 103. (a) 7.0 kg · m²; (b) 7.2 m/s; (c) 71°

CHAPTER 11

CP 1. (a) same; (b) less 2. less (consider the transfer of energy
from rotational kinetic energy to gravitational potential energy) 3.
(draw the vectors, use right-hand rule) (a) $\pm z$; (b) y ; (c) $-x$ 4.
(see Eq. 11-21) (a) 1 and 3 tie; then 2 and 4 tie, then 5 (zero); (b) 2
and 3 5. (see Eqs. 11-23 and 11-16) (a) 3, 1; then 2 and 4 tie (zero);
(b) 3 6. (a) all tie (same τ , same t , thus same ΔL); (b) sphere, disk,
hoop (reverse order of I) 7. (a) decreases; (b) same ($\tau_{\text{net}} = 0$, so L
is conserved); (c) increases
Q 1. a , then b and c tie, then e, d (zero) 3. (a) spins in place; (b)
rolls toward you; (c) rolls away from you 5. (a) 1, 2, 3 (zero); (b) 1
and 2 tie, then 3; (c) 1 and 3 tie, then 2 7. (a) same; (b) increase; (c)
decrease; (d) same, decrease, increase 9. D, B , then A and C tie
P 1. (a) 0; (b) $(22 \text{ m/s})\hat{j}$; (c) $(-22 \text{ m/s})\hat{j}$; (d) 0; (e) $1.5 \times 10^3 \text{ m/s}^2$; (f)
 $1.5 \times 10^3 \text{ m/s}^2$; (g) $(22 \text{ m/s})\hat{i}$; (h) $(44 \text{ m/s})\hat{i}$; (i) 0; (j) 0; (k) 1.5×10^3
 m/s^2 ; (l) $1.5 \times 10^3 \text{ m/s}^2$ 3. -3.15 J 5. 0.020 7. (a) 63 rad/s; (b) 4.0 m
9. 4.8 m 11. (a) $(-4.0 \text{ N})\hat{i}$; (b) $0.60 \text{ kg} \cdot \text{m}^2$ 13. 0.50 15. (a)
 $-(0.11 \text{ m})\omega$; (b) -2.1 m/s^2 ; (c) -47 rad/s^2 ; (d) 1.2 s; (e) 8.6 m; (f) 6.1
 m/s 17. (a) 13 cm/s²; (b) 4.4 s; (c) 55 cm/s; (d) 18 mJ; (e) 1.4 J; (f) 27
rev/s 19. $(-2.0 \text{ N} \cdot \text{m})\hat{j}$ 21. (a) $(6.0 \text{ N} \cdot \text{m})\hat{j} + (8.0 \text{ N} \cdot \text{m})\hat{k}$; (b)
 $(-22 \text{ N} \cdot \text{m})\hat{j}$ 23. (a) $(-1.5 \text{ N} \cdot \text{m})\hat{j} - (4.0 \text{ N} \cdot \text{m})\hat{j} - (1.0 \text{ N} \cdot \text{m})\hat{k}$;
(b) $(-1.5 \text{ N} \cdot \text{m})\hat{j} - (4.0 \text{ N} \cdot \text{m})\hat{j} - (1.0 \text{ N} \cdot \text{m})\hat{k}$ 25. (a) $(50 \text{ N} \cdot \text{m})\hat{k}$;
(b) 90° 27. (a) 0; (b) $(8.0 \text{ N} \cdot \text{m})\hat{i} + (8.0 \text{ N} \cdot \text{m})\hat{k}$ 29. (a) 9.8
 $\text{kg} \cdot \text{m}^2/\text{s}$; (b) $+z$ direction 31. (a) 0; (b) $-22.6 \text{ kg} \cdot \text{m}^2/\text{s}$; (c) -7.84
 $\text{N} \cdot \text{m}$; (d) $-7.84 \text{ N} \cdot \text{m}$ 33. (a) $(-1.7 \times 10^2 \text{ kg} \cdot \text{m}^2/\text{s})\hat{k}$; (b) $(+56$
 $\text{N} \cdot \text{m})\hat{k}$; (c) $(+56 \text{ kg} \cdot \text{m}^2/\text{s}^2)\hat{k}$ 35. (a) $48\hat{k} \text{ N} \cdot \text{m}$; (b) increasing
37. (a) $4.6 \times 10^{-3} \text{ kg} \cdot \text{m}^2$; (b) $1.1 \times 10^{-3} \text{ kg} \cdot \text{m}^2/\text{s}$; (c) 3.9×10^{-3}
 $\text{kg} \cdot \text{m}^2/\text{s}$ 39. (a) $1.47 \text{ N} \cdot \text{m}$; (b) 20.4 rad; (c) -29.9 J ; (d) 19.9 W
41. (a) $1.6 \text{ kg} \cdot \text{m}^2$; (b) $4.0 \text{ kg} \cdot \text{m}^2/\text{s}$ 43. (a) 1.5 m; (b) 0.93 rad/s; (c) 98
J; (d) 8.4 rad/s; (e) $8.8 \times 10^2 \text{ J}$; (f) internal energy of the skaters 45.
(a) 3.6 rev/s; (b) 3.0; (c) forces on the bricks from the man trans-
ferred energy from the man's internal energy to kinetic energy 47.
0.17 rad/s 49. (a) 750 rev/min; (b) 450 rev/min; (c) clockwise 51.
(a) 267 rev/min; (b) 0.667 53. $1.3 \times 10^3 \text{ m/s}$ 55. 3.4 rad/s 57. (a)
18 rad/s; (b) 0.92 59. 11.0 m/s 61. 1.5 rad/s 63. 0.070 rad/s 65. (a)
0.148 rad/s; (b) 0.0123; (c) 181° 67. (a) 0.180 m; (b) clockwise 69.
0.041 rad/s 71. (a) 1.6 m/s^2 ; (b) 16 rad/s^2 ; (c) $(4.0 \text{ N})\hat{i}$ 73. (a) 0; (b)
0; (c) $-30\hat{k} \text{ kg} \cdot \text{m}^2/\text{s}$; (d) $-90\hat{k} \text{ N} \cdot \text{m}$; (e) $30\hat{k} \text{ kg} \cdot \text{m}^2/\text{s}$; (f) $90\hat{k}$
 $\text{N} \cdot \text{m}$ 75. (a) $149 \text{ kg} \cdot \text{m}^2$; (b) $158 \text{ kg} \cdot \text{m}^2/\text{s}$; (c) 0.744 rad/s 77. (a)
 $6.65 \times 10^{-5} \text{ kg} \cdot \text{m}^2/\text{s}$; (b) no; (c) 0; (d) yes 79. (a) 0.333; (b) 0.111
81. (a) 58.8 J; (b) 39.2 J 83. (a) 61.7 J; (b) 3.43 m; (c) no 85. (a)
 $mvR/(I + MR^2)$; (b) $mvR^2/(I + MR^2)$ 87. rotational speed would
decrease; day would be about 0.8 s longer 89. (a) 12.7 rad/s; (b)
clockwise 91. (a) 0.81 mJ; (b) 0.29; (c) $1.3 \times 10^{-2} \text{ N}$ 93. (a) $mR^2/2$;
(b) a solid circular cylinder

CHAPTER 12

CP 1. c, e, f 2. (a) no; (b) at site of $\vec{F}_1$, perpendicular to plane of
figure; (c) 45 N 3. d
Q 1. (a) 1 and 3 tie, then 2; (b) all tie; (c) 1 and 3 tie, then 2 (zero)
3. a and c (forces and torques balance) 5. (a) 12 kg; (b) 3 kg;
(c) 1 kg 7. (a) at C (to eliminate forces there from a torque
equation); (b) plus; (c) minus; (d) equal 9. increase

P 1. (a) 1.00 m; (b) 2.00 m; (c) 0.987 m; (d) 1.97 m 3. (a) 9.4 N;
(b) 4.4 N 5. 7.92 kN 7. (a) $2.8 \times 10^2 \text{ N}$; (b) $8.8 \times 10^2 \text{ N}$; (c) 71°
9. 74.4 g 11. (a) 1.2 kN; (b) down; (c) 1.7 kN; (d) up; (e) left;
(f) right 13. (a) 2.7 kN; (b) up; (c) 3.6 kN; (d) down 15. (a) 5.0 N;
(b) 30 N; (c) 1.3 m 17. (a) 0.64 m; (b) increased 19. 8.7 N
21. (a) 6.63 kN; (b) 5.74 kN; (c) 5.96 kN 23. (a) 192 N; (b) 96.1 N;
(c) 55.5 N 25. 13.6 N 27. (a) 1.9 kN; (b) up; (c) 2.1 kN; (d) down
29. (a) $(-80 \text{ N})\hat{i} + (1.3 \times 10^2 \text{ N})\hat{j}$; (b) $(80 \text{ N})\hat{i} + (1.3 \times 10^2 \text{ N})\hat{j}$
31. 2.20 m 33. (a) 60.0°; (b) 300 N 35. (a) 445 N; (b) 0.50; (c) 315 N
37. 0.34 39. (a) 211 N; (b) 534 N; (c) 320 N 41. (a) slides;
(b) 31°; (c) tips; (d) 34° 43. (a) $6.5 \times 10^6 \text{ N/m}^2$; (b) $1.1 \times 10^{-5} \text{ m}$
45. (a) 0.80; (b) 0.20; (c) 0.25 47. (a) $1.4 \times 10^9 \text{ N}$; (b) 75
49. (a) 866 N; (b) 143 N; (c) 0.165 51. (a) $1.2 \times 10^2 \text{ N}$; (b) 68 N
53. (a) $1.8 \times 10^7 \text{ N}$; (b) $1.4 \times 10^7 \text{ N}$; (c) 16 55. 0.29 57. 76 N
59. (a) 8.01 kN; (b) 3.65 kN; (c) 5.66 kN 61. 71.7 N 63. (a) $L/2$;
(b) $L/4$; (c) $L/6$; (d) $L/8$; (e) $25L/24$ 65. (a) 88 N; (b) $(30\hat{i} + 97\hat{j}) \text{ N}$
67. $2.4 \times 10^9 \text{ N/m}^2$ 69. 60° 71. (a) $\mu < 0.57$; (b) $\mu > 0.57$
73. (a) $(35\hat{i} + 200\hat{j}) \text{ N}$; (b) $(-45\hat{i} + 200\hat{j}) \text{ N}$; (c) $1.9 \times 10^2 \text{ N}$
75. (a) BC, CD, DA ; (b) 535 N; (c) 757 N 77. (a) 1.38 kN; (b) 180 N
79. (a) $a_1 = L/2, a_2 = 5L/8, h = 9L/8$; (b) $b_1 = 2L/3, b_2 = L/2$,
 $h = 7L/6$ 81. $L/4$ 83. (a) 106 N; (b) 64.0°

CHAPTER 13

CP 1. all tie 2. (a) 1, tie of 2 and 4, then 3; (b) line d
3. (a) increase; (b) negative 4. (a) 2; (b) 1 5. (a) path 1
(decreased E (more negative) gives decreased a); (b) less
(decreased a gives decreased T)
Q 1. $3GM^2/d^2$, leftward 3. Gm^2/r^2 , upward 5. b and c tie, then a
(zero) 7. 1, tie of 2 and 4, then 3 9. (a) positive y ; (b) yes, rotates
counterclockwise until it points toward particle B 11. b, d , and f
all tie, then e, c, a
P 1. $\frac{1}{2}$ 3. 19 m 5. 0.8 m 7. -5.00d 9. $2.60 \times 10^5 \text{ km}$
11. (a) $M = m$; (b) 0 13. $8.31 \times 10^{-9} \text{ N}$ 15. (a) -1.88d;
(b) -3.90d; (c) 0.489d 17. (a) 17 N; (b) 2.4 19. $2.6 \times 10^6 \text{ m}$
21. $5 \times 10^{24} \text{ kg}$ 23. (a) 7.6 m/s^2 ; (b) 4.2 m/s^2 25. (a) $(3.0 \times$
 $10^{-7} \text{ N/kg})\hat{m}$; (b) $(3.3 \times 10^{-7} \text{ N/kg})\hat{m}$; (c) $(6.7 \times 10^{-7} \text{ N/kg} \cdot \text{m})\hat{m}r$
27. (a) 9.83 m/s^2 ; (b) 9.84 m/s^2 ; (c) 9.79 m/s^2 29. $5.0 \times 10^9 \text{ J}$
31. (a) 0.74; (b) 3.8 m/s²; (c) 5.0 km/s 33. (a) 0.0451; (b) 28.5
35. $-4.82 \times 10^{-13} \text{ J}$ 37. (a) 0.50 pJ; (b) -0.50 pJ 39. (a) 1.7 km/s;
(b) $2.5 \times 10^5 \text{ m}$; (c) 1.4 km/s 41. (a) 82 km/s; (b) $1.8 \times 10^4 \text{ km/s}$
43. (a) 7.82 km/s; (b) 87.5 min 45. $6.5 \times 10^{23} \text{ kg}$ 47. $5 \times 10^{10} \text{ stars}$
49. (a) $1.9 \times 10^{13} \text{ m}$; (b) $3.6R_p$ 51. (a) $6.64 \times 10^3 \text{ km}$; (b) 0.0136
53. $5.8 \times 10^6 \text{ m}$ 57. 0.71 y 59. $(GM/L)^{0.5}$ 61. (a) $3.19 \times 10^3 \text{ km}$;
(b) lifting 63. (a) 2.8 y; (b) 1.0×10^{-4} 65. (a) $r^{1.5}$; (b) r^{-1} ; (c) $r^{0.5}$;
(d) $r^{-0.5}$ 67. (a) 7.5 km/s; (b) 97 min; (c) $4.1 \times 10^2 \text{ km}$; (d) 7.7 km/s;
(e) 93 min; (f) $3.2 \times 10^{-3} \text{ N}$; (g) no; (h) yes 69. 1.1 s
71. (a) $GMmx(x^2 + R^2)^{-3/2}$; (b) $[2GM(R^{-1} - (R^2 + x^2)^{-1/2})]^{1/2}$
73. (a) $1.0 \times 10^3 \text{ kg}$; (b) 1.5 km/s 75. $3.2 \times 10^{-7} \text{ N}$ 77. $0.037\hat{j} \mu\text{N}$
79. $2\pi^{1.5}G^{-0.5}(M + m/4)^{-0.5}$ 81. (a) $2.2 \times 10^{-7} \text{ rad/s}$; (b) 89 km/s
83. (a) $2.15 \times 10^4 \text{ s}$; (b) 12.3 km/s; (c) 12.0 km/s; (d) $2.17 \times 10^{11} \text{ J}$;
(e) $-4.53 \times 10^{11} \text{ J}$; (f) $-2.35 \times 10^{11} \text{ J}$; (g) $4.04 \times 10^7 \text{ m}$; (h) $1.22 \times$
 10^3 s ; (i) elliptical 85. $2.5 \times 10^4 \text{ km}$ 87. (a) $1.4 \times 10^6 \text{ m/s}$; (b) $3 \times$
 10^6 m/s^2 89. (a) 0; (b) $1.8 \times 10^{32} \text{ J}$; (c) $1.8 \times 10^{32} \text{ J}$; (d) 0.99 km/s
91. (a) Gm^2/R_i ; (b) $Gm^2/2R_i$; (c) $(Gm/R_i)^{0.5}$; (d) $2(Gm/R_i)^{0.5}$;
(e) Gm^2/R_i ; (f) $(2Gm/R_i)^{0.5}$; (g) The center-of-mass frame is an in-
ertial frame, and in it the principle of conservation of energy may
be written as in Chapter 8; the reference frame attached to body A
is noninertial, and the principle cannot be written as in Chapter 8.
Answer d is correct. 93. $2.4 \times 10^4 \text{ m/s}$ 95. $-0.044\hat{j} \mu\text{N}$
97. $GM_E m/12R_E$

CHAPTER 14

CP 1. all tie 2. (a) all tie (the gravitational force on the penguin is the same); (b) $0.95\rho_0$, ρ_0 , $1.1\rho_0$ 3. $13\text{ cm}^3/\text{s}$, outward
 4. (a) all tie; (b) 1, then 2 and 3 tie, 4 (wider means slower); (c) 4, 3, 2, 1 (wider and lower mean more pressure)
Q 1. (a) moves downward; (b) moves downward 3. (a) downward; (b) downward; (c) same 5. b , then a and d tie (zero), then c
 7. (a) 1 and 4; (b) 2; (c) 3 9. B , C , A
P 1. 0.074 3. $1.1 \times 10^5\text{ Pa}$ 5. $2.9 \times 10^4\text{ N}$ 7. (b) 26 kN
 9. (a) $1.0 \times 10^3\text{ torr}$; (b) $1.7 \times 10^3\text{ torr}$ 11. (a) 94 torr ; (b) $4.1 \times 10^2\text{ torr}$; (c) $3.1 \times 10^2\text{ torr}$ 13. $1.08 \times 10^3\text{ atm}$ 15. $-2.6 \times 10^4\text{ Pa}$
 17. $7.2 \times 10^5\text{ N}$ 19. $4.69 \times 10^5\text{ N}$ 21. 0.635 J 23. 44 km
 25. 739.26 torr 27. (a) 7.9 km ; (b) 16 km 29. 8.50 kg 31. (a) $6.7 \times 10^2\text{ kg/m}^3$; (b) $7.4 \times 10^2\text{ kg/m}^3$ 33. (a) $2.04 \times 10^{-2}\text{ m}^3$; (b) 1.57 kN 35. five 37. 57.3 cm 39. (a) 1.2 kg ; (b) $1.3 \times 10^3\text{ kg/m}^3$ 41. (a) 0.10 ; (b) 0.083 43. (a) 637.8 cm^3 ; (b) 5.102 m^3 ; (c) $5.102 \times 10^3\text{ kg}$ 45. 0.126 m^3 47. (a) 1.80 m^3 ; (b) 4.75 m^3
 49. (a) 3.0 m/s ; (b) 2.8 m/s 51. 8.1 m/s 53. 66 W 55. $1.4 \times 10^5\text{ J}$
 57. (a) $1.6 \times 10^{-3}\text{ m}^3/\text{s}$; (b) 0.90 m 59. (a) 2.5 m/s ; (b) $2.6 \times 10^5\text{ Pa}$
 61. (a) 3.9 m/s ; (b) 88 kPa 63. $1.1 \times 10^2\text{ m/s}$ 65. (b) $2.0 \times 10^{-2}\text{ m}^3/\text{s}$ 67. (a) 74 N ; (b) $1.5 \times 10^2\text{ m}^3$ 69. (a) $0.0776\text{ m}^3/\text{s}$; (b) 69.8 kg/s 71. (a) 35 cm ; (b) 30 cm ; (c) 20 cm 73. 1.5 g/cm^3 75. $5.11 \times 10^{-7}\text{ kg}$ 77. 44.2 g 79. $6.0 \times 10^2\text{ kg/m}^3$ 81. 45.3 cm^3
 83. (a) 3.2 m/s ; (b) $9.2 \times 10^4\text{ Pa}$; (c) 10.3 m 85. $1.07 \times 10^3\text{ g}$

CHAPTER 15

CP 1. (sketch x versus t) (a) $-x_m$; (b) $+x_m$; (c) 0 2. a (F must have the form of Eq. 15-10) 3. (a) 5 J ; (b) 2 J ; (c) 5 J 4. all tie (in Eq. 15-29, m is included in I) 5. 1, 2, 3 (the ratio m/b matters; k does not)
Q 1. a and b 3. (a) 2; (b) positive; (c) between 0 and $+x_m$
 5. (a) between D and E ; (b) between $3\pi/2$ rad and 2π rad
 7. (a) all tie; (b) 3, then 1 and 2 tie; (c) 1, 2, 3 (zero); (d) 1, 2, 3 (zero); (e) 1, 3, 2 9. b (infinite period, does not oscillate), c , a
 11. (a) greater; (b) same; (c) same; (d) greater; (e) greater
P 1. (a) 0.50 s ; (b) 2.0 Hz ; (c) 18 cm 3. 37.8 m/s^2 5. (a) 1.0 mm ; (b) 0.75 m/s ; (c) $5.7 \times 10^2\text{ m/s}^2$ 7. (a) 498 Hz ; (b) greater
 9. (a) 3.0 m ; (b) -49 m/s ; (c) $-2.7 \times 10^2\text{ m/s}^2$; (d) 20 rad ; (e) 1.5 Hz ; (f) 0.67 s 11. 39.6 Hz 13. (a) 0.500 s ; (b) 2.00 Hz ; (c) 12.6 rad/s ; (d) 79.0 N/m ; (e) 4.40 m/s ; (f) 27.6 N 15. (a) 0.184 ; (b) same direction 17. (a) 5.58 Hz ; (b) 0.325 kg ; (c) 0.400 m 19. (a) 25 cm ; (b) 2.2 Hz 21. 54 Hz 23. 3.1 cm 25. (a) 0.525 m ; (b) 0.686 s
 27. (a) 0.75 ; (b) 0.25 ; (c) $2^{-0.5}x_m$ 29. 37 mJ 31. (a) 2.25 Hz ; (b) 125 J ; (c) 250 J ; (d) 86.6 cm 33. (a) 1.1 m/s ; (b) 3.3 cm
 35. (a) 3.1 ms ; (b) 4.0 m/s ; (c) 0.080 J ; (d) 80 N ; (e) 40 N
 37. (a) 2.2 Hz ; (b) 56 cm/s ; (c) 0.10 kg ; (d) 20.0 cm 39. (a) 39.5 rad/s ; (b) 34.2 rad/s ; (c) 124 rad/s^2 41. (a) $0.205\text{ kg}\cdot\text{m}^2$; (b) 47.7 cm ; (c) 1.50 s 43. (a) 1.64 s ; (b) equal 45. 8.77 s 47. 0.366 s
 49. (a) 0.845 rad ; (b) 0.0602 rad 51. (a) 0.53 m ; (b) 2.1 s
 53. 0.0653 s 55. (a) 2.26 s ; (b) increases; (c) same 57. 6.0%
 59. (a) 14.3 s ; (b) 5.27 61. (a) $F_m/b\omega$; (b) F_m/b 63. 5.0 cm
 65. (a) $2.8 \times 10^3\text{ rad/s}$; (b) 2.1 m/s ; (c) 5.7 km/s^2 67. (a) 1.1 Hz ; (b) 5.0 cm 69. 7.2 m/s 71. (a) 7.90 N/m ; (b) 1.19 cm ; (c) 2.00 Hz
 73. (a) $1.3 \times 10^2\text{ N/m}$; (b) 0.62 s ; (c) 1.6 Hz ; (d) 5.0 cm ; (e) 0.51 m/s
 75. (a) 16.6 cm ; (b) 1.23% 77. (a) 1.2 J ; (b) 50 79. 1.53 m
 81. (a) 0.30 m ; (b) 0.28 s ; (c) $1.5 \times 10^2\text{ m/s}^2$; (d) 11 J 83. (a) 1.23 kN/m ; (b) 76.0 N 85. 1.6 kg 87. (a) $0.735\text{ kg}\cdot\text{m}^2$; (b) $0.0240\text{ N}\cdot\text{m}$; (c) 0.181 rad/s 89. (a) 3.5 m ; (b) 0.75 s 91. (a) 0.35 Hz ; (b) 0.39 Hz ; (c) 0 (no oscillation) 93. (a) 245 N/m ; (b) 0.284 s

95. $0.079\text{ kg}\cdot\text{m}^2$ 97. (a) $8.11 \times 10^{-5}\text{ kg}\cdot\text{m}^2$; (b) 3.14 rad/s
 99. 14.0° 101. (a) 3.2 Hz ; (b) 0.26 m ; (c) $x = (0.26\text{ m})\cos(20t - \pi/2)$, with t in seconds 103. (a) 0.44 s ; (b) 0.18 m
 105. (a) 0.45 s ; (b) 0.10 m above and 0.20 m below; (c) 0.15 m ; (d) 2.3 J 107. $7 \times 10^2\text{ N/m}$

CHAPTER 16

CP 1. a , 2 , b , 3 ; c , 1 (compare with the phase in Eq. 16-2, then see Eq. 16-5) 2. (a) 2 , 3 , 1 (see Eq. 16-12); (b) 3 , then 1 and 2 tie (find amplitude of dy/dt) 3. (a) same (independent of f); (b) decrease ($\lambda = v/f$); (c) increase; (d) increase 4. 0.20 and 0.80 tie, then 0.60 , 0.45 5. (a) 1 ; (b) 3 ; (c) 2 6. (a) 75 Hz ; (b) 525 Hz
Q 1. (a) 1 , 4 , 2 , 3 ; (b) 1 , 4 , 2 , 3 3. a , upward; b , upward; c , downward; d , downward; e , downward; f , downward; g , upward; h , upward
 5. intermediate (closer to fully destructive) 7. (a) 0 , 0.2 wavelength, 0.5 wavelength (zero); (b) $4P_{\text{avg},1}$ 9. d 11. c , a , b
P 1. 1.1 ms 3. (a) 3.49 m^{-1} ; (b) 31.5 m/s 5. (a) 0.680 s ; (b) 1.47 Hz ; (c) 2.06 m/s 7. (a) 64 Hz ; (b) 1.3 m ; (c) 4.0 cm ; (d) 5.0 m^{-1} ; (e) $4.0 \times 10^2\text{ s}^{-1}$; (f) $\pi/2\text{ rad}$; (g) minus 9. (a) 3.0 mm ; (b) 16 m^{-1} ; (c) $2.4 \times 10^2\text{ s}^{-1}$; (d) minus 11. (a) negative; (b) 4.0 cm ; (c) 0.31 cm^{-1} ; (d) 0.63 s^{-1} ; (e) $\pi\text{ rad}$; (f) minus; (g) 2.0 cm/s ; (h) -2.5 cm/s
 13. (a) 11.7 cm ; (b) $\pi\text{ rad}$ 15. (a) 0.12 mm ; (b) 141 m^{-1} ; (c) 628 s^{-1} ; (d) plus 17. (a) 15 m/s ; (b) 0.036 N 19. 129 m/s 21. 2.63 m
 23. (a) 5.0 cm ; (b) 40 cm ; (c) 12 m/s ; (d) 0.033 s ; (e) 9.4 m/s ; (f) 16 m^{-1} ; (g) $1.9 \times 10^2\text{ s}^{-1}$; (h) 0.93 rad ; (i) plus 27. 3.2 mm
 29. 0.20 m/s 31. $1.41y_m$ 33. (a) 9.0 mm ; (b) 16 m^{-1} ; (c) $1.1 \times 10^3\text{ s}^{-1}$; (d) 2.7 rad ; (e) plus 35. 5.0 cm 37. (a) 3.29 mm ; (b) 1.55 rad ; (c) 1.55 rad 39. 84° 41. (a) 82.0 m/s ; (b) 16.8 m ; (c) 4.88 Hz
 43. (a) 7.91 Hz ; (b) 15.8 Hz ; (c) 23.7 Hz 45. (a) 105 Hz ; (b) 158 m/s
 47. 260 Hz 49. (a) 144 m/s ; (b) 60.0 cm ; (c) 241 Hz 51. (a) 0.50 cm ; (b) 3.1 m^{-1} ; (c) $3.1 \times 10^2\text{ s}^{-1}$; (d) minus 53. (a) 0.25 cm ; (b) $1.2 \times 10^2\text{ cm/s}$; (c) 3.0 cm ; (d) 0 55. 0.25 m 57. (a) 2.00 Hz ; (b) 2.00 m ; (c) 4.00 m/s ; (d) 50.0 cm ; (e) 150 cm ; (f) 250 cm ; (g) 0; (h) 100 cm ; (i) 200 cm
 59. (a) 324 Hz ; (b) eight 61. 36 N 63. (a) 75 Hz ; (b) 13 ms
 65. (a) 2.0 mm ; (b) 95 Hz ; (c) $+30\text{ m/s}$; (d) 31 cm ; (e) 1.2 m/s
 67. (a) 0.31 m ; (b) 1.64 rad ; (c) 2.2 mm 69. (a) $0.83y_1$; (b) 37°
 71. (a) 3.77 m/s ; (b) 12.3 N ; (c) 0; (d) 46.4 W ; (e) 0; (f) 0; (g) $\pm 0.50\text{ cm}$
 73. 1.2 rad 75. (a) 300 m/s ; (b) no 77. (a) $[k\Delta\ell(\ell + \Delta\ell)/m]^{0.5}$
 79. (a) 144 m/s ; (b) 3.00 m ; (c) 1.50 m ; (d) 48.0 Hz ; (e) 96.0 Hz
 81. (a) 1.00 cm ; (b) $3.46 \times 10^3\text{ s}^{-1}$; (c) 10.5 m^{-1} ; (d) plus 83. (a) $2\pi y_m/\lambda$; (b) no 85. (a) 240 cm ; (b) 120 cm ; (c) 80 cm 87. (a) 1.33 m/s ; (b) 1.88 m/s ; (c) 16.7 m/s^2 ; (d) 23.7 m/s^2 89. (a) 0.52 m ; (b) 40 m/s ; (c) 0.40 m 91. (a) 0.16 m ; (b) $2.4 \times 10^2\text{ N}$; (c) $y(x, t) = (0.16\text{ m})\sin[(1.57\text{ m}^{-1})x]\sin[(31.4\text{ s}^{-1})t]$ 93. (c) 2.0 m/s ; (d) $-x$

CHAPTER 17

CP 1. beginning to decrease (example: mentally move the curves of Fig. 17-6 rightward past the point at $x = 42\text{ cm}$) 2. (a) 1 and 2 tie, then 3 (see Eq. 17-28); (b) 3 , then 1 and 2 tie (see Eq. 17-26) 3. second (see Eqs. 17-39 and 17-41) 4. a , greater; b , less; c , can't tell; d , can't tell; e , greater; f , less
Q 1. (a) 0 , 0.2 wavelength, 0.5 wavelength (zero); (b) $4P_{\text{avg},1}$
 3. C , then A and B tie 5. E , A , D , C , B 7. 1 , 4 , 3 , 2 9. 150 Hz and 450 Hz
P 1. (a) 79 m ; (b) 41 m ; (c) 89 m 3. (a) 2.6 km ; (b) 2.0×10^2
 5. $1.9 \times 10^3\text{ km}$ 7. 40.7 m 9. 0.23 ms 11. (a) $76.2\text{ }\mu\text{m}$; (b) 0.333 mm
 13. 960 Hz 15. (a) $2.3 \times 10^2\text{ Hz}$; (b) higher 17. (a) 143 Hz ; (b) 3 ; (c) 5 ; (d) 286 Hz ; (e) 2 ; (f) 3 19. (a) 14 ; (b) 14 21. (a) 343 Hz ; (b) 3 ; (c) 5 ; (d) 686 Hz ; (e) 2 ; (f) 3 23. (a) 0 ; (b) fully constructive; (c) increase; (d) 128 m ; (e) 63.0 m ; (f) 41.2 m 25. 36.8 nm

27. (a) 1.0×10^3 ; (b) 32 29. 15.0 mW 31. $2 \mu\text{W}$ 33. $0.76 \mu\text{m}$
 35. (a) $5.97 \times 10^{-5} \text{ W/m}^2$; (b) 4.48 nW 37. (a) 0.34 nW; (b) 0.68 nW;
 (c) 1.4 nW; (d) 0.88 nW; (e) 0 39. (a) 405 m/s; (b) 596 N; (c) 44.0
 cm; (d) 37.3 cm 41. (a) 833 Hz; (b) 0.418 m 43. (a) 3; (b) 1129 Hz;
 (c) 1506 Hz 45. (a) 2; (b) 1 47. 12.4 m 49. 45.3 N 51. 2.25 ms
 53. 0.020 55. (a) 526 Hz; (b) 555 Hz 57. 0 59. (a) 1.022 kHz;
 (b) 1.045 kHz 61. 41 kHz 63. 155 Hz 65. (a) 2.0 kHz; (b) 2.0
 kHz 67. (a) 485.8 Hz; (b) 500.0 Hz; (c) 486.2 Hz; (d) 500.0 Hz
 69. (a) 42° ; (b) 11 s 71. 1 cm 73. 2.1 m 75. (a) $39.7 \mu\text{W/m}^2$;
 (b) 171 nm; (c) 0.893 Pa 77. 0.25 79. (a) 2.10 m; (b) 1.47 m
 81. (a) 59.7; (b) 2.81×10^{-4} 83. (a) rightward; (b) 0.90 m/s; (c) less
 85. (a) 11 ms; (b) 3.8 m 87. (a) $9.7 \times 10^2 \text{ Hz}$; (b) 1.0 kHz; (c) 60 Hz,
 no 89. (a) 21 nm; (b) 35 cm; (c) 24 nm; (d) 35 cm 91. (a) 7.70 Hz;
 (b) 7.70 Hz 93. (a) 5.2 kHz; (b) 2 95. (a) 10 W; (b) 0.032 W/m^2 ;
 (c) 99 dB 97. (a) 0; (b) 0.572 m; (c) 1.14 m 99. 171 m 101. (a)
 $3.6 \times 10^2 \text{ m/s}$; (b) 150 Hz 103. 400 Hz 105. (a) 14; (b) 12

CHAPTER 18

CP 1. (a) all tie; (b) 50°X , 50°Y , 50°W 2. (a) 2 and 3 tie, then 1,
 then 4; (b) 3, 2, then 1 and 4 tie (from Eqs. 18-9 and 18-10, assume
 that change in area is proportional to initial area) 3. A (see
 Eq. 18-14) 4. c and e (maximize area enclosed by a clockwise
 cycle) 5. (a) all tie (ΔE_{int} depends on i and f , not on path); (b) 4, 3,
 2, 1 (compare areas under curves); (c) 4, 3, 2, 1 (see Eq. 18-26)
 6. (a) zero (closed cycle); (b) negative (W_{net} is negative; see
 Eq. 18-26) 7. b and d tie, then a , c (P_{cond} identical; see Eq. 18-32)
 Q 1. c , then the rest tie 3. B , then A and C tie 5. (a) f , because
 ice temperature will not rise to freezing point and then drop; (b) b
 and c at freezing point, d above, e below; (c) in b liquid partly
 freezes and no ice melts; in c no liquid freezes and no ice melts; in d
 no liquid freezes and ice fully melts; in e liquid fully freezes and no
 ice melts 7. (a) both clockwise; (b) both clockwise 9. (a) greater;
 (b) 1, 2, 3; (c) 1, 3, 2; (d) 1, 2, 3; (e) 2, 3, 1 11. c , b , a
 P 1. 1.366 3. 348 K 5. (a) 320°F ; (b) -12.3°F 7. -92.1°X
 9. 2.731 cm 11. 49.87 cm^3 13. 29 cm^3 15. 360°C 17. 0.26 cm^3
 19. 0.13 mm 21. 7.5 cm 23. 160 s 25. 94.6 L 27. 42.7 kJ
 29. 33 m^2 31. 33 g 33. 3.0 min 35. 13.5°C 37. (a) 5.3°C ; (b) 0;
 (c) 0°C ; (d) 60 g 39. 742 kJ 41. (a) 0°C ; (b) 2.5°C 43. (a) $1.2 \times$
 10^2 J ; (b) 75 J; (c) 30 J 45. -30 J 47. (a) 6.0 cal; (b) -43 cal ;
 (c) 40 cal; (d) 18 cal; (e) 18 cal 49. 60 J 51. (a) 1.23 kW; (b)
 2.28 kW; (c) 1.05 kW 53. 1.66 kJ/s 55. (a) 16 J/s; (b) 0.048 g/s
 57. (a) $1.7 \times 10^4 \text{ W/m}^2$; (b) 18 W/m^2 59. 0.50 min 61. 0.40 cm/h
 63. -4.2°C 65. 1.1 m 67. 10% 69. (a) 80 J; (b) 80 J 71. $4.5 \times$
 $10^2 \text{ J/kg}\cdot\text{K}$ 73. 0.432 cm^3 75. $3.1 \times 10^2 \text{ J}$ 77. 79.5°C 79. 23 J
 81. (a) $11p_1V_1$; (b) $6p_1V_1$ 83. $4.83 \times 10^{-2} \text{ cm}^3$ 85. 10.5°C
 87. (a) 90 W; (b) $2.3 \times 10^2 \text{ W}$; (c) $3.3 \times 10^2 \text{ W}$ 89. (a) 1.87×10^4 ;
 (b) 10.4 h 91. 333 J 93. 8.6 J 95. (a) -45 J ; (b) $+45 \text{ J}$

CHAPTER 19

CP 1. all but c 2. (a) all tie; (b) 3, 2, 1 3. gas A 4. 5 (greatest
 change in T), then tie of 1, 2, 3, and 4 5. 1, 2, 3 ($Q_3 = 0$, Q_2 goes
 into work W_2 , but Q_1 goes into greater work W_1 and increases
 gas temperature)
 Q 1. d , then a and b tie, then c 3. 20 J 5. (a) 3; (b) 1; (c) 4; (d) 2;
 (e) yes 7. (a) 1, 2, 3, 4; (b) 1, 2, 3 9. constant-volume process
 P 1. 0.933 kg 3. (a) 0.0388 mol; (b) 220°C 5. 25 molecules/ cm^3
 7. (a) $3.14 \times 10^3 \text{ J}$; (b) from 9. 186 kPa 11. 5.60 kJ
 13. (a) 1.5 mol; (b) $1.8 \times 10^3 \text{ K}$; (c) $6.0 \times 10^2 \text{ K}$; (d) 5.0 kJ

15. 360 K 17. $2.0 \times 10^5 \text{ Pa}$ 19. (a) 511 m/s; (b) -200°C ; (c) 899°C
 21. $1.8 \times 10^2 \text{ m/s}$ 23. 1.9 kPa 25. (a) $5.65 \times 10^{-21} \text{ J}$; (b) $7.72 \times$
 10^{-21} J ; (c) 3.40 kJ; (d) 4.65 kJ 27. (a) $6.76 \times 10^{-20} \text{ J}$; (b) 10.7
 29. (a) $6 \times 10^9 \text{ km}$ 31. (a) $3.27 \times 10^{10} \text{ molecules/cm}^3$; (b) 172 m
 33. (a) 6.5 km/s; (b) 7.1 km/s 35. (a) 420 m/s; (b) 458 m/s; (c) yes
 37. (a) 0.67; (b) 1.2; (c) 1.3; (d) 0.33 39. (a) $1.0 \times 10^4 \text{ K}$; (b) $1.6 \times$
 10^5 K ; (c) $4.4 \times 10^2 \text{ K}$; (d) $7.0 \times 10^3 \text{ K}$; (e) no; (f) yes 41. (a) 7.0
 km/s; (b) $2.0 \times 10^{-8} \text{ cm}$; (c) $3.5 \times 10^{10} \text{ collisions/s}$ 43. (a) 3.49 kJ;
 (b) 2.49 kJ; (c) 997 J; (d) 1.00 kJ 45. (a) $6.6 \times 10^{-26} \text{ kg}$; (b) 40
 g/mol 47. (a) 0; (b) $+374 \text{ J}$; (c) $+374 \text{ J}$; (d) $+3.11 \times 10^{-22} \text{ J}$
 49. 15.8 J/mol·K 51. 8.0 kJ 53. (a) 6.98 kJ; (b) 4.99 kJ; (c) 1.99 kJ;
 (d) 2.99 kJ 55. (a) 14 atm; (b) $6.2 \times 10^2 \text{ K}$ 57. (a) diatomic;
 (b) 446 K; (c) 8.10 mol 59. -15 J 61. -20 J 63. (a) 3.74 kJ;
 (b) 3.74 kJ; (c) 0; (d) 0; (e) -1.81 kJ ; (f) 1.81 kJ; (g) -3.22 kJ ;
 (h) -1.93 kJ ; (i) -1.29 kJ ; (j) 520 J; (k) 0; (l) 520 J; (m) 0.0246 m^3 ;
 (n) 2.00 atm; (o) 0.0373 m^3 ; (p) 1.00 atm 65. (a) monatomic;
 (b) $2.7 \times 10^4 \text{ K}$; (c) $4.5 \times 10^4 \text{ mol}$; (d) 3.4 kJ; (e) $3.4 \times 10^2 \text{ kJ}$;
 (f) 0.010 67. (a) 2.00 atm; (b) 333 J; (c) 0.961 atm; (d) 236 J
 69. 349 K 71. (a) -374 J ; (b) 0; (c) $+374 \text{ J}$; (d) $+3.11 \times 10^{-22} \text{ J}$
 73. $7.03 \times 10^9 \text{ s}^{-1}$ 75. (a) 900 cal; (b) 0; (c) 900 cal; (d) 450 cal;
 (e) 1200 cal; (f) 300 cal; (g) 900 cal; (h) 450 cal; (i) 0; (j) -900 cal ;
 (k) 900 cal; (l) 450 cal 77. (a) $3/\nu_0^3$; (b) $0.750\nu_0$; (c) $0.775\nu_0$
 79. (a) -2.37 kJ ; (b) 2.37 kJ 81. (b) 125 J; (c) to 83. (a) 8.0 atm;
 (b) 300 K; (c) 4.4 kJ; (d) 3.2 atm; (e) 120 K; (f) 2.9 kJ; (g) 4.6 atm;
 (h) 170 K; (i) 3.4 kJ 85. (a) 38 L; (b) 71 g 87. -3.0 J

CHAPTER 20

CP 1. a, b, c 2. smaller (Q is smaller) 3. c, b, a 4. a, d, c, b 5. b
 Q 1. b , a , c , d 3. unchanged 5. a and c tie, then b and d tie
 7. (a) same; (b) increase; (c) decrease 9. A, first; B, first and
 second; C, second; D, neither
 P 1. (a) 9.22 kJ; (b) 23.1 J/K; (c) 0 3. 14.4 J/K 5. (a) $5.79 \times$
 10^4 J ; (b) 173 J/K 7. (a) 320 K; (b) 0; (c) $+1.72 \text{ J/K}$ 9. $+0.76 \text{ J/K}$
 11. (a) 57.0°C ; (b) -22.1 J/K ; (c) $+24.9 \text{ J/K}$; (d) $+2.8 \text{ J/K}$
 13. (a) -710 mJ/K ; (b) $+710 \text{ mJ/K}$; (c) $+723 \text{ mJ/K}$; (d) -723 mJ/K ;
 (e) $+13 \text{ mJ/K}$; (f) 0 15. (a) -943 J/K ; (b) $+943 \text{ J/K}$; (c) yes
 17. (a) 0.333; (b) 0.215; (c) 0.644; (d) 1.10; (e) 1.10; (f) 0; (g) 1.10;
 (h) 0; (i) -0.889 ; (j) -0.889 ; (k) -1.10 ; (l) -0.889 ; (m) 0; (n) 0.889;
 (o) 0 19. (a) 0.693; (b) 4.50; (c) 0.693; (d) 0; (e) 4.50; (f) 23.0 J/K;
 (g) -0.693 ; (h) 7.50; (i) -0.693 ; (j) 3.00; (k) 4.50; (l) 23.0 J/K
 21. -1.18 J/K 23. 97 K 25. (a) 266 K; (b) 341 K 27. (a) 23.6%;
 (b) $1.49 \times 10^4 \text{ J}$ 29. (a) 2.27 kJ; (b) 14.8 kJ; (c) 15.4%; (d) 75.0%;
 (e) greater 31. (a) 33 kJ; (b) 25 kJ; (c) 26 kJ; (d) 18 kJ
 33. (a) 1.47 kJ; (b) 554 J; (c) 918 J; (d) 62.4% 35. (a) 3.00; (b) 1.98;
 (c) 0.660; (d) 0.495; (e) 0.165; (f) 34.0% 37. 440 W 39. 20 J
 41. 0.25 hp 43. 2.03 47. (a) $W = N!/(n_1! n_2! n_3!)$; (b)
 $[(N/2)! (N/2)!]/[(N/3)! (N/3)! (N/3)!]$; (c) 4.2×10^{16} 49. 0.141 J/K·s
 51. (a) 87 m/s; (b) $1.2 \times 10^2 \text{ m/s}$; (c) 22 J/K 53. (a) 78%; (b) 82 kg/s
 55. (a) 40.9°C ; (b) -27.1 J/K ; (c) 30.3 J/K; (d) 3.18 J/K 57. $+3.59 \text{ J/K}$
 59. $1.18 \times 10^3 \text{ J/K}$ 63. (a) 0; (b) 0; (c) -23.0 J/K ; (d) 23.0 J/K
 65. (a) 25.5 kJ; (b) 4.73 kJ; (c) 18.5% 67. (a) 1.95 J/K; (b) 0.650 J/K;
 (c) 0.217 J/K; (d) 0.072 J/K; (e) decrease 69. (a) 4.45 J/K; (b) no
 71. (a) 1.26×10^{14} ; (b) 4.71×10^{13} ; (c) 0.37; (d) 1.01×10^{29} ;
 (e) 1.37×10^{28} ; (f) 0.14; (g) 9.05×10^{58} ; (h) 1.64×10^{57} ; (i) 0.018;
 (j) decrease 73. (a) 42.6 kJ; (b) 7.61 kJ 75. (a) 1; (b) 1; (c) 3;
 (d) 10; (e) $1.5 \times 10^{-23} \text{ J/K}$; (f) $3.2 \times 10^{-23} \text{ J/K}$

CHAPTER 21

CP 1. C and D attract; B and D attract 2. (a) leftward; (b) leftward; (c) leftward 3. (a) a, c, b ; (b) less than 4. $-15e$ (net charge of $-30e$ is equally shared)
Q 1. 3, 1, 2, 4 (zero) 3. a and b 5. $2kq^2/r^2$, up the page
 7. b and c tie, then a (zero) 9. (a) same; (b) less than; (c) cancel; (d) add; (e) adding components; (f) positive direction of y ; (g) negative direction of y ; (h) positive direction of x ; (i) negative direction of x
P 1. 0.500 3. 1.39 m 5. 2.81 N 7. -4.00 9. (a) $-1.00 \mu\text{C}$; (b) $3.00 \mu\text{C}$ 11. (a) 0.17 N; (b) -0.046 N 13. (a) -14 cm ; (b) 0 15. (a) 35 N; (b) -10° ; (c) -8.4 cm ; (d) $+2.7 \text{ cm}$ 17. (a) 1.60 N; (b) 2.77 N 19. (a) 3.00 cm; (b) 0; (c) -0.444 21. $3.8 \times 10^{-8} \text{ C}$ 23. (a) 0; (b) 12 cm; (c) 0; (d) $4.9 \times 10^{-26} \text{ N}$ 25. 6.3×10^{11} 27. (a) $3.2 \times 10^{-19} \text{ C}$; (b) 2 29. (a) -6.05 cm ; (b) 6.05 cm 31. 122 mA 33. $1.3 \times 10^7 \text{ C}$ 35. (a) 0; (b) $1.9 \times 10^{-9} \text{ N}$ 37. (a) 9B ; (b) 13N ; (c) 12C 39. $1.31 \times 10^{-22} \text{ N}$ 41. (a) $5.7 \times 10^{13} \text{ C}$; (b) cancels out; (c) $6.0 \times 10^5 \text{ kg}$ 43. (b) 3.1 cm 45. 0.19 MC 47. $-45 \mu\text{C}$ 49. 3.8 N 51. (a) 2.00×10^{10} electrons; (b) 1.33×10^{10} electrons 53. (a) 8.99 $\times 10^9 \text{ N}$; (b) 8.99 kN 55. (a) 0.5; (b) 0.15; (c) 0.85 57. $1.7 \times 10^8 \text{ N}$ 59. $-1.32 \times 10^{13} \text{ C}$ 61. (a) $(0.829 \text{ N})\hat{j}$; (b) $(-0.621 \text{ N})\hat{j}$ 63. $2.2 \times 10^{-6} \text{ kg}$ 65. $4.68 \times 10^{-19} \text{ N}$ 67. (a) $2.72L$; (b) 0 69. (a) $5.1 \times 10^2 \text{ N}$; (b) $7.7 \times 10^{28} \text{ m/s}^2$

CHAPTER 22

CP 1. (a) rightward; (b) leftward; (c) leftward; (d) rightward (p and e have same charge magnitude, and p is farther)
 2. (a) toward positive y ; (b) toward positive x ; (c) toward negative y
 3. (a) leftward; (b) leftward; (c) decrease 4. (a) all tie; (b) 1 and 3 tie, then 2 and 4 tie
Q 1. a, b, c 3. (a) yes; (b) toward; (c) no (the field vectors are not along the same line); (d) cancel; (e) add; (f) adding components; (g) toward negative y 5. (a) to their left; (b) no 7. (a) 4, 3, 1, 2; (b) 3, then 1 and 4 tie, then 2 9. a, b, c 11. e, b , then a and c tie, then d (zero)
P 3. (a) $3.07 \times 10^{21} \text{ N/C}$; (b) outward 5. 56 pC 7. $(1.02 \times 10^5 \text{ N/C})\hat{j}$ 9. (a) $1.38 \times 10^{-10} \text{ N/C}$; (b) 180° 11. -30 cm 13. (a) $3.60 \times 10^{-6} \text{ N/C}$; (b) $2.55 \times 10^{-6} \text{ N/C}$; (c) $3.60 \times 10^{-4} \text{ N/C}$; (d) $7.09 \times 10^{-7} \text{ N/C}$; (e) As the proton nears the disk, the forces on it from electrons e , more nearly cancel. 15. (a) 160 N/C; (b) 45° 17. (a) -90° ; (b) $+2.0 \mu\text{C}$; (c) $-1.6 \mu\text{C}$ 19. (a) $qd/4\pi\epsilon_0 r^3$; (b) -90° 23. 0.506 25. (a) $1.62 \times 10^6 \text{ N/C}$; (b) -45° 27. (a) 23.8 N/C; (b) -90° 29. 1.57 31. (a) $-5.19 \times 10^{-14} \text{ C/m}$; (b) $1.57 \times 10^{-3} \text{ N/C}$; (c) -180° ; (d) $1.52 \times 10^{-8} \text{ N/C}$; (e) $1.52 \times 10^{-8} \text{ N/C}$ 35. 0.346 m 37. 28% 39. $-5e$ 41. (a) $1.5 \times 10^3 \text{ N/C}$; (b) $2.4 \times 10^{-16} \text{ N}$; (c) up; (d) $1.6 \times 10^{-26} \text{ N}$; (e) 1.5×10^{10} 43. $3.51 \times 10^{15} \text{ m/s}^2$ 45. $6.6 \times 10^{-15} \text{ N}$ 47. (a) $1.92 \times 10^{12} \text{ m/s}^2$; (b) $1.96 \times 10^5 \text{ m/s}$ 49. (a) 0.245 N; (b) -11.3° ; (c) 108 m; (d) -21.6 m 51. $2.6 \times 10^{-10} \text{ N}$; (b) $3.1 \times 10^{-8} \text{ N}$; (c) moves to stigma 53. 27 μm 55. (a) $2.7 \times 10^6 \text{ m/s}$; (b) 1.0 kN/C 57. (a) $9.30 \times 10^{-15} \text{ C}\cdot\text{m}$; (b) $2.05 \times 10^{-11} \text{ J}$ 59. $1.22 \times 10^{-23} \text{ J}$ 61. $(1/2\pi)(pE/I)^{0.5}$ 63. (a) $8.87 \times 10^{-15} \text{ N}$; (b) 120 65. 217° 67. 61 N/C 69. (a) 47 N/C; (b) 27 N/C 71. 38 N/C 73. (a) -1.0 cm ; (b) 0; (c) 10 pC 75. $+1.00 \mu\text{C}$ 77. (a) 6.0 mm; (b) 180° 79. 9:30 81. (a) -0.029 C ; (b) repulsive forces would explode the sphere 83. (a) $-1.49 \times 10^{-26} \text{ J}$; (b) $(-1.98 \times 10^{-26} \text{ N}\cdot\text{m})\hat{k}$; (c) $3.47 \times 10^{-26} \text{ J}$ 85. (a) top row: 4, 8, 12; middle row: 5, 10, 14; bottom row: 7, 11, 16; (b) $1.63 \times 10^{-19} \text{ C}$ 87. (a) $(-1.80 \text{ N/C})\hat{j}$; (b) $(43.2 \text{ N/C})\hat{j}$; (c) $(-6.29 \text{ N/C})\hat{j}$

CHAPTER 23

CP 1. (a) $+EA$; (b) $-EA$; (c) 0; (d) 0 2. (a) 2; (b) 3; (c) 1 3. (a) equal; (b) equal; (c) equal 4. 3 and 4 tie, then 2, 1
Q 1. (a) $8 \text{ N}\cdot\text{m}^2/\text{C}$; (b) 0 3. all tie 5. all tie 7. a, c , then b and d tie (zero) 9. (a) 2, 1, 3; (b) all tie ($+4q$)
P 1. $-0.015 \text{ N}\cdot\text{m}^2/\text{C}$ 3. (a) 0; (b) $-3.92 \text{ N}\cdot\text{m}^2/\text{C}$; (c) 0; (d) 0 5. $3.01 \text{ nN}\cdot\text{m}^2/\text{C}$ 7. $2.0 \times 10^5 \text{ N}\cdot\text{m}^2/\text{C}$ 9. (a) $8.23 \text{ N}\cdot\text{m}^2/\text{C}$; (b) 72.9 pC; (c) $8.23 \text{ N}\cdot\text{m}^2/\text{C}$; (d) 72.9 pC 11. -1.70 nC 13. $3.54 \mu\text{C}$ 15. (a) 0; (b) 0.0417 17. (a) $37 \mu\text{C}$; (b) $4.1 \times 10^6 \text{ N}\cdot\text{m}^2/\text{C}$ 19. (a) $4.5 \times 10^{-7} \text{ C/m}^2$; (b) $5.1 \times 10^4 \text{ N/C}$ 21. (a) $-3.0 \times 10^{-6} \text{ C}$; (b) $+1.3 \times 10^{-5} \text{ C}$ 23. (a) $0.32 \mu\text{C}$; (b) $0.14 \mu\text{C}$ 25. $5.0 \mu\text{C/m}$ 27. $3.8 \times 10^{-8} \text{ C/m}^2$ 29. (a) 0.214 N/C; (b) inward; (c) 0.855 N/C; (d) outward; (e) $-3.40 \times 10^{-12} \text{ C}$; (f) $-3.40 \times 10^{-12} \text{ C}$ 31. (a) $2.3 \times 10^6 \text{ N/C}$; (b) outward; (c) $4.5 \times 10^5 \text{ N/C}$; (d) inward 33. (a) 0; (b) 0; (c) $(-7.91 \times 10^{-11} \text{ N/C})\hat{j}$ 35. -1.5 37. (a) $5.3 \times 10^7 \text{ N/C}$; (b) 60 N/C 39. 5.0 nC/m^2 41. 0.44 mm 43. (a) 0; (b) $1.31 \mu\text{N/C}$; (c) $3.08 \mu\text{N/C}$; (d) $3.08 \mu\text{N/C}$ 45. (a) $2.50 \times 10^4 \text{ N/C}$; (b) $1.35 \times 10^4 \text{ N/C}$ 47. -7.5 nC 49. (a) 0; (b) 56.2 mN/C; (c) 112 mN/C; (d) 49.9 mN/C; (e) 0; (f) 0; (g) -5.00 fC ; (h) 0 51. $1.79 \times 10^{-11} \text{ C/m}^2$ 53. (a) 7.78 fC; (b) 0; (c) 5.58 mN/C; (d) 22.3 mN/C 55. $6K\epsilon_0 r^3$ 57. (a) 0; (b) $2.88 \times 10^4 \text{ N/C}$; (c) 200 N/C 59. (a) 5.4 N/C; (b) 6.8 N/C 61. (a) 0; (b) $q_a/4\pi\epsilon_0 r^2$; (c) $(q_a + q_b)/4\pi\epsilon_0 r^2$ 63. -1.04 nC 65. (a) 0.125; (b) 0.500 67. (a) $+2.0 \text{ nC}$; (b) -1.2 nC ; (c) $+1.2 \text{ nC}$; (d) $+0.80 \text{ nC}$ 69. $(5.65 \times 10^4 \text{ N/C})\hat{j}$ 71. (a) $-2.53 \times 10^{-2} \text{ N}\cdot\text{m}^2/\text{C}$; (b) $+2.53 \times 10^{-2} \text{ N}\cdot\text{m}^2/\text{C}$ 75. 3.6 nC 77. (a) $+4.0 \mu\text{C}$; (b) $-4.0 \mu\text{C}$ 79. (a) 693 kg/s; (b) 693 kg/s; (c) 347 kg/s; (d) 347 kg/s; (e) 575 kg/s 81. (a) 0.25R; (b) 2.0R

CHAPTER 24

CP 1. (a) negative; (b) increase 2. (a) positive; (b) higher 3. (a) rightward; (b) 1, 2, 3, 5; positive; 4, negative; (c) 3, then 1, 2, and 5 tie, then 4 4. all tie 5. a, c (zero), b 6. (a) 2, then 1 and 3 tie; (b) 3; (c) accelerate leftward
Q 1. $-4q/4\pi\epsilon_0 d$ 3. (a) 1 and 2; (b) none; (c) no; (d) 1 and 2, yes; 3 and 4, no 5. (a) higher; (b) positive; (c) negative; (d) all tie 7. (a) 0; (b) 0; (c) 0; (d) all three quantities still 0 9. (a) 3 and 4 tie, then 1 and 2 tie; (b) 1 and 2, increase; 3 and 4, decrease
P 1. (a) $3.0 \times 10^5 \text{ C}$; (b) $3.6 \times 10^6 \text{ J}$ 3. 2.8×10^5 5. 8.8 mm 7. -32.0 V 9. (a) $1.87 \times 10^{-21} \text{ J}$; (b) -11.7 mV 11. (a) -0.268 mV ; (b) -0.681 mV 13. (a) 3.3 nC; (b) 12 nC/m^2 15. (a) 0.54 mm; (b) 790 V 17. 0.562 mV 19. (a) 6.0 cm; (b) -12.0 cm 21. 16.3 μV 23. (a) 24.3 mV; (b) 0 25. (a) -2.30 V ; (b) -1.78 V 27. 13 kV 29. 32.4 mV 31. 47.1 μV 33. 18.6 mV 35. $(-12 \text{ V/m})\hat{i} + (12 \text{ V/m})\hat{j}$ 37. 150 N/C 39. $(-4.0 \times 10^{-16} \text{ N})\hat{i} + (1.6 \times 10^{-16} \text{ N})\hat{j}$ 41. (a) 0.90 J; (b) 4.5 J 43. -0.192 pJ 45. 2.5 km/s 47. 22 km/s 49. 0.32 km/s 51. (a) $+6.0 \times 10^4 \text{ V}$; (b) $-7.8 \times 10^5 \text{ V}$; (c) 2.5 J; (d) increase; (e) same; (f) same 53. (a) 0.225 J; (b) $A 45.0 \text{ m/s}^2$, $B 22.5 \text{ m/s}^2$; (c) $A 7.75 \text{ m/s}$, $B 3.87 \text{ m/s}$ 55. $1.6 \times 10^{-9} \text{ m}$ 57. (a) 3.0 J; (b) -8.5 m 59. (a) proton; (b) 65.3 km/s 61. (a) 12; (b) 2 63. (a) $-1.8 \times 10^2 \text{ V}$; (b) 2.9 kV; (c) -8.9 kV 65. $2.5 \times 10^{-8} \text{ C}$ 67. (a) 12 kN/C; (b) 1.8 kV; (c) 5.8 cm 69. (a) 64 N/C; (b) 2.9 V; (c) 0 71. $p/2\pi\epsilon_0 r^3$ 73. (a) $3.6 \times 10^5 \text{ V}$; (b) no 75. $6.4 \times 10^8 \text{ V}$ 77. 2.90 kV 79. $7.0 \times 10^5 \text{ m/s}$ 81. (a) 1.8 cm; (b) $8.4 \times 10^5 \text{ m/s}$; (c) $2.1 \times 10^{-17} \text{ N}$; (d) positive; (e) $1.6 \times 10^{-17} \text{ N}$; (f) negative 83. (a) $+7.19 \times 10^{-10} \text{ V}$; (b) $+2.30 \times 10^{-28} \text{ J}$; (c) $+2.43 \times 10^{-29} \text{ J}$ 85. $2.30 \times 10^{-28} \text{ J}$ 87. 2.1 days 89. $2.30 \times 10^{-22} \text{ J}$ 91. $1.48 \times 10^7 \text{ m/s}$ 93. -1.92 MV

95. (a) $Q/4\pi\epsilon_0 r$; (b) $(\rho/3\epsilon_0)(1.5r_2^2 - 0.50r^2 - r_1^3 r^{-1})$,
 $\rho = Q/[(4\pi/3)(r_2^3 - r_1^3)]$; (c) $(\rho/2\epsilon_0)(r_2^2 - r_1^2)$, with ρ as in (b); (d) yes
 101. (a) 0.484 MeV; (b) 0 103. -1.7 105. (a) 38 s; (b) 280 days

CHAPTER 25

- CP 1. (a) same; (b) same 2. (a) decreases; (b) increases;
 (c) decreases 3. (a) $V, q/2$; (b) $V/2, q$
 Q 1. $a, 2, b, 1, c, 3$ 3. (a) no; (b) yes; (c) all tie 5. (a) same;
 (b) same; (c) more; (d) more 7. a , series; b , parallel; c , parallel
 9. (a) increase; (b) same; (c) increase; (d) increase; (e) increase;
 (f) increase 11. parallel, C_1 alone, C_2 alone, series
 P 1. (a) 3.5 pF; (b) 3.5 pF; (c) 57 V 3. (a) 144 pF; (b) 17.3 nC
 5. 0.280 pF 7. 6.79×10^{-4} F/m² 9. 315 mC 11. 3.16 μ F
 13. 43 pF 15. (a) 3.00 μ F; (b) 60.0 μ C; (c) 10.0 V; (d) 30.0 μ C;
 (e) 10.0 V; (f) 20.0 μ C; (g) 5.00 V; (h) 20.0 μ C 17. (a) 789 μ C;
 (b) 78.9 V 19. (a) 4.0 μ F; (b) 2.0 μ F 21. (a) 50 V; (b) 5.0×10^{-5} C;
 (c) 1.5×10^{-4} C 23. (a) 4.5×10^{14} ; (b) 1.5×10^{14} ; (c) 3.0×10^{14} ; (d)
 4.5×10^{14} ; (e) up; (f) up 25. 3.6 pC 27. (a) 9.00 μ C;
 (b) 16.0 μ C; (c) 9.00 μ C; (d) 16.0 μ C; (e) 8.40 μ C; (f) 16.8 μ C;
 (g) 10.8 μ C; (h) 14.4 μ C 29. 72 F 31. 0.27 J 33. 0.11 J/m³
 35. (a) 9.16×10^{-18} J/m³; (b) 9.16×10^{-6} J/m³; (c) 9.16×10^6 J/m³;
 (d) 9.16×10^{18} J/m³; (e) ∞ 37. (a) 16.0 V; (b) 45.1 pJ; (c) 120 pJ; (d)
 75.2 pJ 39. (a) 190 V; (b) 95 mJ 41. 81 pF/m 43. Pyrex
 45. 66 μ J 47. 0.63 m² 49. 17.3 pF 51. (a) 10 kV/m; (b) 5.0 nC;
 (c) 4.1 nC 53. (a) 89 pF; (b) 0.12 nF; (c) 11 nC; (d) 11 nC;
 (e) 10 kV/m; (f) 2.1 kV/m; (g) 88 V; (h) -0.17 μ J 55. (a) 0.107 nF;
 (b) 7.79 nC; (c) 7.45 nC 57. 45 μ C 59. 16 μ C 61. (a) 7.20 μ C;
 (b) 18.0 μ C; (c) Battery supplies charges only to plates to which it is
 connected; charges on other plates are due to electron transfers
 between plates, in accord with new distribution of voltages across
 the capacitors. So the battery does not directly supply charge on
 capacitor 4. 63. (a) 10 μ C; (b) 20 μ C 65. 1.06 nC 67. (a) 2.40 μ F;
 (b) 0.480 mC; (c) 80 V; (d) 0.480 mC; (e) 120 V 69. 4.9%
 71. (a) 0.708 pF; (b) 0.600; (c) 1.02×10^{-9} J; (d) sucked in 73. 5.3 V
 75. 40 μ F 77. (a) 200 kV/m; (b) 200 kV/m; (c) 1.77 μ C/m²;
 (d) 4.60 μ C/m²; (e) -2.83 μ C/m²

CHAPTER 26

- CP 1. 8 A, rightward 2. (a)-(c) rightward 3. a and c tie, then b
 4. device 2 5. (a) and (b) tie, then (d), then (c)
 Q 1. tie of A, B , and C , then tie of $A + B$ and $B + C$, then
 $A + B + C$ 3. (a) top-bottom, front-back, left-right; (b) top-
 bottom, front-back, left-right; (c) top-bottom, front-back, left-right;
 (d) top-bottom, front-back, left-right 5. a, b , and c all tie, then d
 7. (a) B, A, C ; (b) B, A, C 9. (a) C, B, A ; (b) all tie; (c) A, B, C ;
 (d) all tie
 P 1. (a) 1.2 kC; (b) 7.5×10^{21} 3. 6.7 μ C/m² 5. (a) 6.4 A/m²;
 (b) north; (c) cross-sectional area 7. 0.38 mm 9. 18.1 μ A
 11. (a) 1.33 A; (b) 0.666 A; (c) J_a 13. 13 min 15. 2.4 Ω
 17. 2.0×10^6 ($\Omega \cdot \text{m}$)⁻¹ 19. 2.0×10^{-8} $\Omega \cdot \text{m}$ 21. $(1.8 \times 10^3)^\circ \text{C}$
 23. 8.2×10^{-4} $\Omega \cdot \text{m}$ 25. 54 Ω 27. 3.0 29. 3.35×10^{-7} C
 31. (a) 6.00 mA; (b) 1.59×10^{-8} V; (c) 21.2 n Ω 33. (a) 38.3 mA;
 (b) 109 A/m²; (c) 1.28 cm/s; (d) 227 V/m 35. 981 k Ω 39. 150 s
 41. (a) 1.0 kW; (b) US\$0.25 43. 0.135 W 45. (a) 10.9 A;
 (b) 10.6 Ω ; (c) 4.50 MJ 47. (a) 5.85 m; (b) 10.4 m 49. (a) US\$4.46;
 (b) 144 Ω ; (c) 0.833 A 51. (a) 5.1 V; (b) 10 V; (c) 10 W; (d) 20 W
 53. (a) 28.8 Ω ; (b) 2.60×10^{19} s⁻¹ 55. 660 W 57. 28.8 kC
 59. (a) silver; (b) 51.6 n Ω 61. (a) 2.3×10^{12} ; (b) 5.0×10^3 ; (c) 10 MV
 63. 2.4 kW 65. (a) 1.37; (b) 0.730 67. (a) -8.6%; (b) smaller
 69. 146 kJ 71. (a) 250°C; (b) yes 73. 3.0×10^6 J/kg 75. 560 W

CHAPTER 27

- CP 1. (a) rightward; (b) all tie; (c) b , then a and c tie;
 (d) b , then a and c tie 2. (a) all tie; (b) R_1, R_2, R_3 3. (a) less;
 (b) greater; (c) equal 4. (a) $V/2, i$; (b) $V, i/2$ 5. (a) 1, 2, 4, 3;
 (b) 4, tie of 1 and 2, then 3
 Q 1. (a) equal; (b) more 3. parallel, R_2, R_1 , series 5. (a) series;
 (b) parallel; (c) parallel 7. (a) less; (b) less; (c) more 9. (a) same;
 (b) same; (c) less; (d) more 11. (a) all tie; (b) 1, 3, 2
 P 1. (a) 0.50 A; (b) 1.0 W; (c) 2.0 W; (d) 6.0 W; (e) 3.0 W; (f) sup-
 plied; (g) absorbed 3. (a) 14 V; (b) 1.0×10^2 W; (c) 6.0×10^2 W;
 (d) 10 V; (e) 1.0×10^2 W 5. 11 kJ 7. (a) 80 J; (b) 67 J; (c) 13 J
 9. (a) 12.0 eV; (b) 6.53 W 11. (a) 50 V; (b) 48 V; (c) negative
 13. (a) 6.9 km; (b) 20 Ω 15. 8.0 Ω 17. (a) 0.004 Ω ; (b) 1
 19. (a) 4.00 Ω ; (b) parallel 21. 5.56 A 23. (a) 50 mA; (b) 60 mA;
 (c) 9.0 V 25. 3d 27. 3.6×10^3 A 29. (a) 0.333 A; (b) right;
 (c) 720 J 31. (a) -11 V; (b) -9.0 V 33. 48.3 V 35. (a) 5.25 V;
 (b) 1.50 V; (c) 5.25 V; (d) 6.75 V 37. 1.43 Ω 39. (a) 0.150 Ω ;
 (b) 240 W 41. (a) 0.709 W; (b) 0.050 W; (c) 0.346 W; (d) 1.26 W;
 (e) -0.158 W 43. 9 45. (a) 0.67 A; (b) down; (c) 0.33 A; (d) up;
 (e) 0.33 A; (f) up; (g) 3.3 V 47. (a) 1.11 A; (b) 0.893 A; (c) 126 m
 49. (a) 0.45 A 51. (a) 55.2 mA; (b) 4.86 V; (c) 88.0 Ω ; (d) decrease
 53. -3.0% 57. 0.208 ms 59. 4.61 61. (a) 2.41 μ s; (b) 161 pF
 63. (a) 1.1 mA; (b) 0.55 mA; (c) 0.55 mA; (d) 0.82 mA; (e) 0.82 mA;
 (f) 0; (g) 4.0×10^2 V; (h) 6.0×10^2 V 65. 411 μ A 67. 0.72 M Ω
 69. (a) 0.955 μ C/s; (b) 1.08 μ W; (c) 2.74 μ W; (d) 3.82 μ W
 71. (a) 3.00 A; (b) 3.75 A; (c) 3.94 A 73. (a) 1.32×10^7 A/m²;
 (b) 8.90 V; (c) copper; (d) 1.32×10^7 A/m²; (e) 51.1 V; (f) iron
 75. (a) 3.0 kV; (b) 10 s; (c) 11 G Ω 77. (a) 85.0 Ω ; (b) 915 Ω
 81. 4.0 V 83. (a) 24.8 Ω ; (b) 14.9 k Ω 85. the cable 87. -13 μ C
 89. 20 Ω 91. (a) 3.00 A; (b) down; (c) 1.60 A; (d) down; (e) supply;
 (f) 55.2 W; (g) supply; (h) 6.40 W 93. (a) 1.0 V; (b) 50 m Ω
 95. 3 99. (a) 1.5 mA; (b) 0; (c) 1.0 mA

CHAPTER 28

- CP 1. $a, +z$; $b, -x$; $c, \vec{F}_B = 0$ 2. (a) 2, then tie of 1 and 3 (zero);
 (b) 4 3. (a) electron; (b) clockwise 4. -y 5. (a) all tie; (b) 1 and
 4 tie, then 2 and 3 tie
 Q 1. (a) no, because $\vec{v}$ and $\vec{F}_B$ must be perpendicular; (b) yes;
 (c) no, because $\vec{B}$ and $\vec{F}_B$ must be perpendicular
 3. (a) $+z$ and $-z$ tie, then $+y$ and $-y$ tie, then $+x$ and $-x$ tie (zero);
 (b) $+y$ 5. (a) $\vec{F}_E$; (b) $\vec{F}_B$ 7. (a) $\vec{B}_1$; (b) $\vec{B}_1$ into page, $\vec{B}_2$ out of page;
 (c) less 9. (a) positive; (b) $2 \rightarrow 1$ and $2 \rightarrow 4$ tie, then $2 \rightarrow 3$ (which is
 zero) 11. (a) negative; (b) equal; (c) equal; (d) half-circle
 P 1. (a) 400 km/s; (b) 835 eV 3. (a) $(6.2 \times 10^{-14} \text{ N})\hat{k}$;
 (b) $(-6.2 \times 10^{-14} \text{ N})\hat{k}$ 5. -2.0 T 7. $(-11.4 \text{ V/m})\hat{j} - (6.00 \text{ V/m})\hat{j} +$
 $(4.80 \text{ V/m})\hat{k}$ 9. $(-0.267 \text{ mT})\hat{k}$ 11. 0.68 MV/m 13. 7.4 μ V
 15. (a) $(-600 \text{ mV/m})\hat{k}$; (b) 1.20 V 17. (a) 2.60×10^6 m/s;
 (b) 0.109 μ s; (c) 0.140 MeV; (d) 70.0 kV 19. 1.2×10^{-9} kg/C
 21. (a) 2.05×10^7 m/s; (b) 467 μ T; (c) 13.1 MHz; (d) 76.3 ns
 23. 21.1 μ T 25. (a) 0.978 MHz; (b) 96.4 cm 27. (a) 495 mT;
 (b) 22.7 mA; (c) 8.17 MJ 29. 65.3 km/s 31. 5.07 ns
 33. (a) 0.358 ns; (b) 0.166 mm; (c) 1.51 mm 35. (a) 200 eV;
 (b) 20.0 keV; (c) 0.499% 37. 2.4×10^2 m 39. (a) 28.2 N;
 (b) horizontally west 41. (a) 467 mA; (b) right 43. (a) 0; (b) 0.138 N;
 (c) 0.138 N; (d) 0 45. $(-2.50 \text{ mN})\hat{j} + (0.750 \text{ mN})\hat{k}$ 47. (a) 0.12 T;
 (b) 31° 49. $(-4.3 \times 10^{-3} \text{ N}\cdot\text{m})\hat{j}$ 51. 2.45 A 55. (a) 2.86 A $\cdot$ m²;
 (b) 1.10 A $\cdot$ m² 57. (a) 12.7 A; (b) 0.0805 N $\cdot$ m 59. (a) 0.30 A $\cdot$ m²;
 (b) 0.024 N $\cdot$ m 61. (a) -72.0 μ J; (b) $(96.0\hat{i} + 48.0\hat{k}) \mu$ N $\cdot$ m
 63. (a) $(-9.7 \times 10^{-4} \text{ N}\cdot\text{m})\hat{i} - (7.2 \times 10^{-4} \text{ N}\cdot\text{m})\hat{j} + (8.0 \times 10^{-4} \text{ N}\cdot\text{m})\hat{k}$;
 (b) -6.0×10^{-4} J 65. (a) 90°; (b) 1; (c) 1.28×10^{-7} N $\cdot$ m

67. (a) 20 min; (b) $5.9 \times 10^{-2} \text{ N} \cdot \text{m}$ 69. 8.2 mm 71. 127 μ
 73. (a) $6.3 \times 10^{14} \text{ m/s}^2$; (b) 3.0 mm 75. (a) 1.4; (b) 1.0
 77. $(-500 \text{ V/m})\hat{j}$ 79. (a) 0.50; (b) 0.50; (c) 14 cm; (d) 14 cm
 81. $(0.80\hat{j} - 1.1\hat{k}) \text{ mN}$ 83. -40 mC 85. (a) $(12.8\hat{i} + 6.41\hat{j}) \times 10^{-22} \text{ N}$; (b) 90° ; (c) 173°

CHAPTER 29

- CP 1. *b, c, a* 2. *d*, tie of *a* and *c*, then *b* 3. *d, a*, tie of *b* and *c* (zero)
 Q 1. *c, a, b* 3. *c, d*, then *a* and *b* tie (zero) 5. *a, c, b*
 7. *c* and *d* tie, then *b, a* 9. *b, a, d, c* (zero) 11. (a) 1, 3, 2; (b) less
 P 1. (a) $3.3 \mu\text{T}$; (b) yes 3. (a) 16 A; (b) east 5. (a) 1.0 mT;
 (b) out; (c) 0.80 mT; (d) out 7. (a) $0.102 \mu\text{T}$; (b) out
 9. (a) opposite; (b) 30 A 11. (a) 4.3 A; (b) out 13. 50.3 nT
 15. (a) $1.7 \mu\text{T}$; (b) into; (c) $6.7 \mu\text{T}$; (d) into 17. 132 nT 19. 5.0 μT
 21. 256 nT 23. $(-7.75 \times 10^{-23} \text{ N})\hat{i}$ 25. 2.00 rad 27. 61.3 mA
 29. $(80 \mu\text{T})\hat{j}$ 31. (a) $20 \mu\text{T}$; (b) into 33. $(22.3 \text{ pT})\hat{j}$ 35. 88.4 pN/m
 37. $(-125 \mu\text{N/m})\hat{i} + (41.7 \mu\text{N/m})\hat{j}$ 39. 800 nN/m
 41. $(3.20 \text{ mN})\hat{j}$ 43. (a) 0; (b) 0.850 mT; (c) 1.70 mT; (d) 0.850 mT
 45. (a) $-2.5 \mu\text{T} \cdot \text{m}$; (b) 0 47. (a) 0; (b) $0.10 \mu\text{T}$; (c) $0.40 \mu\text{T}$
 49. (a) 533 μT ; (b) 400 μT 51. 0.30 mT 53. 0.272 A
 55. (a) 4.77 cm; (b) 35.5 μT 57. (a) $2.4 \text{ A} \cdot \text{m}^2$; (b) 46 cm
 59. $0.47 \text{ A} \cdot \text{m}^2$ 61. (a) $79 \mu\text{T}$; (b) $1.1 \times 10^{-6} \text{ N} \cdot \text{m}$ 63. (a) $(0.060 \text{ A} \cdot \text{m}^2)\hat{j}$;
 (b) $(96 \text{ pT})\hat{j}$ 65. 1.28 mm 69. (a) 15 A; (b) $-z$ 71. 7.7 mT
 73. (a) $15.3 \mu\text{T}$ 75. (a) $(0.24\hat{i}) \text{ nT}$; (b) 0; (c) $(-43\hat{k}) \text{ pT}$; (d) $(0.14\hat{k}) \text{ nT}$
 79. (a) 4.8 mT; (b) 0.93 mT; (c) 0 83. $(-0.20 \text{ mT})\hat{k}$
 87. (a) $\mu_0 i r / 2\pi c^2$; (b) $\mu_0 i / 2\pi r$; (c) $\mu_0 i (a^2 - r^2) / 2\pi (a^2 - b^2) r$; (d) 0

CHAPTER 30

- CP 1. *b*, then *d* and *e* tie, and then *a* and *c* tie (zero) 2. *a* and *b*
 tie, then *c* (zero) 3. *c* and *d* tie, then *a* and *b* tie 4. *b*, out; *c*, out; *d*,
 into; *e*, into 5. *d* and *e* 6. (a) 2, 3, 1 (zero); (b) 2, 3, 1
 7. *a* and *b* tie, then *c*
 Q 1. out 3. (a) all tie (zero); (b) 2, then 1 and 3 tie (zero) 5. *d*
 and *c* tie, then *b, a* 7. (a) more; (b) same; (c) same; (d) same (zero)
 9. (a) all tie (zero); (b) 1 and 2 tie, then 3; (c) all tie (zero)
 P 1. 0 3. 30 mA 5. 0 7. (a) 31 mV; (b) left 9. 0.198 mV
 11. (b) 0.796 m^2 13. 29.5 mC 15. (a) 21.7 V; (b) counterclock-
 wise 17. (a) $1.26 \times 10^{-4} \text{ T}$; (b) 0; (c) $1.26 \times 10^{-4} \text{ T}$; (d) yes;
 (e) $5.04 \times 10^{-8} \text{ V}$ 19. 5.50 kV 21. (a) 40 Hz; (b) 3.2 mV
 23. (a) $\mu_0 i R^2 \pi r^2 / 2x^3$; (b) $3\mu_0 i \pi R^2 r^2 v / 2x^4$; (c) counterclockwise
 25. (a) $13 \mu\text{Wb/m}$; (b) 17%; (c) 0 27. (a) $80 \mu\text{V}$; (b) clockwise
 29. (a) 48.1 mV; (b) 2.67 mA; (c) 0.129 mW 31. 3.68 μW
 33. (a) $240 \mu\text{V}$; (b) 0.600 mA; (c) $0.144 \mu\text{W}$; (d) $2.87 \times 10^{-8} \text{ N}$;
 (e) $0.144 \mu\text{W}$ 35. (a) 0.60 V; (b) up; (c) 1.5 A; (d) clockwise;
 (e) 0.90 W; (f) 0.18 N; (g) 0.90 W 37. (a) $71.5 \mu\text{V/m}$; (b) $143 \mu\text{V/m}$
 39. 0.15 V/m 41. (a) 2.45 mWb; (b) 0.645 mH 43. 1.81 $\mu\text{H/m}$
 45. (a) decreasing; (b) 0.68 mH 47. (b) $L_{\text{eq}} = \sum L_j$, sum from $j = 1$
 to $j = N$ 49. 59.3 mH 51. 46 Ω 53. (a) 8.45 ns; (b) 7.37 mA
 55. 6.91 57. (a) 1.5 s 59. (a) $i[1 - \exp(-Rt/L)]$; (b) $(L/R) \ln 2$
 61. (a) 97.9 H; (b) 0.196 mJ 63. 25.6 ms 65. (a) 18.7 J; (b) 5.10 J;
 (c) 13.6 J 67. (a) 34.2 J/m^3 ; (b) 49.4 mJ 69. $1.5 \times 10^8 \text{ V/m}$
 71. (a) 1.0 J/m^3 ; (b) $4.8 \times 10^{-15} \text{ J/m}^3$ 73. (a) 1.67 mH; (b) 6.00 mWb
 75. 13 μH 77. (b) have the turns of the two solenoids wrapped in
 opposite directions 79. (a) 2.0 A; (b) 0; (c) 2.0 A; (d) 0; (e) 10 V;
 (f) 2.0 A/s; (g) 2.0 A; (h) 1.0 A; (i) 3.0 A; (j) 10 V; (k) 0; (l) 0
 81. (a) $10 \mu\text{T}$; (b) out; (c) $3.3 \mu\text{T}$; (d) out 83. 0.520 ms
 85. (a) $(4.4 \times 10^7 \text{ m/s}^2)\hat{j}$; (b) 0; (c) $(-4.4 \times 10^7 \text{ m/s}^2)\hat{j}$
 87. (a) 0.40 V; (b) 20 A 89. (a) 10 A; (b) $1.0 \times 10^2 \text{ J}$ 91. (a) 0;
 (b) $8.0 \times 10^2 \text{ A/s}$; (c) 1.8 mA; (d) $4.4 \times 10^2 \text{ A/s}$; (e) 4.0 mA; (f) 0
 93. 1.15 W 95. (a) 20 A/s; (b) 0.75 A 97. 12 A/s

CHAPTER 31

- CP 1. (a) $T/2$; (b) T ; (c) $T/2$; (d) $T/4$ 2. (a) 5 V; (b) $150 \mu\text{J}$
 3. (a) remains the same; (b) remains the same 4. (a) *C, B, A*; (b) 1,
A; 2, *B*; 3, *S*; 4, *C*; (c) *A* 5. (a) remains the same; (b) increases;
 (c) remains the same; (d) decreases 6. (a) 1, lags; 2, leads; 3, in
 phase; (b) 3 ($\omega_d = \omega$ when $X_L = X_C$) 7. (a) increase (circuit is
 mainly capacitive; increase *C* to decrease X_C to be closer to reso-
 nance for maximum P_{avg}); (b) closer 8. (a) greater; (b) step-up
 Q 1. *b, a, c* 3. (a) $T/4$; (b) $T/4$; (c) $T/2$; (d) $T/2$ 5. *c, b, a* 7. *a*
 inductor; *b* resistor; *c* capacitor 9. (a) positive; (b) decreased (to
 decrease X_L and get closer to resonance); (c) decreased (to increase
 X_C and get closer to resonance) 11. (a) rightward, increase (X_L
 increases, closer to resonance); (b) rightward, increase (X_C decreases,
 closer to resonance); (c) rightward, increase (ω_d/ω increases, closer to
 resonance)
 P 1. (a) $1.17 \mu\text{J}$; (b) 5.58 mA 3. (a) 6.00 μs ; (b) 167 kHz; (c) 3.00 μs
 5. 45.2 mA 7. (a) 1.25 kg; (b) 372 N/m; (c) $1.75 \times 10^{-4} \text{ m}$;
 (d) 3.02 mm/s 9. $7.0 \times 10^{-4} \text{ s}$ 11. (a) 6.0; (b) 36 pF; (c) 0.22 mH
 13. (a) 0.180 mC; (b) 70.7 μs ; (c) 66.7 W 15. (a) 3.0 nC; (b) 1.7 mA;
 (c) 4.5 nJ 17. (a) 275 Hz; (b) 365 mA 21. (a) 356 μs ; (b) 2.50 mH;
 (c) 3.20 mJ 23. (a) 1.98 μJ ; (b) 5.56 μC ; (c) 12.6 mA; (d) -46.9° ;
 (e) $+46.9^\circ$ 25. 8.66 m Ω 29. (a) 95.5 mA; (b) 11.9 mA
 31. (a) 0.65 kHz; (b) 24 Ω 33. (a) 6.73 ms; (b) 11.2 ms;
 (c) inductor; (d) 138 mH 35. 89 Ω 37. 7.61 A 39. (a) 267 Ω ;
 (b) -41.5° ; (c) 135 mA 41. (a) 206 Ω ; (b) 13.7° ; (c) 175 mA
 43. (a) 218 Ω ; (b) 23.4° ; (c) 165 mA 45. (a) yes; (b) 1.0 kV
 47. (a) 224 rad/s; (b) 6.00 A; (c) 219 rad/s; (d) 228 rad/s; (e) 0.040
 49. (a) 796 Hz; (b) no change; (c) decreased; (d) increased
 53. (a) 12.1 Ω ; (b) 1.19 kW 55. 1.84 A 57. (a) 117 μF ; (b) 0;
 (c) 90.0 W; (d) 0° ; (e) 1; (f) 0; (g) -90° ; (h) 0 59. (a) 2.59 A;
 (b) 38.8 V; (c) 159 V; (d) 224 V; (e) 64.2 V; (f) 75.0 V; (g) 100 W; (h) 0;
 (i) 0 61. (a) 0.743; (b) lead; (c) capacitive; (d) no; (e) yes; (f) no;
 (g) yes; (h) 33.4 W 63. (a) 2.4 V; (b) 3.2 mA; (c) 0.16 A
 65. (a) 1.9 V; (b) 5.9 W; (c) 19 V; (d) $5.9 \times 10^2 \text{ W}$; (e) 0.19 kV; (f) 59 kW
 67. (a) 6.73 ms; (b) 2.24 ms; (c) capacitor; (d) 59.0 μF
 69. (a) -0.405 rad ; (b) 2.76 A; (c) capacitive 71. (a) 64.0 Ω ;
 (b) 50.9 Ω ; (c) capacitive 73. (a) 2.41 μH ; (b) 21.4 pJ; (c) 82.2 nC
 75. (a) 39.1 Ω ; (b) 21.7 Ω ; (c) capacitive 79. (a) 0.577 Ω ; (b) 0.152
 81. (a) 45.0° ; (b) 70.7 Ω 83. 1.84 kHz 85. (a) 0.689 μH ;
 (b) 17.9 pJ; (c) 0.110 μC 87. (a) 165 Ω ; (b) 313 mH; (c) 14.9 μF

CHAPTER 32

- CP 1. *d, b, c, a* (zero) 2. *a, c, b, d* (zero) 3. tie of *b, c*, and *d*, then *a*
 4. (a) 2; (b) 1 5. (a) away; (b) away; (c) less 6. (a) toward;
 (b) toward; (c) less
 Q 1. 1 *a*, 2 *b*, 3 *c* and *d* 3. *a*, decreasing; *b*, decreasing
 5. supplied 7. (a) *a* and *b* tie, then *c, d*; (b) none (because plate
 lacks circular symmetry, $\vec{B}$ not tangent to any circular loop);
 (c) none 9. (a) 1 up, 2 up, 3 down; (b) 1 down, 2 up, 3 zero
 11. (a) 1, 3, 2; (b) 2
 P 1. +3 Wb 3. (a) 47.4 μWb ; (b) inward 5. $2.4 \times 10^{13} \text{ V/m} \cdot \text{s}$
 7. (a) $1.18 \times 10^{-19} \text{ T}$; (b) $1.06 \times 10^{-19} \text{ T}$ 9. (a) $5.01 \times 10^{-22} \text{ T}$;
 (b) $4.51 \times 10^{-22} \text{ T}$ 11. (a) 1.9 pT 13. $7.5 \times 10^5 \text{ V/s}$
 17. (a) 0.324 V/m; (b) $2.87 \times 10^{-16} \text{ A}$; (c) 2.87×10^{-18}
 19. (a) 75.4 nT; (b) 67.9 nT 21. (a) 27.9 nT; (b) 15.1 nT
 23. (a) 2.0 A; (b) $2.3 \times 10^{11} \text{ V/m} \cdot \text{s}$; (c) 0.50 A; (d) $0.63 \mu\text{T} \cdot \text{m}$
 25. (a) 0.63 μT ; (b) $2.3 \times 10^{12} \text{ V/m} \cdot \text{s}$ 27. (a) 0.71 A; (b) 0; (c) 2.8 A
 29. (a) 7.60 μA ; (b) 859 kV $\cdot \text{m/s}$; (c) 3.39 mm; (d) 5.16 pT 31. 55 μT
 33. (a) 0; (b) 0; (c) 0; (d) $\pm 3.2 \times 10^{-25} \text{ J}$; (e) $-3.2 \times 10^{-34} \text{ J} \cdot \text{s}$;
 (f) $2.8 \times 10^{-23} \text{ J/T}$; (g) $-9.7 \times 10^{-25} \text{ J}$; (h) $\pm 3.2 \times 10^{-25} \text{ J}$
 35. (a) $-9.3 \times 10^{-24} \text{ J/T}$; (b) $1.9 \times 10^{-23} \text{ J/T}$ 37. (b) $+x$;

(c) clockwise; (d) $+x$ 39. yes 41. 20.8 mJ/T 43. (b) K_i/B ;
(c) $-z$; (d) 0.31 kA/m 47. (a) 1.8×10^2 km; (b) 2.3×10^{-5}
49. (a) $3.0 \mu\text{T}$; (b) 5.6×10^{-10} eV 51. 5.15×10^{-24} A·m²
53. (a) 0.14 A; (b) $79 \mu\text{C}$ 55. (a) 6.3×10^8 A; (b) yes; (c) no
57. 0.84 kJ/T 59. (a) $(1.2 \times 10^{-13} \text{ T}) \exp[-t/(0.012 \text{ s})]$;
(b) 5.9×10^{-15} T 63. (a) 27.5 mm; (b) 110 mm 65. 8.0 A
67. (a) -8.8×10^{15} V/m·s; (b) 5.9×10^{-7} T·m 69. (b) sign is
minus; (c) no, because there is compensating positive flux through
open end nearer to magnet 71. (b) $-x$; (c) counterclockwise;
(d) $-x$ 73. (a) 7; (b) 7; (c) $3h/2\pi$; (d) $3eh/4\pi m$; (e) $3.5h/2\pi$;
(f) 8 75. (a) 9; (b) 3.71×10^{-23} J/T; (c) $+9.27 \times 10^{-24}$ J;
(d) -9.27×10^{-24} J

CHAPTER 33

CP 1. (a) (Use Fig. 33-5.) On right side of rectangle, $\vec{E}$ is in
negative y direction; on left side, $\vec{E} + d\vec{E}$ is greater and in same
direction; (b) $\vec{E}$ is downward. On right side, $\vec{B}$ is in negative z
direction; on left side, $\vec{B} + d\vec{B}$ is greater and in same direction.
2. positive direction of x 3. (a) same; (b) decrease 4. a, d, b, c
(zero) 5. a
Q 1. (a) positive direction of z ; (b) x 3. (a) same; (b) increase;
(c) decrease 5. (a) and (b) $A=1, n=4, \theta=30^\circ$ 7. a, b, c 9. B 11. none
P 1. 7.49 GHz 3. (a) 515 nm; (b) 610 nm; (c) 555 nm;
(d) 5.41×10^{14} Hz; (e) 1.85×10^{-15} s 5. 5.0×10^{-21} H 7. 1.2 MW/m²
9. 0.10 MJ 11. (a) 6.7 nT; (b) y ; (c) negative direction of y
13. (a) 1.03 kV/m; (b) $3.43 \mu\text{T}$ 15. (a) 87 mV/m; (b) 0.29 nT;
(c) 6.3 kW 17. (a) 6.7 nT; (b) 5.3 mW/m^2 ; (c) 6.7 W 19. 1.0×10^7 Pa
21. 5.9×10^{-8} Pa 23. (a) 4.68×10^{11} W; (b) any chance
disturbance could move sphere from directly above source—the
two force vectors no longer along the same axis 27. (a) 1.0×10^8 Hz;
(b) 6.3×10^8 rad/s; (c) 2.1 m^{-1} ; (d) $1.0 \mu\text{T}$; (e) z ; (f) 1.2×10^2 W/m²;
(g) 8.0×10^{-7} N; (h) 4.0×10^{-7} Pa 29. 1.9 mm/s 31. (a) 0.17 m/s;
(b) toward the Sun 33. 3.1% 35. 4.4 W/m^2 37. (a) 2 sheets;
(b) 5 sheets 39. (a) 1.9 V/m; (b) 1.7×10^{-11} Pa 41. 20° or 70°
43. 0.67 45. 1.26 47. 1.48 49. 180° 51. (a) 56.9° ; (b) 35.3°
55. 1.07 m 57. 182 cm 59. (a) 48.9° ; (b) 29.0° 61. (a) 26.8° ;
(b) yes 63. (a) $(1 + \sin^2 \theta)^{0.5}$; (b) $2^{0.5}$; (c) yes; (d) no 65. 23.2°
67. (a) 1.39; (b) 28.1° ; (c) no 69. 49.0° 71. (a) 0.50 ms; (b) 8.4 min;
(c) 2.4 h; (d) 5446 B.C. 73. (a) $(16.7 \text{ nT}) \sin[(1.00 \times 10^6 \text{ m}^{-1})z +$
 $(3.00 \times 10^{14} \text{ s}^{-1})t]$; (b) $6.28 \mu\text{m}$; (c) 20.9 fs; (d) 33.2 mW/m^2 ; (e) x ;
(f) infrared 75. 1.22 77. (c) 137.6° (d) 139.4° (e) 1.7°
81. (a) z axis; (b) 7.5×10^{14} Hz; (c) 1.9 kW/m^2 83. (a) white;
(b) white dominated by red end; (c) no refracted light
85. 1.5×10^{-9} m/s² 87. (a) $3.5 \mu\text{W/m}^2$; (b) 0.78 W/m^2 ;
(c) 1.5×10^{-17} W/m²; (d) 1.1×10^{-7} V/m; (e) 0.25 fT 89. (a) 55.8° ;
(b) 55.5° 91. (a) 83 W/m^2 ; (b) 1.7 MW 93. 35°

CHAPTER 34

CP 1. 0.2d, 1.8d, 2.2d 2. (a) real; (b) inverted; (c) same
3. (a) e ; (b) virtual, same 4. virtual, same as object, diverging
Q 1. (a) a ; (b) c 3. (a) a and c ; (b) three times; (c) you
5. convex 7. (a) all but variation 2; (b) 1, 3, 4; right, inverted; 5, 6;
left, same 9. d (infinite), tie of a and b , then c
P 1. 9.10 m 3. 1.11 5. 351 cm 7. 10.5 cm 9. (a) $+24$ cm;
(b) $+36$ cm; (c) -2.0 ; (d) R; (e) I; (f) same 11. (a) -20 cm;
(b) -4.4 cm; (c) $+0.56$; (d) V; (e) NI; (f) opposite 13. (a) $+36$ cm;
(b) -36 cm; (c) $+3.0$; (d) V; (e) NI; (f) opposite 15. (a) -16 cm;
(b) -4.4 cm; (c) $+0.44$; (d) V; (e) NI; (f) opposite 17. (b) plus;
(c) $+40$ cm; (e) -20 cm; (f) $+2.0$; (g) V; (h) NI; (i) opposite
19. (a) convex; (b) -20 cm; (d) $+20$ cm; (f) $+0.50$; (g) V; (h) NI;

(i) opposite 21. (a) concave; (c) $+40$ cm; (e) $+60$ cm; (f) -2.0 ;
(g) R; (h) I; (i) same 23. (a) convex; (b) minus; (c) -60 cm;
(d) $+1.2$ m; (e) -24 cm; (g) V; (h) NI; (i) opposite 25. (a) concave;
(b) $+8.6$ cm; (c) $+17$ cm; (e) $+12$ cm; (f) minus; (g) R; (i) same
27. (a) convex; (c) -60 cm; (d) $+30$ cm; (f) $+0.50$; (g) V; (h) NI;
(i) opposite 29. (b) -20 cm; (c) minus; (d) $+5.0$ cm; (e) minus;
(f) $+0.80$ (g) V; (h) NI; (i) opposite 31. (b) 0.56 cm/s; (c) 11 m/s;
(d) 6.7 cm/s 33. (c) -33 cm; (e) V; (f) same 35. (d) -26 cm; (e) V;
(f) same 37. (c) $+30$ cm; (e) V; (f) same 39. (a) 2.00; (b) none
41. (a) $+40$ cm; (b) ∞ 43. 5.0 mm 45. 1.86 mm 47. (a) 45 mm;
(b) 90 mm 49. 22 cm 51. (a) -48 cm; (b) $+4.0$; (c) V; (d) NI;
(e) same 53. (a) -4.8 cm; (b) $+0.60$; (c) V; (d) NI; (e) same
55. (a) -8.6 cm; (b) $+0.39$; (c) V; (d) NI; (e) same 57. (a) $+36$ cm;
(b) -0.80 ; (c) R; (d) I; (e) opposite 59. (a) $+55$ cm; (b) -0.74 ;
(c) R; (d) I; (e) opposite 61. (a) -18 cm; (b) $+0.76$; (c) V; (d) NI;
(e) same 63. (a) -30 cm; (b) $+0.86$; (c) V; (d) NI; (e) same
65. (a) -7.5 cm; (b) $+0.75$; (c) V; (d) NI; (e) same 67. (a) $+84$ cm;
(b) -1.4 ; (c) R; (d) I; (e) opposite 69. (a) C; (d) -10 cm; (e) $+2.0$;
(f) V; (g) NI; (h) same 71. (a) D; (b) -5.3 cm; (d) -4.0 cm; (f) V; (g)
NI; (h) same 73. (a) C; (b) $+3.3$ cm; (d) $+5.0$ cm; (f) R; (g) I;
(h) opposite 75. (a) D; (b) minus; (d) -3.3 cm; (e) $+0.67$; (f) V;
(g) NI 77. (a) C; (b) $+80$ cm; (d) -20 cm; (f) V; (g) NI; (h) same
79. (a) C; (b) plus; (d) -13 cm; (e) $+1.7$; (f) V; (g) NI; (h) same
81. (a) $+24$ cm; (b) $+6.0$; (c) R; (d) NI; (e) opposite
83. (a) $+3.1$ cm; (b) -0.31 ; (c) R; (d) I; (e) opposite 85. (a) -4.6 cm;
(b) $+0.69$; (c) V; (d) NI; (e) same 87. (a) -5.5 cm; (b) $+0.12$; (c) V;
(d) NI; (e) same 89. (a) 13.0 cm; (b) 5.23 cm; (c) -3.25 ; (d) 3.13;
(e) -10.2 91. (a) 2.35 cm; (b) decrease 93. (a) 3.5; (b) 2.5
95. (a) $+8.6$ cm; (b) $+2.6$; (c) R; (d) NI; (e) opposite
97. (a) $+7.5$ cm; (b) -0.75 ; (c) R; (d) I; (e) opposite 99. (a) $+24$ cm;
(b) -0.58 ; (c) R; (d) I; (e) opposite 105. (a) 3.00 cm; (b) 2.33 cm
107. (a) 40 cm; (b) 20 cm; (c) -40 cm; (d) 40 cm 109. (a) 20 cm;
(b) 15 cm 111. (a) 6.0 mm; (b) 1.6 kW/m^2 ; (c) 4.0 cm

CHAPTER 35

CP 1. b (least n), c, a 2. (a) top; (b) bright intermediate illumina-
tion (phase difference is 2.1 wavelengths) 3. (a) $3\lambda, 3$; (b) $2.5\lambda, 2.5$
4. a and d tie (amplitude of resultant wave is $4E_0$), then b and c tie
(amplitude of resultant wave is $2E_0$) 5. (a) 1 and 4; (b) 1 and 4
Q 1. (a) decrease; (b) decrease; (c) decrease; (d) blue 3. (a) $2d$;
(b) (odd number) $\lambda/2$; (c) $\lambda/4$ 5. (a) intermediate closer to
maximum, $m=2$; (b) minimum, $m=3$; (c) intermediate closer to
maximum, $m=2$; (d) maximum, $m=1$ 7. (a) maximum;
(b) minimum; (c) alternates 9. (a) peak; (b) valley 11. c, d
P 1. (a) 155 nm; (b) 310 nm 3. (a) $3.60 \mu\text{m}$; (b) intermediate
closer to fully constructive 5. 4.55×10^7 m/s 7. 1.56
9. (a) $1.55 \mu\text{m}$; (b) $4.65 \mu\text{m}$ 11. (a) 1.70; (b) 1.70; (c) 1.30;
(d) all tie 13. (a) 0.833; (b) intermediate closer to fully
constructive 15. 648 nm 17. 16 19. 2.25 mm 21. 72 μm
23. 0 25. $7.88 \mu\text{m}$ 27. $6.64 \mu\text{m}$ 29. 2.65 31. $27 \sin(\omega t + 8.5^\circ)$
33. $(17.1 \mu\text{V/m}) \sin[(2.0 \times 10^{14} \text{ rad/s})t]$ 35. 120 nm 37. 70.0 nm
39. (a) 0.117 μm ; (b) 0.352 μm 41. 161 nm 43. 560 nm
45. 478 nm 47. 509 nm 49. 273 nm 51. 409 nm 53. 338 nm
55. (a) 552 nm; (b) 442 nm 57. 608 nm 59. 528 nm 61. 455 nm
63. 248 nm 65. 339 nm 67. 329 nm 69. $1.89 \mu\text{m}$ 71. 0.012°
73. 140 75. $[(m + \frac{1}{2})\lambda R]^{0.5}$, for $m=0, 1, 2, \dots$ 77. 1.00 m
79. 588 nm 81. 1.00030 83. (a) 50.0 nm; (b) 36.2 nm 85. 0.23°
87. (a) 1500 nm; (b) 2250 nm; (c) 0.80 89. $x = (D/2a)(m + 0.5)\lambda$,
for $m=0, 1, 2, \dots$ 91. (a) 22° ; (b) refraction reduces θ 93. 600 nm
95. (a) $1.75 \mu\text{m}$; (b) 4.8 mm 97. $I_m \cos^2(2\pi x/\lambda)$ 99. (a) 42.0 ps;
(b) 42.3 ps; (c) 43.2 ps; (d) 41.8 ps; (e) 4 101. 33 μm

CHAPTER 36

- CP** 1. (a) expand; (b) expand 2. (a) second side maximum; (b) 2.5 3. (a) red; (b) violet 4. diminish 5. (a) left; (b) less
- Q** 1. (a) $m = 5$ minimum; (b) (approximately) maximum between the $m = 4$ and $m = 5$ minima 3. (a) A, B, C ; (b) A, B, C
5. (a) 1 and 3 tie, then 2 and 4 tie; (b) 1 and 2 tie, then 3 and 4 tie
7. (a) larger; (b) red 9. (a) decrease; (b) same; (c) remain in place
11. (a) A ; (b) left; (c) left; (d) right
- P** 1. (a) 2.5 mm; (b) 2.2×10^{-4} rad 3. (a) 70 cm; (b) 1.0 mm
5. (a) 700 nm; (b) 4; (c) 6 7. $60.4 \mu\text{m}$ 9. 1.77 mm
11. 160° 13. (a) 0.18° ; (b) 0.46 rad ; (c) 0.93 15. (d) 52.5° ; (e) 10.1° ; (f) 5.06° 17. (b) 0; (c) -0.500 ; (d) 4.493 rad ; (e) 0.930 ; (f) 7.725 rad ; (g) 1.96 19. (a) 19 cm; (b) larger 21. (a) $1.1 \times 10^4 \text{ km}$; (b) 11 km 23. (a) $1.3 \times 10^{-4} \text{ rad}$; (b) 10 km 25. 50 m
27. $1.6 \times 10^3 \text{ km}$ 29. (a) $8.8 \times 10^{-7} \text{ rad}$; (b) $8.4 \times 10^7 \text{ km}$; (c) 0.025 mm
31. (a) 0.346° ; (b) 0.97° 33. (a) 17.1 m; (b) 1.37×10^{-10}
35. 5 37. 3 39. (a) $5.0 \mu\text{m}$; (b) $20 \mu\text{m}$ 41. (a) 7.43×10^{-3} ; (b) between the $m = 6$ minimum (the seventh one) and the $m = 7$ maximum (the seventh side maximum); (c) between the $m = 3$ minimum (the third one) and the $m = 4$ minimum (the fourth one)
43. (a) 9; (b) 0.255 45. (a) 62.1° ; (b) 45.0° ; (c) 32.0° 47. 3
49. (a) $6.0 \mu\text{m}$; (b) $1.5 \mu\text{m}$; (c) 9; (d) 7; (e) 6 51. (a) 2.1° ; (b) 21° ; (c) 11 53. (a) 470 nm; (b) 560 nm 55. 3.65×10^3
57. (a) $0.032^\circ/\text{nm}$; (b) 4.0×10^4 ; (c) $0.076^\circ/\text{nm}$; (d) 8.0×10^4 ; (e) $0.24^\circ/\text{nm}$; (f) 1.2×10^5 59. 0.15 nm 61. (a) $10 \mu\text{m}$; (b) 3.3 mm
63. $1.09 \times 10^3 \text{ rulings/mm}$ 65. (a) 0.17 nm; (b) 0.13 nm
67. (a) 25 pm; (b) 38 pm 69. 0.26 nm 71. (a) 15.3° ; (b) 30.6° ; (c) 3.1° ; (d) 37.8° 73. (a) $0.7071a_0$; (b) $0.4472a_0$; (c) $0.3162a_0$; (d) $0.2774a_0$; (e) $0.2425a_0$ 75. (a) 625 nm; (b) 500 nm; (c) 416 nm
77. 3.0 mm 83. (a) 13; (b) 6 85. 59.5 pm 87. 4.9 km 89. 1.36×10^4
91. 2 93. 4.7 cm 97. 36 cm 99. (a) fourth; (b) seventh

CHAPTER 37

- CP** 1. (a) same (speed of light postulate); (b) no (the start and end of the flight are spatially separated); (c) no (because his measurement is not a proper time) 2. (a) Eq. 2; (b) $+0.90c$; (c) 25 ns; (d) -7.0 m 3. (a) right; (b) more 4. (a) equal; (b) less
- Q** 1. c 3. b 5. (a) C' ; (b) C'_1 7. (a) 4 s; (b) 3 s; (c) 5 s; (d) 4 s; (e) 10 s 9. (a) a tie of 3, 4, and 6, then a tie of 1, 2, and 5; (b) 1, then a tie of 2 and 3, then 4, then a tie of 5 and 6; (c) 1, 2, 3, 4, 5, 6; (d) 2 and 4; (e) 1, 2, 5 11. (a) 3, tie of 1 and 2, then 4; (b) 4, tie of 1 and 2, then 3; (c) 1, 4, 2, 3
- P** 1. 0.990 50 3. (a) 0.999 999 50 5. 0.446 ps 7. $2.68 \times 10^3 \text{ y}$
9. (a) 87.4 m; (b) 394 ns 11. 1.32 m 13. (a) 26.26 y; (b) 52.26 y; (c) 3.705 y 15. (a) 0.999 999 15; (b) 30 ly 17. (a) 138 km; (b) $-374 \mu\text{s}$ 19. (a) $25.8 \mu\text{s}$; (b) small flash 21. (a) $\gamma[1.00 \mu\text{s} - \beta(400 \text{ m})/(2.998 \times 10^8 \text{ m/s})]$; (d) 0.750; (e) $0 < \beta < 0.750$; (f) $0.750 < \beta < 1$; (g) no 23. (a) 1.25; (b) $0.800 \mu\text{s}$
25. (a) 0.480; (b) negative; (c) big flash; (d) $4.39 \mu\text{s}$ 27. $0.81c$
29. (a) 0.35; (b) 0.62 31. $1.2 \mu\text{s}$ 33. (a) 1.25 y; (b) 1.60 y; (c) 4.00 y
35. 22.9 MHz 37. 0.13c 39. (a) 550 nm; (b) yellow
41. (a) 196.695; (b) 0.999 987 43. (a) 1.0 keV; (b) 1.1 MeV
45. 110 km 47. $1.01 \times 10^7 \text{ km}$ 49. (a) 0.222 cm; (b) 701 ps; (c) 7.40 ps 51. $2.83mc$ 53. $\gamma(2\pi m/lq|B|)$; (b) no; (c) 4.85 mm; (d) 15.9 mm; (e) 16.3 ps; (f) 0.334 ns 55. (a) 0.707; (b) 1.41; (c) 0.414 57. 18 smu/y 59. (a) 2.08 MeV; (b) -1.21 MeV
61. (d) 0.801 63. (a) $vt \sin \theta$; (b) $t[1 - (v/c) \cos \theta]$; (c) $3.24c$
67. (b) $+0.44c$ 69. (a) 1.93 m; (b) 6.00 m; (c) 13.6 ns; (d) 13.6 ns; (e) 0.379 m; (f) 30.5 m; (g) -101 ns ; (h) no; (i) 2; (k) no; (l) both
71. (a) $5.4 \times 10^4 \text{ km/h}$; (b) 6.3×10^{-10} 73. 189 MeV
75. $8.7 \times 10^{-3} \text{ ly}$ 77. 7 79. $2.46 \text{ MeV}/c$ 81. 0.27c
83. (a) 5.71 GeV; (b) 6.65 GeV; (c) $6.58 \text{ GeV}/c$; (d) 3.11 MeV; (e) 3.62 MeV; (f) $3.59 \text{ MeV}/c$ 85. 0.95c 87. (a) 256 kV; (b) 0.745c

Figures are noted by page numbers in *italics*, tables are indicated by t following the page number.

A

- a_g (gravitational acceleration), 334
 - variation with altitude, 334t
- absolute pressure, 363
- absolute zero, 476
- absorption, of heat by solids and liquids, 485–488
- absorption event, photons, 1058
 - in lasers, 1132
- absorption lines, 1100, 1101
- ac (alternating current), 826, 835
- acceleration, 263t. *See also* angular acceleration; forces
 - constant, 22, 22–24
 - free-fall, 25–26
 - graphical integration in motion analysis, 27, 27–28
 - negative, 20
 - and Newton's first law, 87–91
 - Newton's laws applications, 100–105
 - and Newton's second law, 91–94
 - one-dimensional motion, 18–26
 - principle of equivalence (with gravitation), 347
 - projectile motion, 64–70
 - relating linear to angular, 251, 251t
 - relative motion in one dimension, 73–74
 - relative motion in two dimensions, 74–75
 - rockets, 225, 224–226
 - rolling down ramp, 278–279, 279
 - sign of, 20
 - simple harmonic motion, 389, 389
 - system of particles, 206–209
 - two- and three-dimensional motion, 73–75
 - uniform circular motion, 70–72, 71, 124
 - as vector quantity, 38
 - yo-yo, 281
- acceleration amplitude, in simple harmonic motion, 389
- acceleration vectors, 38
- accelerators, 747–749, 1219, 1220, 1220
- acceptor atoms, 1153
- acre-foot, 9
- action at a distance, 580
- activity, of radioactive sample, 1175
- addition
 - of vectors by components, 44–46
 - of vectors geometrically, 39, 39–40, 40
- adiabat, 526, 526
- adiabatic expansion, 491–492, 492
 - ideal gas, 526, 526–528
- adiabatic processes
 - first law of thermodynamics for, 491–493, 492t
 - summarized, 528, 528t
- adiabatic wind, 534
- air
 - bulk modulus, 446
 - density, 360t
 - dielectric properties at 1 atm, 669, 669t
 - and drag force, 121–123
 - effect on projectile motion, 68–69, 68
 - electric breakdown, 593, 593, 602
 - electric breakdown field, 580t
 - index of refraction at STP, 906t
 - speed of sound in, 446–447, 447t
 - terminal speeds in, 122t
 - thermal conductivity, 494t
- air conditioners, 548
- airspeed, 84
- alpha decay, 1177–1179, 1178
- alpha particles, 572, 644, 1166–1168, 1166, 1177
 - binding energy per nucleon, 1172
- magic nucleon number, 1186
- radiation dosage, 1184
- in thermonuclear fusion, 1209
- alternating current (ac), 826, 835
- alternating current circuits, 835. *See also* transformers
 - capacitive load, 838–840, 839
 - forced oscillations, 836, 836
 - inductive load, 840–841, 841
 - LC oscillations, 826–829, 827
 - leading and lagging in, 841, 842t, 843
 - phase and amplitude relationships, 842t
 - power in, 847–849
 - resistive load, 836–838, 837
 - RLC circuit damped oscillations, 833, 833–834
 - series RLC circuits, 842–847, 843
- alternating current generator, 835
 - with capacitive load, 838–840, 839
 - with inductive load, 840, 840–842, 841
 - with resistive load, 836–838, 837
 - with transformer, 850, 850–853
- ammeters, 720, 720
- Ampère, André-Marie, 771
- Ampere-Maxwell law, 864, 864–866
 - Maxwell's equation form, 869t
- Ampere's law, 771–774
- ampere (unit), 565, 683, 770
- Amperian loop, 771, 771–774
- amplitude
 - alternating current, 842t
 - of emf in ac, 835
 - exponentially decaying in RLC circuits, 834
 - LC oscillations, 826
 - simple harmonic motion, 387–388, 388
 - waves, 415, 415–416, 416
- amplitude ratio, traveling electromagnetic waves, 892
- amusement park rides
 - roller coaster headache, 252
 - roller coasters, 20
 - Rotor, 249
- analyzer, 903
- Andromeda Galaxy, 330–331, 331
- angle of incidence, 906, 906
- angle of minimum deviation prism, 920
- rainbow, 922
- angle of reflection, 906, 906
- angle of refraction, 906, 906
- angles, 43
- angular acceleration, 243, 263t
 - relating to linear, 251, 251t
 - rolling wheel, 278, 278
 - rotation with constant, 248–249
- angular amplitude
 - simple pendulum, 396
- angular displacement, 242, 242
 - nonvector nature of, 247
- angular frequency
 - circulating charged particle, 745
 - damped harmonic oscillator, 400
 - of emf, 835
 - LC oscillations, 826, 831–832, 832
 - natural, 402
 - simple harmonic motion, 386–389, 388
 - sound waves, 449
 - waves, 416
- angular magnification
 - compound microscope, 944
 - refracting telescope, 945
 - simple magnifying lens, 943
- angular momentum, 284, 284–285, 289t
 - conservation of, 290–293, 291, 292
 - at equilibrium, 305
 - Newton's second law in angular form, 285
 - rigid body rotating about fixed axis, 288–290, 289
 - system of particles, 288
- angular motion, 242
- angular position, 242–243, 242, 263t
 - relating to linear, 250
- angular simple harmonic motion, 394, 394–395
- angular simple harmonic oscillator, 394, 394–395
- angular speed, 243, 244
 - relating to linear, 250–251
 - in rolling, 275–277, 276
- angular velocity, 243–246, 263t
 - vector nature of, 247, 246–247
- angular wave number, 415, 1072
 - sound waves, 449
- annihilation, 1222
 - electron-positron, 572, 572, 1222
 - proton-antiproton, 1224t, 1224
- annular cylinder
 - rotational inertia, 255t
- antenna, 891, 891
- antibaryons, 1237
- antiderivative, 24
- antihydrogen, 1108, 1222

- antimatter, 1222
 - energy released by 1 kg, 1195t
- antineutrino, 1180
- antinodes, 431, 431, 433
- antiparticles, 1222, 1224
 - formation in early universe, 1240
- antiprotons, 1222
 - antihydrogen, 1108, 1222
 - discovery of, 1248
 - proton-antiproton annihilation, 1224t, 1224
- antisolar point, 909, 909
- aphelion distance, 344
- apparent weight, 96
 - in fluids, 369–370
- Archimedes' principle, 367, 367–370
- area vector, 606
- astronomical Doppler effect, 1040–1042, 1041
- astronomical unit, 12
- atmosphere (atm), 361
- atmospheric pressure, 361t
- atomic bomb, 1172, 1201
 - rating, 1215
 - as trigger for fusion bomb, 1211
- atomic clocks, 5–6
 - time dilation tests, 1029
- atomic mass, 1171
 - selected nuclides, 1168t
- atomic mass units, 7, 1171
- atomic number, 572, 1117, 1168
- atoms, 1083. *See also* electrons; neutrons; protons
 - absorption and emission of light, 1113
 - angular momentum, 1113–1114, 1113, 1115–1117, 1116
 - Bohr model, 1096–1099, 1097
 - Einstein-de Haas experiment, 1114, 1114
 - electron spin, 1115
 - formation in early universe, 1240
 - magnetic resonance, 1121, 1120–1121
 - magnetism, 1113–1114, 1113
 - matter wave interference, 1068, 1068
 - orbital angular momentum, 1115–1117, 1116, 1117
 - properties of, 1112–1114, 1113
 - spin angular momentum, 1116–1117, 1117
 - Stern-Gerlach experiment, 1118, 1118–1120
- attractive forces, 331
- aurora, 561
- automobiles. *See also* race cars
 - magnet applications, 735
 - spark discharge from, 645, 645
 - tire pressure, 361t
- autotransformer, 859, 859
- average acceleration
 - one-dimensional motion, 18
 - two- and three-dimensional motion, 62–64
- average angular acceleration, 243
- average angular velocity, 243
- average power, 155, 185
 - engines, 547
 - traveling wave on stretched string, 423
- average speed
 - of gas molecules, 516–518
 - one-dimensional motion, 16
- average velocity
 - constant acceleration, 23
 - one-dimensional motion, 14–15
 - two- and three-dimensional motion, 60
- Avogadro's number, 507–508, 685
- axis of rotation, 242, 242
- B**
- Babinet's principle, 1016
- ball
 - projectile motion, 64–66, 65
- ballet dancing
 - grand jeté, 208, 208
 - tour jeté, 292, 292
- ballistic pendulum, 220, 220
- balloons, lifting capacity, 534
- Balmer series, 1097, 1099–1101, 1100
- band-gap pattern
 - crystalline solid, 1144
 - insulator, 1144
 - metal, 1145
 - semiconductor, 1151
- bands, energy bands in crystalline solids, 1144, 1144
- bar magnets
 - Earth as, 870, 870
 - magnetic dipole moment of small, 755, 755t
 - magnetic field, 862, 862–863
 - magnetic field lines, 738, 738–739
- barrier tunneling, 1074, 1074–1076
- baryonic matter, 1240
- baryon number, conservation
 - of, 1229
- baryons, 1221, 1228–1229, 1245
 - conservation of baryon number, 1229
 - and eightfold way, 1231t, 1231–1232
 - and quark model, 1232, 1237
- baseball
 - collision of ball with bat, 211, 211
 - fly ball, air resistance to, 68, 68
- base quantities, 1
- base standards, 1
- basilisk lizards, 233, 233
- basketball free throws, 58
- bats, navigation using ultrasonic waves, 464
- batteries. *See also* emf
 - connected to capacitors, 658, 658, 666
 - and current, 682–684, 683
 - as emf devices, 705, 706–707
 - in multiloop circuits, 714, 714–719
 - multiple batteries in multiloop circuit, 716–717, 717
 - potential difference across, 711–712, 712
 - and power in circuits, 695, 695–696
 - in RC circuits, 720, 720–724
 - recharging, 712
 - in RL circuits, 808–810
 - in single-loop circuits, 707–710
 - work and energy, 706, 706–707
- beam, 892
- beam expander, 957
- beam splitter, 980, 1067, 1067
- beats, 459, 459–460
- becquerel, 1175
- bends, 507
 - during flight, 379
- Bernoulli's equation, 374–377
- beta decay, 577, 1180–1182, 1181
 - and quark model, 1234
- beta-minus decay, 1180
- beta-plus decay, 1180
- bi-concave lens, 957
- bi-convex lens, 957
- bicycle wheels
 - rolling, 275–277, 276, 277
 - rolling with friction, 278–280, 279
- Big Bang, 1237, 1240–1242, 1241
- binding energy, *See* nuclear binding energy
- binding energy per nucleon, 1171, 1172, 1198
- Biot-Savart law, 765–766, 771, 779
- bivalent atom, 1145
- blackbody radiator, 496
- black holes, 330
 - event horizon, 337
 - gravitational lensing caused by, 348, 348
 - miniature, 351
 - and quasars, 1239
 - supermassive, 330
- blocks
 - acceleration of falling, 261
 - connected to massless, frictionless pulleys, 97, 98, 100, 100–101
 - floating, 370
 - forces on stationary, 117, 116–118
 - friction of sliding, 97, 97
 - hanging and sliding, 100, 100–101
 - Newton's laws applications, 92, 100–105
 - normal forces, 96–97, 97
 - power used in work on, 156, 156
 - stable static equilibrium, 305–306, 307, 309–313
 - third-law force pair, 98–99, 99
 - work done by external force with friction, 181, 181
- block-spring oscillator, 830
- block-spring systems
 - damped oscillating systems, 400, 400–401
 - and electrical-mechanical analogy, 829t, 829
 - kinetic energy, 149, 149–151, 151
 - oscillating systems, 390, 390–392
 - potential energy, 167–168, 167, 171
- blood pressure
 - normal systolic, 360t
- blue shift, 1041
- bob, of pendulum, 395
- body armor, 444, 444
- body diagonal, 56
- Bohr, Niels, 1089, 1185, 1199
- Bohr magneton, 872, 873, 1117
- Bohr model, of hydrogen, 1096–1199, 1097

- Bohr radius, 1098, 1106
boiling point, 486
 selected substances, 487t
Boltzmann, Ludwig, 1133
 entropy equation, 553
Boltzmann constant, 508
Bose, Satyendra Nath, 1220
Bose-Einstein condensate, 1221, 1221
bosons, 1220–1221, 1221
bottomness, 1231
bottom quark, 1233t, 1234
boundary condition, 1103
Bragg angle, 1012–1013
Bragg's law, 1012–1013
Brahe, Tycho, 342
branches, circuits, 714
breakdown potential, 670
breakeven, in magnetic confinement, 1212
Brewster angle, 912, 912
Brewster's law, 913
bright bands, *See* bright fringes
bright fringes
 double-slit interference, 964, 964, 966
 single-slit diffraction, 992–993, 993
British thermal unit (Btu), 485
Brookhaven accelerator, 1219
bubble chambers, 572, 572–573, 737, 737
 gamma ray track, 1069, 1069–1070
 proton-antiproton annihilation event, 1223, 1223
buildings
 natural angular frequency, 402
 submerged during earthquakes, 7–8
 swaying in wind, 393–394, 434
bulk modulus, 317, 446
bungee-cord jumping, 166, 166
buoyant force, 367–370, 368
- C**
c. *See* speed of light
calorie (cal), 484
Calorie (Cal) (nutritional), 484
cameras, 943, 955
cancer radiation therapy, 1165
 with ^{60}Co , 1190
capacitance, 656–658
calculating, 659–662
 LC oscillations, 826–829, 827
 parallel circuits, 662–663, 715t
 RC circuits, 720, 720–724
 RLC circuits, 833–834
 series circuits, 663–666, 715t, 842–844
capacitive reactance, 838
capacitive time constant
 RC circuits, 721, 721
capacitors, 656, 656–658, 658
 with ac generator, 838, 838–840
 charging, 657, 658
 charging in *RC* circuits, 720, 720–721
 with dielectrics, 669, 669–671
 discharging, 658
 discharging in *RC* circuits, 720, 720–721, 722–724
 displacement current, 866–869, 867
 electric field calculation, 659
 energy density, 668
 energy stored in electric field, 667–669
 Faraday's, 669, 669
 induced magnetic field, 863, 865–866
 LC oscillations, 826–829, 827
 in parallel, 662–663, 663, 715t
 phase and amplitude relationships for ac circuits, 842t
 potential difference calculation, 659–662
 RC circuits, 720, 720–724
 RLC circuits, 833–834, 833
 in series, 663–667, 664, 715t, 842–844, 843
 series *RLC* circuits, 842–844, 843
carbon cycle, in stars, 1217
carbon¹⁴ dating, 1183
carbon dioxide
 molar specific heat at constant volume, 520t
 RMS speed at room temperature, 512t
carbon disulfide
 index of refraction, 906t
Carnot cycle, 543, 543, 544
Carnot engines, 543–546, 543
 efficiency, 545–546, 549–550
Carnot refrigerators, 548, 549–550
cars, *See* automobiles; race cars
cascade, of decay processes, 1230
cat, terminal speed of falling, 122, 122–123
cathode ray tube, 740, 740–741
Celsius temperature scale, 479–481, 480
center of curvature
 spherical mirrors, 928, 928
 spherical refracting surfaces, 934, 934
center of gravity, 308–309, 308
center of mass
 and center of gravity, 308–309
 one-dimensional inelastic collisions, 218–220, 219, 220
 rolling wheel, 276, 276
 solid bodies, 203–206, 205
 system of particles, 202–203, 202, 206–209
center of momentum frame, 1056
center of oscillation, physical pendulum, 397
centigrade temperature scale, 479
central axis
 spherical mirrors, 929, 929
central configuration peak, 552
central diffraction maximum, 998, 998
central interference maximum, 966
central line, 1007
central maximum
 diffraction patterns, 990, 990
centripetal acceleration, 70–71
centripetal force, 124–129, 125
Cerenkov counters, 1248
CERN accelerator, 1219, 1236
 antihydrogen, 1222
 pion beam experiments, 1024
cesium
 work function, 1079
cesium₁₃₇, in fallout, 1194
chain-link conversion, of units, 3
chain reaction, 1201
change in entropy, *See* entropy changes
characteristic nuclear time, 1194
characteristic x-ray spectrum, 1128, 1128
charge, *See* electric charge
charge carrier density, 685
 silicon and copper compared, 696t, 696–697
charge carriers, 684, 1151
 doped semiconductors, 1152–1154
 drift speed, 686, 685–688
 semiconductors, 696
charged disk
 electric field due to, 591
 electric potential due to, 640, 640
charged isolated conductor
 with cavity, 612, 613
 electric potential, 644–645, 645
 in external electric field, 645, 645
 Gauss' law for, 612–615
 spark discharge from, 561–562
charged objects, 561–562, 580
charge number, 1117
charge quantum number, 1224
charging
 capacitors, 657–658
 capacitors in *RC* circuits, 720, 720–721
charm, 1231
charm quark, 1233t, 1234
Chernobyl reactor, 1194
chip (integrated circuits), 1159–1160
chocolate crumb explosion mystery, 626, 653, 680, 703
chromatic aberration, 946
chromatic dispersion, 907–908, 908
circuit elements, 658
circuits, 658, 658, 705, 715t. *See also* alternating current circuits
 capacitive load, 838–840, 839
 capacitors in parallel, 662–663, 663, 715t
 capacitors in series, 663, 663–666, 715t, 842–844, 843
 current direction, 714
 current in single-loop, 707, 707–709
 direct-current (DC), 705
 grounding, 712, 712
 inductive load, 840–842, 841
 inductors in series, 842–844, 843
 integrated, 1159–1160, 1159
 internal resistance, 709, 709
 multiloop, 708, 714, 714–719
 oscillating, 826
 parallel capacitors, 662–663, 663, 715t
 parallel inductors, 822
 parallel resistors, 714, 714–719, 715t
 potential difference between two points, 711, 711–714, 712
 power, potential, and emf, 712–713

- circuits (*cont.*)
 power in, 695–696
RC, 720, 720–724
 resistive load, 836–838, 837
 resistors in parallel, 714, 714–719, 715t
 resistors in series, 710, 710, 715t, 842–844, 843
RL, 807, 807–810, 809
RLC, 833, 833–834
 series capacitors, 663, 663–666, 715t, 842–844, 843
 series inductors, 842–844, 843
 series resistors, 710, 710, 715t, 842–844, 843
 single-loop, 707–710, 836
 work, energy, and emf, 706, 706–707
- circular aperture
 diffraction patterns, 1000, 1000–1003
- circular arc charge distributions, 590
- clocks
 event measurement array, 1025, 1025
 time dilation tests, 1029–1030, 1057
- closed circuit, 709, 709
- closed cycle processes
 first law of thermodynamics for, 492, 492t
- closed path, 168–169, 169
- closed-path test, for conservative force, 168
- closed shell, 1186
- closed subshell, 1126
- closed system, 207
 entropy, 542
 linear momentum conservation, 215
- COBE (Cosmic Background Explorer) satellite, 1242
- coefficient of kinetic friction, 119–121
- coefficient of linear expansion, 482
 selected materials, 482t
- coefficient of performance
 refrigerators, 548
- coefficient of static friction, 119–121
- coefficient of volume expansion, 482
- coherence, 969
- coherence length, 1131
- coherent light, 969, 1131
- coils, 753. *See also* inductors
 in ideal transformers, 850, 850–851
 magnetic field, 779, 779–780
 mutual induction, 813–815, 814
 self-induction, 806, 806–807
- cold-weld, 118, 118
- collective model, of nucleus, 1185
- collimated slit, 1118
- collimator, 1008, 1118
- collisions, 211–214
 elastic in one dimension, 221, 221–223
 glancing, 224, 224
 impulse of series of collisions, 213–214, 214
 impulse of single collision, 212, 211–212
 inelastic in one dimension, 218, 218–220, 219
 momentum and kinetic energy, 217–218
 two dimensions, 224, 224
- color force, 1236–1237
- color-neutral quarks, 1236–1237
- color-shifting inks, 958
- comets
 dust and ion tails, 917, 917
 stimulated emission from, 1140–1141, 1141
- commutator, 752, 752
- compass, 869
- completely inelastic collisions, 218, 218–220, 219
- complex conjugate, 1071
- component notation, vectors, 42
- components, of light, 907–908
- components, of vectors, 41, 41–44
 adding vectors by, 45, 44–46
- composite slab, conduction through, 494–495, 495
- compound microscope, 944, 944
- compound nucleus, 1185
- compressibility, 318, 361
- compressive stress, 316–317
- Compton scattering, 1062, 1062–1065, 1063
- Compton shift, 1062, 1062–1065
- Compton wavelength, 1064
- computers
 preventing sparks when working on, 732
- concave lenses, 957
- concave mirrors, 929, 929
 images, 931
- concrete
 coefficient of linear expansion, 482t
 elastic properties, 317, 317t
- condensing, 486
- conducting devices, 692
- conducting plates
 eddy currents, 800
 Gauss' law, 617, 617–618
- conduction, 494–495, 494, 495
- conduction band, 1151, 1151
- conduction electrons, 563, 682, 1145–1150
- conduction rate, 494
- conductivity, 690, 1146
- conductors, 563–564, 682. *See also* electric current
 drift speed in, 685–688
 Ohm's law, 692–695
- configurations, in statistical mechanics, 550–553
- confinement principle, 1084
- conservation of angular momentum, 290–294, 291, 292
- conservation of baryon number, 1229
- conservation of electric charge, 572–573
- conservation of energy, 140, 183–186, 184
- conservation of lepton number, 1228
- conservation of linear momentum, 215–217
- conservation of strangeness, 1231
- conservative forces, 168, 629
 path independence of, 168–170, 169
- constant acceleration
 one-dimensional motion, 22, 22–24
- constant angular acceleration
 rotation with, 248–249
- constant linear acceleration, 248
- constant-pressure molar specific heat, 521–522
- constant-pressure processes, 489, 489–490
 summarized, 528, 528t
 work done by ideal gases, 511
- constant-pressure specific heat, 486
- constant-temperature processes
 summarized, 528, 528t
 work done by ideal gases, 509, 509–510
- constant-volume gas thermometer, 478, 478–479
- constant-volume molar specific heat, 519–520
- constant-volume processes, 489, 489–490
 first law of thermodynamics for, 492t, 492
 summarized, 528, 528t
 work done by ideal gases, 510
- constant-volume specific heat, 486
- contact potential difference, 1155
- continuous charge distribution, 586
 electric potential due to, 639, 639–640, 640
- continuous x-ray spectrum, 1127, 1127–1128
- contracted length, 1032
- convection, 495
- converging lens, 936–937, 937
- conversion factors, 3
- convex lenses, 957
- convex mirrors, 929, 929
 images, 931
- Coordinated Universal Time (UTC), 5–6
- copper
 coefficient of linear expansion, 482t
 conduction electrons, 563
 electric properties compared to silicon, 696t, 696–697, 1143t, 1150–1151
 energy levels, 1143, 1143
 Fermi energy, 1145, 1162
 Fermi speed, 1145
 heats of transformation, 487t
 mean free time, 694
 resistivity, 689t, 691, 691, 1151
 rubbing rod with wool, 563
 temperature coefficient of resistivity, 1151
 unit cell, 1142, 1142
- copper wire
 aluminum competes with, 700
 as conductor, 563, 563, 683, 683
 drift speed in, 686
 electric field inside, 580t
 magnetic force on current carrying, 750, 750–751
- cord (unit of wood), 11
- core (Earth), 353, 353

- density, 335–336, 335, 336, 361t
 pressure, 361t
 core (Sun)
 density, 360t
 pressure, 361t
 speed distribution of photons in, 517
 corona discharge, 645
 correspondence principle, 1089, 1106
 cosine, 43, 43
 cosine-squared rule, for intensity of transmitted polarized light, 902
 Cosmic Background Explorer (COBE) satellite, 1242
 cosmic background radiation, 1239, 1241, 1242–1243, 1243
 cosmic ray protons, 577
 cosmic rays, 701
 Coulomb barrier, 1207
 coulomb per second, 683
 Coulomb's law, 565–572, 565 and Gauss' law, 612 spherical conductors, 566, 566–570
 coulomb (unit), 565
 Crab nebula pulsar, 268
 critical angle, for total internal reflection, 911
 crossed fields
 discovery of electron, 740–741
 Hall effect, 741–744, 741, 750
 crossed sheets, polarizers, 903, 903
 cross product, 49–51, 51
 crust (Earth), 353, 353
 density, 335, 360t
 crystal defects, 577
 crystalline solids
 electrical properties, 1142, 1142–1143
 energy levels, 1143, 1143
 crystal planes, 1011, 1011
 crystals
 matter waves incident after x ray and electron scattering, 1068, 1068, 1069
 polycrystalline solids, 880
 x-ray diffraction, 1011, 1011–1013
 curie (unit), 1175
 Curie constant, 878
 Curie's law, 878
 Curie temperature, 879
 currency, anti-counterfeiting measures, 958
 current, *See* electric current
 current amplitude
 series *RLC* circuits, 842–844, 843
 current-carrying wire
 magnetic field due to, 764–769, 765
 magnetic field inside long straight, 773, 773–774
 magnetic field outside long straight, 772, 772–773
 magnetic force, 750, 750–751
 magnetic force between parallel, 770, 770–771
 current density, 685–688, 685
 current law, Kirchhoff's, 714
 current-length element, 764, 764
 current loops, 683, 683
 electrons, 873–874, 873, 874
 Faraday's law of induction, 792–794
 Lenz's law for finding direction of current, 794, 794–797, 795
 as magnetic dipoles, 778–780, 779
 solenoids and toroids, 774–778, 775
 torque on, 752, 752–753
 curvature, of space, 347–348, 347, 1241
 cutoff frequency, photoelectric effect, 1060
 cutoff wavelength
 continuous x-ray spectrum, 1127
 photoelectric effect, 1060
 cycle
 engines, 543
 simple harmonic motion, 387
 thermodynamic, 489, 490, 492
 cyclotrons, 748, 748–749
 cylinders
 rotational inertia, 255t
 tracer study of flow around, 371
 calculating capacitance, 660, 660–661
 cylindrical symmetry, Gauss' law, 615, 615–616
- D**
 damped oscillations
RLC circuits, 833–834
 damped simple harmonic motion, 400, 400–401, 401
 damped simple harmonic oscillator, 400, 400
 damping constant, simple harmonic motion, 400
 damping force, simple harmonic motion, 400
 dark bands, *See* dark fringes
 dark energy, 1243
 dark fringes
 double-slit interference, 964, 964–965, 965, 967
 single-slit diffraction, 992, 992–994, 998
 dark matter, 1240
 daughter nuclei, 572, 1189
 dc (direct current), 705, 835
 de Broglie wavelength, 1067, 1085
 decay, *See* radioactive decay
 decay constant, 1174
 decay rate, 1174–1177
 deceleration, 20
 decibel, 454–456
 dees, 748
 de-excitation, of electrons to lower energy levels, 1086
 deformation, 315, 315
 degenerate energy levels, 1095
 degrees of freedom, ideal gas molecules, 523–524
 density
 defined, 7
 fluids, 359–361
 linear density of stretched string, 420
 selected engineering materials, 317t
 selected materials and objects, 360t
 uniform, for solid bodies, 203–204
 density gradient, 1155
 density of occupied states, 1149, 1149–1150
 density of states, 1146–1147, 1147
 depletion zone
 MOSFET, 1160
p-n junctions, 1155
 derived units, 2
 deuterium, 1182
 heavy water, 1217
 deuterium-tritium fuel pellets, 1212, 1212
 deuterons, 1190, 1211
 in cyclotron, 749
 deuteron-triton reaction, 1211
 diamagnetic material, 875
 diamagnetism, 875–876, 876
 diamond
 as insulator, 1144, 1151
 unit cell, 1142, 1142
 diamond lattice, 1142, 1142
 diatomic molecules, 520
 degrees of freedom, 523, 523–524, 524t
 molar specific heats at constant volume, 520t
 potential energy, 193
 dielectric constant, 669
 selected materials, 669t
 dielectrics
 atomic view, 671, 671
 with capacitors, 669–670
 and Gauss' law, 672–674, 672
 polarization of light by reflection, 912
 dielectric strength, 669–670
 selected materials, 669t
 differential equation, 830
 diffraction. *See also* interference
 circular aperture, 1000, 1000–1002
 double-slit, 1003, 1003–1005, 1004
 Fresnel bright spot, 991, 991
 intensity in double-slit, 1003, 1003–1004
 intensity in single-slit, 997–999, 998
 interference compared and contrasted, 1005
 by organized layers, 1011–1013, 1012
 single-slit, 963–964, 964, 992, 992–995, 993
 and wave theory of light, 990–991
 x rays, 1011–1013, 1012
 diffraction factor, 1004
 diffraction gratings, 1006, 1006–1008
 dispersion, 1009, 1009–1011
 resolving power, 1009–1011, 1010
 three-slit, 1019
 x rays, 1011
 diffraction patterns
 defined, 990–991
 double-slit, 1003, 1004
 single-slit, 1003, 1004
 diffusion current, *p-n* junctions, 1155
 dimensional analysis, 420
 dip meter, 870
 dip north pole, 870
 dipole antenna, 890–893, 891
 dipole axis, 584, 870
 dip-slip, 57
 direct current (dc), 835
 direct-current (dc) circuits, 705, 835

- direction
 of acceleration in one-dimensional motion, 19
 of acceleration in two- and three-dimensional motion, 62–63
 of displacement in one-dimensional motion, 13–14
 of vector components, 41
 of vectors, 39–40, 39, 40
 of velocity in one-dimensional motion, 16
 of velocity in two- and three-dimensional motion, 60
- discharging
 capacitors, 658
 capacitors in *RC* circuits, 720, 720–724
 a charged object, 563
- disintegration, 1169
- disintegration constant, 1174
- disintegration energy, 1177
- disks
 diffraction by circular aperture, 1000, 1000–1003
 rotational inertia, 251t
- dispersion
 chromatic, 907–908, 908
 by diffraction gratings, 1009, 1009–1011
- displacement. *See also* angular displacement; work
 damped harmonic oscillator, 400–401, 401
 one-dimensional motion, 14
 simple harmonic motion, 388, 388, 389
 traveling waves, 417
 two- and three-dimensional motion, 58–60, 59
 as vector quantity, 14, 38
 waves on vibrating string, 414–417, 415
- displacement amplitude
 forced oscillations, 402, 402
 sound waves, 449, 449
- displacement current, 866–868, 867
- displacement vector, 38, 38
- dissipated energy, in resistors, 695, 707
 ac current circuits, 847–848
- distortion parameter, 1200
- distribution of molecular speeds, 516–519, 517
- diverging lens, 936–937, 937
- donor atoms, 1152–1153
- doped semiconductors, 1152, 1152–1154
- doping, of semiconductors, 696, 1152–1154
- Doppler effect, 461–465, 1025
 detector moving, source stationary, 462, 462
 for light, 1040–1042, 1041
 source moving, detector stationary, 463, 463
- dose equivalent, radiation, 1184
- dot product, 48, 48
- double-slit diffraction, 1003, 1003–1005, 1004
- double-slit interference
 intensity, 969–972, 970
 from matter waves, 1067–1068, 1068
 single-photon, wide-angle version, 1067, 1067
 single-photon version, 1066
 Young's experiment, 964, 964–968
- doubly magic nuclides, 1186
- down quark, 578, 655, 1232–1234, 1233t
- drag coefficient, 122
- drag force, 121–123
 damped simple harmonic motion, 400
 mechanical energy not conserved in presence of, 175
 as nonconservative force, 168
- drain, FETs, 1159, 1159
- drift current, *p-n* junctions, 1155
- drift speed, 686–688
 Hall effect for determining, 741, 741–742
- driven oscillations, 402, 836, 836
- driving angular frequency, 835
- driving frequency, of emf, 835
- d* subshells, 1125, 1127
- E**
- E (exponent of 10), 2
- Earth, 330, 1243. *See also* core; crust; gravitational force; mantle
 angular momentum effects
 if polar ice caps melted, 304
 atmospheric electric field, 580t, 656
 average density, 360t
 density as function of distance from center, 335
 eccentricity of orbit, 343
 effective magnetic dipole moment, 1117
 ellipsoidal shape of, 335
 escape speed, 341t, 341–342
 gravitation near surface, 334–337, 336
 interior of, 353, 353
 Kepler's law of periods, 344t
 level of compensation, 380
 magnetic dipole moment, 754, 754t, 887
 magnetism, 869–871
 nonuniform distribution of mass, 335, 335–337
 rotation, 336, 336
 satellite orbits and energy, 345–346, 345
 variation in length of day over 4-year period, 6
- earthquakes
 building oscillations during, 386
 buildings submerged during, 7
 natural angular frequency of buildings, 402, 402
 S and P waves, 468
- Earth's magnetic field, 738, 870, 870–871
 magnetic latitude, 888
 polarity reversal, 870, 870–871
 at surface, 738t
- eccentricity, of orbits, 342–343, 343
 and orbital energy, 345
 planets of Solar System, 344t
- eddy currents, 800
- edge effect, 618
- effective cross-sectional area, 122
- effective magnetic dipole moment, 1117
- effective phase difference, optical interference, 962
- efficiency
 Carnot engines, 545–546
 real engines, 546, 549–550
 Stirling engines, 546
- eightfold way, 1231, 1231t, 1231–1232
- Einstein, Albert, 87, 893, 1022, 1022, 1025, 1043. *See also* relativity
 Bose-Einstein condensate, 1221, 1221
 and bosons, 1221
 and lasers, 1132
 view of gravitation, 347, 347–348
- work on photoelectric effect, 1059–1062
 work on photons, 1057–1059
- Einstein-de Haas experiment, 1114, 1114
- Einstein ring, 348, 348
- elastic bodies, 315
- elastic collisions
 defined, 217
 in one dimension with moving target, 222–223, 223
 in one dimension with stationary target, 221, 221–222
 in two dimensions, 224, 224
- elasticity, 305, 315–318, 316
 and wave speed on stretched string, 420
- elastic potential energy, 167
 determining, 171
 traveling wave on stretched string, 421–422, 421
- electrical breakdown, 5801t, 593, 593
- electrically isolated object, 562
- electrically neutral objects, 562
- electrical-mechanical analogy, 829t, 829
- electric charge, 561–562. *See also* circuits conductors and insulators, 563–564
 conservation of, 572–573
 and Coulomb's law, 565–572, 565
 induced, 564
 LC oscillations, 826, 830
 measures of, 586, 586t
 quantization, 571
- electric circuits. *See* circuits
- electric current, 682–685, 683
 calculating in single-loop circuits, 707, 707–709
 current density, 685–688, 685
- direction in circuits, 684, 684–685
 induced, 791, 797–800
 LC oscillations, 826, 831
 magnetic field due to, 764–769, 765
 time-varying in *RC* circuits, 720–724
- electric dipole, 582, 754
 in electric field, 594–596
 electric field due to, 584, 584–585
 electric potential due to, 637–638, 638
 induced, 638
 potential energy, 594–595
- electric dipole antenna, 891, 891–892

- electric dipole moment, 585, 585, 594
 - dielectrics, 671
 - permanent, 638
- electric displacement, 673
- electric field, 580–581, 736
 - calculating from potential, 641, 641–642
 - calculating potential from, 633, 633–634
- capacitors, 659
- crossed fields, 741–744, 741
- and current, 682
- due to charged disk, 591, 640, 640
- due to electric dipole, 584, 584–585
- due to line of charge, 586–590, 586
- due to point charge, 582, 582–583
- electric dipole in, 594–596
- energy stored in capacitor, 667–669
- equipotential surfaces, 631, 631–632, 632
- flux, 606, 606–608, 884
- Gauss' law for, 605, 608–620, 771, 862, 869t
- Hall effect, 741–744, 741, 750
- induced, 800–804, 801
- LC oscillations, 826
- point charge in, 592–593
- polarized light, 902, 902–904, 903
- system of point charges, 642–644, 643
- traveling electromagnetic waves, 890–893, 891, 893
- work done by, 628
- electric field lines, 581, 581–582
- electric fish, 718
- electric flux, 606, 606–608, 884
- electric force, 562, 735
- electric generator, 705
- electric motors, 752, 752–753, 869
- electric potential, 629–631
 - calculating field from, 641, 641–642
 - calculating from field, 633, 633–634
- charged isolated sphere, 645, 645
- due to charged disk, 640, 640
- due to continuous charge distribution, 639, 639–640, 640
- due to electric dipole, 637–638, 638
- due to group of point charges, 636, 636–637
- due to line of charge, 639, 639–640
- due to point charge, 635, 635
- and induced electric field, 803–804
- electric potential difference, 630
- electric potential energy, 628–629, 629
 - system of point charges, 642–644, 643
- electric quadrupole, 600, 600
- electric spark, 593, 593
 - airborne dust explosions set off by, 668–669
 - dangers of, 645, 645
 - and pit stop fuel dispenser fire, 723, 723–724
- electric wave component, of electromagnetic waves, 891–893, 893
- electromagnetic energy, 832
- electromagnetic force, 1221, 1235–1236
 - messenger particle, 1235–1237
- electromagnetic oscillations, 826–829, 827
 - damped in *RLC* circuits, 833–834
 - forced, 836, 836–838
- electromagnetic radiation, 890
- electromagnetic spectrum, 889–890, 890
- electromagnetic waves, 413, 889–890. *See also* reflection; refraction
 - energy transport, 897–899, 898
 - induced electric field, 894, 894–896
 - induced magnetic field, 895, 895–896
 - intensity variation with distance, 898, 898–899
 - polarization, 901, 901–904, 902, 903
 - polarization by reflection, 912, 912–913
 - Poynting vector, 897–899, 898
 - radiation pressure, 899–900
 - speed of travel, 413, 893
 - traveling, 890–896, 893, 894
- electromagnetism, 764, 869
- electromagnets, 735, 735
 - magnetic field near big, 738t
- electromotive force (emf), 705. *See also* emf
- electron capture, 1180
- electron diffraction, 1069
- electron microscope, 1080
- electron neutrinos, 1227, 1227t
- electron-positron annihilation, 572, 572, 1222
- electrons, 563, 1219
 - accelerator studies, 748
 - barrier tunneling, 1074, 1074–1076, 1075
 - Bohr model, 1096–1099, 1097
 - bubble chamber tracks, 572, 572, 738
 - charge, 571, 571t
 - Compton scattering, 1063, 1063–1065, 1063
 - conduction, 1145–1150
 - discovery by Thomson, 740, 740–741, 1165
 - as fermions, 1220
 - kinetic energy dependence on speed, 1023, 1023
 - as leptons, 1221, 1227, 1227t
 - loop model for orbits, 873–874, 873, 874
 - magnetic dipole moment, 754, 754t
 - and magnetism, 869–874
 - majority carrier in *n*-type semiconductors, 1153, 1153t
 - matter waves, 1067–1070, 1068, 1069, 1083–1084
 - minority carrier in *p*-type semiconductors, 1153, 1153t
 - orbital angular momentum, 872–873, 873
 - from proton-antiproton annihilation, 1224t
 - radiation dosage, 1184
 - spin, 1115, 1115t, 1220–1221, 1221
 - spin angular momentum, 871–872, 872
 - spin-flip, 883
 - in superconductors, 697–698
- electron spin, 1115, 1115t, 1220–1221, 1221
- electron traps, 1115
 - energies of, 1084–1088
 - finite well, 1091, 1091–1093
 - multiple electrons in rectangular, 1121–1124
 - nanocrystallites, 1093, 1093
 - one-dimensional, 1084–1095
 - quantum corrals, 1094, 1094
 - quantum dots, 1083, 1094, 1094
 - two- and three-dimensional, 1095, 1095–1096
- wave functions, 1088–1091, 1089
- electron-volt, 630
- electroplaques, 718, 718–719
- electrostatic constant, 565
- electrostatic equilibrium, 613
- electrostatic force, 562, 580
 - and Coulomb's law, 565–572, 565
- electric field due to point charge, 582, 582–583
- point charge in electric field, 592–593
- work done by, 628, 630–631
- electroweak force, 1236
- elementary charge, 570
- measuring, 592
- emf
 - alternating current, 835
 - defined, 705
 - and energy and work, 706, 706–707
 - induced, 792
 - potential and power in circuits, 711–714
 - self-induced, 806
- emf devices, 705. *See also* batteries
 - internal dissipation rate, 712
 - real and ideal, 706–706, 707
- emf rule, 709
- emission event, photons, 1058
 - lasers, 1132, 1132–1133
- emission lines, 1006, 1006, 1100–1101
- emissivity, 496
- endothermic reactions, 1227
- energy. *See also* gravitational potential energy; kinetic energy; mechanical energy; potential energy; thermal energy; work
 - conservation of, 140, 183–186, 184
 - dark energy, 1243
 - defined, 140
 - and relativity, 1043–1047
 - scalar nature of, 38
 - in simple harmonic motion, 392–394, 393
 - as state property, 537
 - stored in electric field, 667–669
 - stored in magnetic field, 811–812
 - transport by electromagnetic waves, 897–899, 898
 - traveling wave on stretched string, 421, 421–423
- energy bands, 1143–1144, 1144

- energy density
 capacitors, 668
 electric field, 668
 kinetic energy density of fluids, 374
 magnetic field, 812–813
 energy gap, 1143–1144, *1144*
 energy-level diagrams, 1085, *1085*
 one-electron, 1123, *1123*
 energy levels
 excitation and de-excitation, 1086
 full, empty, and partially occupied, 1122
 hydrogen, 1099–1100, *1099*, *1100*
 multiple electron traps, 1121–1124
 nuclear, 1172
 single electron traps, 1084–1088, *1085*
 energy method, of calculating current in single-loop circuits, 707
 engines
 Carnot, 543–546, *543*, 549–550
 efficiency, *543*, 545–546, *549*, 549–550, *550*
 ideal, 543–544
 perfect, *546*, *546*
 Stirling, *546*, *546*
 entoptic halos, 1014, 1016
 entropy
 engines, 543–546
 force due to, 542
 and irreversible processes, 536
 and probability, 552–553
 refrigerators, 548–549
 and second law of thermodynamics, 541–542
 as state function, 537, 539–540
 statistical mechanics view of, 550–553
 entropy changes, 536, 537–541
 Carnot engines, 545
 Stirling engines, 545
 entropy postulate, 536
 equation of continuity, 372–374, *372*
 equations of motion
 constant acceleration, 23, 23t
 constant linear and angular acceleration contrasted, 248t
 free-fall, 25
 equilibrium, 92. *See also* static equilibrium
 defined, 305–306
 electrostatic, 613
 examples of static, 309–313
 indeterminate structures, 314, *314*
 protons, 569
 requirements of, 306–307
 secular, 1191
 thermal, 477
 equilibrium charge, capacitors in *RC* circuits, 720
 equilibrium points, in potential energy curves, 176–177
 equilibrium position, simple pendulum, 395
 equilibrium separation, atoms in diatomic molecules, 193
 equipartition of energy, 523
 equipotential surfaces, *631*, 631–633
 equivalence, of gravity and acceleration, 347
 equivalent capacitance/capacitor
 parallel capacitances, 662–663, *663*, 715t
 series capacitances, 663–666, 715t
 equivalent inductance/inductor
 parallel inductances, 822
 series inductances, 822
 equivalent resistance/resistor
 parallel resistances, *714*, 714–719, 715t
 series resistances, 710, 715t
 escape speed, 341–342
 selected astronomical bodies, 341t
 event horizon, 337
 events, 1022
 Lorentz factor, *1028*, 1028, 1029, 1043
 Lorentz transformation, *1035*, 1035–1036
 measuring, 1024–1025, *1025*
 relativistic reversal, 1051
 relativity of length, *1031*, 1031–1034, 1038
 relativity of simultaneity, 1025–1026, *1026*, 1037
 relativity of time, *1027*, 1027–1031, 1037
 relativity of velocities, *1039*, 1039
 exchange coupling, 879, 887
 excitation, of electrons, 1086, *1086*
 excitation energy, 1110
 excited states, 1085, *1086*
 expansion, of universe, 1238, 1242–1243, *1243*
 explosions
 one-dimensional, *216*, 216–217
 two-dimensional, 216–217, *217*
 extended objects, 100, 927, *927*
 locating images by drawing rays, 939, *939*
 external electric field, 591
 Gaussian surfaces, *612*, 612–614
 isolated conductor in, 645, *645*
 external forces, 92
 collisions and internal energy transfers, 184–185
 system of particles, 206–209
 work done with friction, 180–182
 work done without friction, 180
 external magnetic field
 and diamagnetism, 875–876
 and ferromagnetism, 875, 879–880, *879*
 and paramagnetism, 875, 877–878
 external torque, 288
 and conservation of angular momentum, 290, 291
 eye, *See* human eye
 eyepiece
 compound microscope, *944*, 944
 refracting telescope, 945
F
 face-centered cubic, 1142, *1142*
 Fahrenheit temperature scale, 479–481, *480*
 farad, 657
 Faraday, Michael, 561, 581, 669, 791, 792, 805
 Faraday's law of induction, 792–794, 863
 Faraday's experiments, 792
 and Lenz's law, *794*, 794–797, 795
 Maxwell's equation form, 869t
 mutual induction, 814
 reformulation, 802
 self-induction, *806*, 806–807
 transformers, 850
 femtometer, 1170
 fermi (unit), 1170
 Fermi, Enrico, 1196, 1206, 1220
 Fermi-Dirac statistics, 1148
 Fermi energy, 1145, 1146, 1148
 calculating, 1150
 effect of doping on, 1163–1164, *1164*
 Fermilab accelerator, 1219, 1234
 Fermi level, 1145
 fermions, 1220–1221, *1221*
 quarks as, 1233t
 Fermi speed, 1145
 ferromagnetic materials, 875, 879–880, *879*, *880*
 magnetically hard and soft, 883
 ferromagnetism, 875, 879–880, *879*. *See also* iron
 FET (field-effect-transistor), 1159, *1159*
 field declination, 870
 field-effect-transistor (FET), 1159, *1159*
 field inclination, 870
 field of view
 refracting telescope, 945
 spherical mirrors, 928
 final state, 488, *489*, 520
 finite well electron traps, *1091*, 1091–1092
 fires
 pit stop fuel dispenser fire, 723, 723–724
 first law of thermodynamics, 491
 special cases of, 492t, 492–493
 first-order line, 1006
 first reflection point, *921*, 921
 fish, electric, 718–719
 fission fragments, 1196–1198
 fixed axis, 241
 angular momentum of rigid body rotation about, 288–290, 289
 flavors, of quarks, 1232, 1236–1237
 floaters, 991, 1017
 floating, 367, 368
 fluids, 121
 apparent weight in, 369–370
 Archimedes' principle, 367, 367–370
 Bernoulli's equation, 374–377
 defined, 359
 density, 359–360
 equation of continuity, 372, 372–373
 hydraulic lever, 366, 366–367
 motion of ideal, 371–372, *372*
 Pascal's principle, 366, 366–367
 pressure, 360–361

- pressure measurement,
365–367, 366
at rest, 362, 362–364
fluid streamlines, 371–372, 372
flux, 605–606, 606
of electric field, 606,
606–608, 884
of magnetic field, 862
focal length
compound microscope, 944,
944–945
refracting telescope, 945, 945
simple magnifying lens, 943,
943–944
spherical mirrors, 928,
928–930
thin lenses, 936–937, 937
focal plane, 967
focal point
compound microscope, 944,
944–945
refracting telescope, 945,
945–946
simple magnifying lens, 943,
943–944
spherical mirrors, 928,
928–929
thin lenses, 936–937, 937
two-lens system, 939–940,
940
focus, *See* focal point
force constant, 149
forced oscillations, 402, 836,
836
forced oscillators, 402, 402
forces, 289t. *See also* accelera-
tion; angular momen-
tum; collisions; drag
force; electrostatic
force; external forces;
friction; gravitational
force; internal forces;
linear momentum; mag-
netic force; spring force;
torque; variable force;
work
attractive, 331
buoyant, 367, 367–370
centripetal, 124–128, 125
conservative, 168–170, 169,
629
defined, 87
due to entropy, 542
equilibrium, 92
and motion, 13
net, 92
and Newton's first law, 87–91
Newton's laws applications,
100–105
and Newton's second law,
91–94
and Newton's third law, 98
nonconservative, 168
normal, 96–97, 97
path independence of con-
servative, 168–170, 169
principle of superposition
for, 88
resultant, 92
of rolling, 278–280, 279
tension, 97–98, 98
unit of, 88, 88–90
as vector quantities, 88
and weight, 95–96
forward-bias connection, junc-
tion rectifiers,
1156–1157, 1156, 1157
Franklin, Benjamin, 562, 570,
572, 701
Fraunhofer lines, 1141
free-body diagrams, 92–94, 93
Newton's laws applications,
100–105
free charge, 673
free-electron model, 1115,
1145
of Ohm's law, 693
free electrons, 682
free expansion
first law of thermodynamics
for, 492t, 492
ideal gases, 527–528, 537,
537–541, 538
free-fall acceleration (*g*),
25–26, 25
measuring with physical
pendulum, 397
free-fall flight, 25
free oscillations, 402, 836
free particle, 1071
and Heisenberg's uncer-
tainty principle, 1072,
1073
matter waves, 1084
free space, 890
freezing point, 486
frequency, 414–417
circulating charged particle,
744–745
and index of refraction, 961
photons, 1058
simple harmonic motion,
386–388, 388
sound waves, 449
wave on stretched string, 421
waves, 418, 418
Fresnel bright spot, 991, 991
friction, 97, 97, 117, 116–121.
See also kinetic fric-
tional force; static fric-
tional force
cold-weld, 118, 118, 119
as nonconservative force
(kinetic friction), 168
and rolling, 278–279, 279
work done by external force
with, 180–182, 180, 181
frictionless surface, 88, 97
fringing, 618
fuel charge, nuclear reactor,
1205
fuel rods, 1202, 1205
fulcrum, 321
full levels, 1122
full width at half maximum,
1016
fully charged capacitor, 658
fully constructive interference,
426, 427, 427t, 431
sound waves, 451
fully destructive interference,
426, 427, 427t, 431
sound waves, 451
fundamental mode, 434, 457
fur
rubbing plastic rod with, 562
fused quartz
coefficient of linear expan-
sion, 482t
index of refraction, 906t
index of refraction as func-
tion of wavelength, 907
resistivity, 689t
fusion, 1045, 1172, 1207–1208
controlled, 1211–1213
in Sun and stars, 1207,
1209–1210, 1209
fusion reaction, 1045
G
g units, of acceleration, 20
g, free-fall acceleration, 25–26
measuring with physical
pendulum, 397
G, gravitational constant, 331
galaxies, 330
Doppler shift, 1041
Doppler shift of NGC 7319,
1052, 1052
formation in early universe,
1241
gravitational lensing caused
by, 348, 348
preponderance of matter
over antimatter, 1222
recession, and expansion of
universe, 1238
superluminal jets, 1054
Galilean transformation equa-
tions, 1035
Galileo, 355
gamma-ray photons, 1222
in fusion, 1209
gamma rays, 573, 737, 890
bubble chamber track, 1069,
1069
radiation dosage, 1184
ultimate speed, 1023
gas constant, 508
gases, 507. *See also* ideal gases;
kinetic theory of gases
compressibility, 360
confined to cylinder with
movable piston,
488–490, 489
density of selected, 360t
as fluids, 359
specific heats of selected,
485t
speed of sound in, 447t
thermal conductivity of se-
lected, 494t
gas state, 486
gauge (wire), 700
gauge pressure, 363
gauss (unit), 738
Gauss, Carl Friedrich, 605
Gaussian form, of thin-lens
formula, 956
Gaussian surfaces
capacitors, 659–662
defined, 605, 605
electric field flux through,
606, 606–608
external electric field,
613–614, 614
and Gauss' law for magnetic
fields, 862
Gauss' law, 605
charged isolated conductor,
612–615
conducting plates, 617,
617–618
and Coulomb's law, 612
cylindrical symmetry, 615,
615–616
dielectrics, 672–674, 672
for electric fields, 605,
608–620, 862, 869t
isolated conductor with cav-
ity, 612, 613
for magnetic fields, 862–863,
863, 869t
Maxwell's equation forms,
869t
nonconducting sheet, 617,
617
planar symmetry, 617–618,
617
spherical symmetry,
619–620, 619
Geiger counter, 627, 627, 1165
general theory of relativity,
347–348, 1022, 1029

- geomagnetic pole, 738, 870
 geometrical optics, 904, 990
 diffraction as limit to, 963–964
 geosynchronous orbit, 354
 glass. *See also* Pyrex glass
 coefficient of linear expansion, 482t
 index of refraction, 906t
 as insulator, 563
 polarization of light by reflection, 912
 rubbing rod with silk, 562, 562, 563
 shattering by sound waves, 454
 Global Positioning System (GPS), 1, 1022
 g-LOC (*g*-induced loss of consciousness), 72
 in dogfights, 380
 gluons, 747, 1233, 1236
 gold, 1129
 alpha particle scattering, 1166
 impact with alpha particle, 644
 isotopes, 1169
 grand jeté, 208, 208
 grand unification theories (GUTs), 1237
 graphical integration, for one-dimensional motion, 27, 27–28
 grating spacing, 1006–1007
 grating spectroscope, 1008, 1008
 gravitation, 330
 and Big Bang, 1241
 Einstein's view of, 347–348, 348
 inside Earth, 337–338
 near Earth's surface, 334–337, 335
 Newton's law of, 330–332, 331, 342
 gravitational acceleration (a_g), 334
 variation with altitude, 334t
 gravitational constant (G), 331
 gravitational force, 94, 1221
 center of gravity, 308, 308–309
 electrostatic force contrasted, 572
 and Newton's law of gravitation, 330–332, 331
 pendulums, 395, 395
 and potential energy, 340–341
 and principle of superposition, 333–334
 work done by, 146, 146–148
 gravitational lensing, 348, 348
 gravitational potential energy, 166, 339–342, 339
 determining, 171
 and escape speed, 341–342
 and gravitational force, 340–341
 graviton, 348
 gray (unit), 1184
 grounding, 563
 ground speed, 84
 ground state, 1085, 1086, 1086
 wave function of hydrogen, 1102–1103, 1103
 zero-point energy, 1090
g subshells, 1125
 gyroscope precession, 294–295, 295
- H**
 hadrons, 1221, 1228–1230
 half-life, 1170, 1175, 1219
 and radioactive dating, 1183
 half-width of diffraction grating lines, 1006–1007, 1007
 resonance curves, 845–846, 846
 Hall effect, 741–744, 741, 750, 1142
 Hall potential difference, 742
 halo nuclides, 1170
 hanging blocks, 100, 100–101
 hard reflection, of traveling waves at boundary, 432
 harmonic motion, 387
 harmonic number, 434
 musical sounds, 456–459
 harmonic series, 434
 hearing threshold, 454t
 heat, 547–548. *See also* thermodynamics
 absorption by solids and liquids, 485–488
 defined, 484
 first law of thermodynamics, 491
 path-dependent quantity, 491
 signs for, 483–484
 and temperature, 483–484
 heat capacity, 485
 heat engines, 543–548
 heat of fusion, 487
 selected substances, 487t
 heat of vaporization, 486
 selected substances, 487t
 heat pumps, 548
 heats of transformation, 486–488
 selected substances, 487t
 heat transfer, 494–497
 Heisenberg's uncertainty principle, 1072, 1073, 1235–1236
 helical paths, charged particles, 745–746, 746
 helium-neon gas laser, 1133, 1133–1135
 Helmholtz coil, 787, 788, 790
 helium-burning, in fusion, 1210
 henry (unit), 805
 hertz, 386
 holes, 1128, 1151
 majority carrier in *p*-type semiconductors, 1153, 1153t
 minority carrier in *n*-type semiconductors, 1153, 1153t
 holograms, 1131
 home-base level, for spectral series, 1101
 Hooke, Robert, 149
 Hooke's law, 149, 176
 hoop
 rotational inertia, 255t
 horizontal motion, in projectile motion, 66–67, 67
 horizontal range, in projectile motion, 65, 66
 horsepower (hp), 155
h subshells, 1125
 Hubble constant, 1238
 Hubble's law, 1238
 human body
 as conductor, 563
 physiological emf devices, 706
 human eye, 943, 955, 955
 floaters, 991, 1017
 image production, 926
 sensitivity to different wavelengths, 890, 890
 Huygens' principle, 959, 959–960
 Huygens' wavelets, 959, 992, 995
 hydraulic compression, 317
 hydraulic engineering, 359
 hydraulic jack, 367
 hydraulic lever, 366, 366–367
 hydraulic stress, 316, 316–318
 hydrogen
 Bohr model, 1096–1099, 1097
 electric field within, 580t
 emission lines, 1008, 1008
 energy levels and spectra, 1099–1101
 formation in early universe, 1241
 in fusion, 1045, 1207–1213
 heats of transformation, 487t
 quantum numbers, 1101t, 1101–1103
 RMS speed at room temperature, 512t
 and Schrödinger's equation, 1099–1106, 1115
 speed of sound in, 447t
 thermal conductivity, 494t
 wave function of ground state, 1102–1103, 1103
 hydrogen bomb (thermonuclear bomb), 1211, 1217
 hydrostatic pressures, 362–364
 hysteresis
 ferromagnetic materials, 880–881
 hysteresis loop, 880, 880
- I**
 ideal emf devices, 706
 ideal engines, 543–544
 ideal fluids
 motion, 371–372, 372
 ideal gases, 508–511
 adiabatic expansion, 526–528
 average speed of molecules, 516–519
 free expansion, 537, 537–541, 538
 internal energy, 519–522
 molar specific heat and degrees of freedom, 523–524
 molar specific heat at constant pressure, 521–522
 molar specific heat at constant volume, 519–520
 molar specific heats, 519–522
 most probable speed of molecules, 517
 RMS speed, 511, 511–513, 512t, 516
 translational kinetic energy, 513–514
 work done at constant pressure, 510
 work done at constant temperature, 509, 509–510
 work done at constant volume, 510
 ideal gas law, 508
 ideal gas temperature, 479
 ideal inductor, 807
 ideal refrigerators, 548

- ideal solenoid, 776–777
- ideal spring, 149
- ideal toroids, 778
- ideal transformers, 850, 850–852
- ignition, in magnetic confinement, 1212
- image distances, 926
- images
 - extended objects, 927, 927, 939, 939
 - locating by drawing rays, 932, 932–933, 939, 939
 - from plane mirrors, 926, 926–928
 - from spherical mirrors, 928, 928, 930, 930–934, 946, 946
 - from spherical refracting surfaces, 934, 934–935, 946–947, 946
 - from thin lenses, 938, 938–942, 939, 947–948, 947
 - types of, 924–925
- impedance, 843
 - matching in transformers, 852
- impulse, 212
 - series of collisions, 213, 213
 - single collision, 211, 211–212
- incident ray, 904–906, 905
- incoherent light, 969
- incompressible flow, 371
- indefinite integral, 24
- independent particle model, of nucleus, 1185–1186
- indeterminate structures
 - equilibrium, 314, 314
- index of refraction
 - and chromatic dispersion, 907–908
 - common materials, 906t
 - defined, 906, 960
 - and wavelength, 960–962
 - induced charge, 564
 - induced current, 791
 - induced electric dipole moment, 638
- induced electric fields, 800–804, 801
 - from traveling electromagnetic waves, 894, 894–896
- induced emf, 792
- induced magnetic fields, 863–866, 864
 - displacement current, 866–869, 867
 - finding, 867–868, 868
 - from traveling electromagnetic waves, 895, 895–896
- inductance, 805–806
 - LC* oscillations, 826–829
 - RL* circuits, 807–810
 - RLC* circuits, 833–834
 - solenoids, 805–806, 806
- induction, 792. *See also* Faraday's law of induction
 - energy transfer, 797–800, 798
 - mutual induction, 813–815, 814
 - self-induction, 806, 806–807, 813
- inductive reactance, 841
- inductive time constant, 808–809
- inductors, 805, 805–807
 - with ac generator, 840, 840–841, 841
 - LC* oscillations, 826–829, 827
 - in parallel, 822
 - phase and amplitude relationships for ac circuits, 842t
 - RL* circuits, 807–810
 - RLC* circuits, 833, 833–834
 - in series, 822, 844–853
 - series *RLC* circuits, 842–844, 843
- inelastic collisions
 - defined, 218
 - in one dimension, 218, 218–220, 219
 - in two dimensions, 224
- inertial confinement, 1212
- inertial reference frames, 89–90, 1022
- inexact differentials, 491
- infinitely deep potential energy well, 1085, 1085
- infinite potential well, 1085
- inflation, of early universe, 1241
- infrared radiation, 899
- initial conditions, 391
- initial state, 488, 489, 519
- in phase
 - ac circuits, 842t
 - resistive load, 838
 - sound waves, 451
 - thin-film interference, 973–974
 - traveling electromagnetic waves, 890–891
 - waves, 425, 428
- instantaneous acceleration
 - one-dimensional motion, 18
 - two- and three-dimensional motion, 62–63
- instantaneous angular acceleration, 243
- instantaneous angular velocity, 243
- instantaneous power, 155, 185
- instantaneous velocity
 - one-dimensional motion, 17–18
 - two- and three-dimensional motion, 60
- insulators, 563–564, 697
 - electrical properties, 1144, 1144
 - resistivities of selected, 689t
 - unit cell, 1142
- integrated circuits, 1159–1160
- intensity
 - diffraction gratings, 1006
 - double-slit diffraction, 1003, 1003–1005
 - double-slit interference, 969–972, 970
 - electromagnetic waves, 898, 898–899
 - single-slit diffraction, 995–999, 996, 998
 - of transmitted polarized light, 902–904, 903
- intensity, of sound waves, 452–456, 453
- interference, 4257, 425–428, 426 *See also* diffraction and coherence, 969
 - combining more than two waves, 972
 - diffraction compared and contrasted, 1003–1004
 - double-slit from from matter waves, 1067–1069, 1068
 - double-slit from single photons, 1065, 1065–1067
 - intensity in double-slit, 969–972, 970
 - Michelson's interferometer, 980, 980
 - Newton's rings, 987, 987
 - and rainbows, 962, 962
 - sound waves, 451, 451–452
 - thin films, 973, 973–979
 - Young's double-slit experiment, 964, 964–968, 965
- interference factor, 1004
- interference fringes, 965, 965
- interference pattern, 964, 964, 967
- interfering waves, 425
- interferometer, 980
- intermediate interference, 426, 427, 427t
- sound waves, 451
- internal energy, 476
 - and conservation of total energy, 183
 - and external forces, 184–185
 - and first law of thermodynamics, 491
 - of ideal gas by kinetic theory, 519–522
- internal forces, 92
 - system of particles, 206–9
- internal resistance
 - ammeters, 720
 - circuits, 709, 709
 - emf devices, 712–713
- internal torque, 288
- International Bureau of Weights and Standards, 3, 6
- International System of Units, 2
- interplanar spacing, 1012
- intrinsic angular momentum
 - electrons, 871
 - protons and neutrons, 872
- inverse cosine, 43, 43
- inverse sine, 43, 43
- inverse tangent, 43, 43
- inverse trig functions, 43, 43
- inverted images, 930, 930
- ionization energy, 1113, 1113
 - plot of elements, 1113
- ionized atoms, 1101
- iron, 1126–1127. *See also* steel
 - Curie temperature, 879
 - ferromagnetic material, 875, 879–880
 - quantum corral, 1094, 1094
 - radius of nucleus, 571
 - resistivity, 689t
- iron filings
 - bar magnet's effect on, 862, 862
 - current-carrying wire's effect on, 765
- irreversible processes, 536. *See also* entropy and second law of thermodynamics, 541–542
- irrotational flow, 371, 374
- island of stability, 1170
- isobaric processes summarized, 528, 528t
- isobars, 1170
- isochoric processes summarized, 528, 528t
- isolated spherical capacitors, 669
 - calculating capacitance, 661–662

- isolated system, 173
 conservation of total energy, 183–184
 linear momentum conservation, 215
- isotherm, 509, 509
- isothermal compression, 509
 Carnot engine, 543, 543
- isothermal expansion, 509
 Carnot engine, 543, 543
 entropy change, 537–538, 538
- isothermal processes
 summarized, 528, 528t
- isotopes, 1169
- isotopic abundance, 1168t
- isotropic materials, 690
- isotropic point source, 898
- isotropic sound source, 454
- J**
- joint, in rock layers, 131
- Josephson junction, 1075
- joule (J), 141, 485
- junction diodes, 697
- junction lasers, 1158, 1158
- junction plane, 1154, 1155
- junction rectifiers, 1156, 1156–1157, 1164
- junction rule, Kirchoff's, 714
- junctions, circuits, 714
- K**
- kaons, 1030, 1219
 and eightfold way, 1231t
 and strangeness, 1231
- kelvins, 476, 478, 479, 482
- Kelvin temperature scale, 476, 476, 480
- Kepler, Johannes, 342
- Kepler's first law (law of orbits), 342–343, 343
- Kepler's second law (law of areas), 343, 343
- Kepler's third law (law of periods), 344, 344
 for planets in Solar System, 344t
- kilogram, 6
- kilowatt-hour, 155
- kinematics, 13
- kinetic energy, 263t. *See also* collisions; rotational kinetic energy; translational kinetic energy; work-kinetic energy theorem
- collisions, 217–218
 and conservation of mechanical energy, 173–175
 and conservation of total energy, 183–186
 defined, 141
 and momentum, 1046
 and relativity, 1045, 1045–1047
 of rolling, 277, 277–280
 of rotation, 253–254, 254
 satellites in orbit, 345–346, 346
 simple harmonic motion, 393, 393
 traveling wave on stretched string, 421, 421, 422
 and work, 142–146, 143
 yo-yo, 281
- kinetic energy density, of fluids, 374
- kinetic energy function, 176
- kinetic frictional force, 118, 118
 as nonconservative force, 168
 rolling wheel, 278
- kinetic theory of gases. *See also* gases; ideal gases
 average speed of molecules, 516–518
 and Avogadro's number, 507–508
 distribution of molecular speeds, 516–519, 517
 mean free path, 514, 514–515
 molar specific heat and degrees of freedom, 523–524
 most probable speed of molecules, 517
 and quantum theory, 523, 525
 RMS speed, 511, 511–513, 512t, 516
 translational kinetic energy, 513–514
- Kirchoff's current law, 714
- Kirchoff's junction rule, 714
- Kirchoff's loop rule, 708
- Kirchoff's voltage law, 708
- K* shell, 1128, 1129, 1136
- L**
- lagging, in ac circuits, 841, 842t, 843
- lagging waves, 428
- lambda particle and eightfold way, 1231t
- lambda-zero particle, 1230
- laminar flow, 371
- lasers, 1131–1132
 coherence, 969
 helium-neon gas, 1133, 1133–1135
 junction lasers, 1158, 1158
 nuclear-pumped x-ray, 1017
 operation, 1132, 1132–1135
 for plasma compression, 916
 radiation pressure, 899–900
 surgery applications, 1131, 1131
- laser thickness monitoring, 1015, 1015
- lasing, 1134, 1140
- lateral magnification
 compound microscope, 944
 spherical mirrors, 931
 two-lens system, 939
- lateral manipulation, using STM, 1075
- lattice, 315, 315, 1142, 1142
- law of areas (Kepler's second law), 343, 343
- law of Biot and Savart, 765–766, 771, 779
- law of conservation of angular momentum, 290, 290–294, 291, 292
- law of conservation of electric charge, 572–573
- law of conservation of energy, 183–186, 184
- law of conservation of linear momentum, 215
- law of orbits (Kepler's first law), 342–343, 343
- law of periods (Kepler's third law), 344, 344
 for planets in Solar System, 344t
- law of reflection, 906
- law of refraction, 906, 959, 959–962
- Lawson's criteria, 1211
- LC* oscillations
 and electrical-mechanical analogy, 830t, 830
 qualitative aspects, 826–829, 827
 quantitative aspects, 830–833
- LC* oscillators, 830t, 830–833
 radio wave creation, 890–893, 891
- lead
 coefficient of linear expansion, 482t
 heats of transformation, 487t
 specific heats, 485t
 thermal conductivity, 494t
- leading, in ac circuits, 841, 842t, 843
- leading waves, 428
- Leidenfrost effect, 504
- length
 relativity of, 1031, 1031–1034, 1037
 length contraction, 1032–1034
 consequences of Lorentz transformation equations, 1037–1038
 length units, 3–4
 lens, 936. *See also* thin lenses
 diffraction by, 1000
 lens maker's equation, 936
 Lenz's law, 794, 794–797, 795
 self-induction, 806
- lepton number, 1228
 conservation of, 1228
- leptons, 1221, 1227t, 1227–1228
 conservation of lepton number, 1228
 formation in early universe, 1241
- lifetime, subatomic particles, 1029
- lifting capacity, balloons, 534
- light, 413, 894. *See also* diffraction; interference; photons; reflection; refraction
- Doppler effect, 461
 in early universe, 1241
 Huygens' principle, 959, 959–960
 law of reflection, 906
 law of refraction, 906, 959, 959–962
 as probability wave, 1065–1067
 travel through media of different indices of refraction, 961, 961
 as wave, 959, 959–963
 wave theory of, and diffraction, 990–991
- light-emitting diodes (LEDs), 1157, 1157–1158, 1158
- light-gathering power refracting telescope, 945
- lightning, 561, 656
 role in creation of lode-stones, 881
 side flash, 728, 728
 upward streamers, 616, 616
- light quantum, 1057–1058
- light wave, 893
- light-year, 12
- line (spectral), 1101
- linear acceleration, *See* acceleration
- linear charge density, 586, 586t
- linear density, of stretched string, 420, 421
- linear expansion, 482, 482

- linear momentum, 210, 289t
 - completely inelastic collisions in one dimension, 218–219
 - conservation of, 215–217
 - elastic collisions in one dimension with moving target, 222–223
 - elastic collisions in one dimension with stationary target, 221–222
 - elastic collisions in two dimensions, 224
 - at equilibrium, 305
 - and impulse of series of collisions, 213
 - and impulse of single collision, 211–212
 - inelastic collisions in one dimension, 218, 218–220, 219
 - inelastic collisions in two dimensions, 224
 - photons, 1062, 1062–1065, 1063
 - system of particles, 211
- linear momentum-impulse theorem, 212
- linear motion, 241
- linear oscillator, 390–392
- linear simple harmonic oscillators, 390, 390–392
- line integral, 633
- line of action, of torque, 259, 259
- line of charge
 - electric field due to, 586–590, 586
 - electric potential due to, 639, 639–640
- line of symmetry, center of
 - mass of solid bodies with, 203
- lines, diffraction gratings, 1006–1007
- lines of force, 581
- lines shapes, diffraction gratings, 1010
- liquefaction, of ground during earthquakes, 7
- liquids
 - compressibility, 317, 360
 - density of selected, 360t
 - as fluids, 359
 - heat absorption, 485–488
 - speed of sound in, 447t
 - thermal expansion, 481–483
- liquid state, 486
- Local Group, 330
- Local Supercluster, 330
- lodestones, 561, 869, 881
- longitudinal motion, 414
- longitudinal waves, 413–414, 414
- long jump
 - conservation of angular momentum, 291, 291
- long wave radiation, 890
- loop model, for electron orbits, 873–874, 873, 874
- loop rule, 708, 714
 - RL* circuits, 808, 808
- Lorentz factor, 1028, 1028, 1029, 1043
- Lorentz transformation, 1035, 1035–1036
- Lorentz transformation equations, 1035–1036
 - consequences of, 1037–1038
 - pairs of events, 1036t
- Loschmidt number, 535
- L* shell, 1128–1129, 1128
- Lyman series, 1100, 1101, 1104
- M**
- Mach cone, 465, 465
- Mach cone angle, 465, 465
- Mach number, 465
- magic electron numbers, 1185–1186
- magic nucleon numbers, 1185–1186
- magnetically hard material, 883
- magnetically soft material, 883
- magnetic confinement, 1212
- magnetic dipole moment, 754, 754–755, 1114, 1114
 - diamagnetic materials, 875–876
 - effective, 1117
 - ferromagnetic materials, 875, 879
 - orbital, 873
 - paramagnetic materials, 875, 877–878
 - spin, 871–872, 872
- magnetic dipoles, 738, 753–755, 754, 862, 862–863
- magnetic domains, 879–880, 880
 - and hysteresis, 880
- magnetic energy, 811–812
- magnetic energy density, 812–813
- magnetic field, 736. *See also*
 - Earth's magnetic field
 - Ampere's law, 771, 771–774
 - circulating charged particle, 744–747, 745, 746
 - crossed fields, 740, 741, 744
 - cyclotrons, 748, 748
 - defined, 735, 737, 739
 - displacement current, 866–868, 867
 - due to current in circular arc of wire, 767, 767–768
 - due to current in long straight wire, 765–767, 766
 - Earth, 870, 870–871
 - energy density of, 812–813
 - energy stored in, 811–812
 - and Faraday's law of induction, 792–794, 793
 - Gauss' law for, 862–863, 869t
 - Hall effect, 741, 741–744, 750
 - induced, 863–865, 864
 - inside long straight current carrying wire, 773, 773–774
 - LC* oscillations, 826
 - and Lenz' law, 794, 794–797, 795
 - outside long straight current carrying wire, 772, 772–773
 - parallel currents, 770, 770–771
 - selected objects and situations, 738t
 - solenoids, 776–777, 775
 - synchrotrons, 749
 - toroids, 777, 777–778
 - torque on current loops, 752, 752–753
 - traveling electromagnetic waves, 890–893, 891, 893
- magnetic field lines, 738, 738–739, 739
 - current-carrying wire, 766–767
- magnetic flux, 862
 - and Faraday's law, 792–793
- magnetic flux linkage, 805
- magnetic force, 561, 736
 - circulating charged particle, 744–747, 745, 746
 - current-carrying wire, 750, 750–751, 770, 770–771
 - particle in magnetic field, 736–738, 737
- magnetic latitude, 887
- magnetic levitation, 697, 697
- magnetic materials, 861, 862, 875
 - diamagnetism, 875, 875–876
 - ferromagnetism, 875, 879–880, 879
- magnetically hard and soft, 883
- paramagnetism, 875, 877, 877–878
- magnetic monopole, 736, 862
- magnetic potential energy, 753–754, 811
- magnetic resonance, 1120–1121, 1121
- magnetic resonance imaging, *See* MRI
- magnetic wave component, of electromagnetic waves, 892–893, 893
- magnetism. *See also* Earth's magnetic field
 - Earth, 870, 870–871
 - electrons, 871–874
 - protons and neutrons, 872
- magnetization
 - ferromagnetic materials, 879, 879–880
 - paramagnetic materials, 877, 877–878
- magnetization curves
 - ferromagnetic materials, 879, 879–880
 - hysteresis, 880, 880
 - paramagnetic materials, 868, 877
- magnetometers, 870
- magnets, 561, 735, 735–739, 738, 869–871
 - applications, 735
- magnetic dipoles, 862, 862–863
- magnitude
 - of acceleration in one-dimensional motion, 19
 - of acceleration in two- and three-dimensional motion, 62
 - of displacement in one-dimensional motion, 14
 - of free-fall acceleration, 25
 - of vectors, 39, 39
 - of velocity in one-dimensional motion, 15
 - of velocity in two- and three-dimensional motion, 62–63
- magnitude-angle notation, vectors, 42
- magnitude ratio, traveling electromagnetic waves, 893
- majority carriers, 1153
 - motions of, 1154–1156, 1155
- mantle (Earth), 353, 353
- density, 335
- mass, 263t. *See also* center of mass
 - defined, 90–91
 - scalar nature of, 38, 91
 - of selected objects, 7t

- mass (*cont.*)
 units, 6–7
 and wave speed on stretched string, 420
 weight contrasted, 96
 mass dampers, 394
 mass energy, 1043–1044
 energy equivalents of particles and objects, 1043t
 mass excess, 1171
 mass flow rate, 373
 massless, frictionless pulleys, 98, 98
 Newton's laws applications, 100, 100–101
 massless cord, 98, 98
 massless spring, 149
 mass number, 572, 1168
 selected nuclides, 1168t
 mass spectrometer, 747, 747
 matter
 antimatter, 1222
 baryonic matter, 1240
 dark matter, 1240
 energy released by 1 kg, 1195t
 nuclear, 1174
 particle nature of, 1069–1070, 1069
 wave nature of, 1067–1069
 matter waves, 413
 electrons, 1067–1070, 1068, 1069, 1070, 1083, 1084
 maxima
 diffraction patterns, 990, 990
 double-slit interference, 964, 964, 967, 970
 single-slit diffraction, 992–995
 thin-film interference, 975
 Maxwell, James Clerk, 516, 523, 561, 771, 863, 890, 899, 958, 1236
 Maxwellian electromagnetism, 1218
 Maxwell's equations, 861, 869t, 1071
 Maxwell's law of induction, 863
 Maxwell's rainbow, 889–890
 Maxwell's speed distribution law 516–519, 517
 mean free time, 694
 mean life, radioactive decay, 1175, 1219
 measurement, 1
 mechanical energy
 conservation of, 173–175
 and conservation of total energy, 183
 damped harmonic oscillator, 400
 satellites in orbit, 345, 345
 simple harmonic motion, 392–393, 393
 mechanical waves, 413
 medium, 893
 melting point, 486
 selected substances, 487t
 meniscus concave lens, 957
 meniscus convex lens, 957
 mercury barometer, 361, 365, 365
 mercury thermometer, 482
 mesons, 1221, 1228–1229
 and eightfold way, 1231t, 1231–1232
 and quark model, 1232–1234, 1237
 underlying structure suggested, 1232
 messenger particles, 1235–1237
 metallic conductors, 682, 696
 metal-oxide-semiconductor-field effect-transistor (MOSFET), 1159, 1159
 metals
 coefficients of linear expansion, 482t
 density of occupied states, 1149, 1149–1150
 density of states, 1146–1148, 1147
 elastic properties of selected, 317t
 electrical properties, 1145–1150
 lattice, 315, 315
 occupancy probability, 1147–1148, 1148
 resistivities of selected, 689t
 speed of sound in, 447t
 thermal conductivity of selected, 494t
 unit cell, 1142
 metastable states, 1132
 meter (m), 1, 3–4
 metric system, 2
 mho (unit), 690
 Michelson's interferometer, 980, 980
 microfarad, 657
 microscopes, 943
 compound, 944, 944
 electron microscopes, 1080
 microscopic clocks, time dilation tests, 1029
 microstates, in statistical mechanics, 550–552
 microwave cooking, 595
 microwaves, 413, 889
 Doppler effect, 461
 Mid-Atlantic Ridge, magnetism, 870, 870
 Milky Way galaxy, 330
 Millikan oil-drop experiment, 592, 592
 millimeter of mercury (mm Hg), 361
 millimeter-wave radar, 1017
 miniature black holes, 351
 minima
 circular aperture diffraction, 1000, 1000
 diffraction patterns, 990, 990–991, 991
 double-slit interference, 964, 964, 966, 970
 single-slit diffraction, 992–995, 995–997, 996
 thin-film interference, 976
 minority carriers, 1153
 motions of, 1154–1155
 mirage, 925, 925
 mirrors, 926
 plane, 926, 926–928
 spherical, 928–934, 928, 929, 946, 946–947
 mobile radio, 889
 moderators, for nuclear reactors, 1202
 modulus of elasticity, 315
 Mohole, 353
 molar mass, 508
 molar specific heat, 485
 at constant pressure, 521–522
 at constant volume, 519–520
 and degrees of freedom, 523, 523–524
 selected materials, 485t
 molecular mass, 508
 molecular speeds, Maxwell's distribution of, 516–519, 517
 molecules, 1112
 matter wave interference, 1069
 moment arm, 259, 259
 moment of inertia, 253
 momentum, 210. *See also* angular momentum; linear momentum
 center of momentum frame, 1056
 and kinetic energy, 1046
 and relativity, 1042–1043
 monatomic molecules, 519
 degrees of freedom, 523, 523–524, 524t
 molar specific heats at constant volume, 520t
 monochromatic light, 907
 lasers, 1131
 monovalent atom, 1145
 Moon, 330
 corona around, 1018, 1018
 potential effect on humans, 351
 radioactive dating of rocks, 1183
 more capacitive than inductive circuit, 844
 more inductive than capacitive circuit, 844
 Moseley plot, 1129, 1129–1130
 MOSFET (metal-oxide-semiconductor-field-effect-transistor), 1159, 1159
 most probable configuration, 552
 most probable speed in fusion, 1207
 of gas molecules, 517
 motion, 13. *See also* acceleration; collisions; displacement; forces; one-dimensional motion; position; uniform circular motion; velocity
 graphical integration, 27, 27–28
 projectile, 64–70, 65
 relative in one dimension, 73, 73–74
 relative in two dimensions, 74–75, 75
 two- and three-dimensional, 58–64
 MRI (magnetic resonance imaging)
 effect on magnetic tattoo inks, 861, 861
M shell, 1128, 1128
 multiloop circuits, 707, 714, 714–720
 resistances in parallel, 714, 714–720
 multimeter, 720
 multiplication, of vectors, 47–52, 48
 multiplication factor, nuclear reactors, 1203
 multiplicity, of configurations in statistical mechanics, 551
 muon neutrinos, 1227, 1227t
 muons, 1029–1030, 1054, 1219, 1227t, 1227
 decay, 1225
 from proton-antiproton annihilation, 1224t, 1224
 musical sounds, 456–459
 mutual induction, 813–815, 814

- N**
- nanotechnology, 1083
- National Institute of Standards and Technology (NIST), 5
- natural angular frequency, 402, 836
- series *RLC* circuits, 846, 846
- NAVSTAR satellites, 1022
- n* channel, in MOSFET, 1159
- near point, 943, 943
- negative charge, 562, 562
- negative charge carriers, 684
- drift speed, 686
- negative direction, 13, 13
- negative lift, in race cars, 126–127, 127
- negative terminal, batteries, 658, 658, 706
- net force, 92
- net torque, 259
- net wave, 425
- net work, 144
- net work per cycle
- Carnot engine, 544
- neutral equilibrium, potential energy curves, 178
- neutral pion, 1023
- neutrinos, 1180
- and beta decay, 1180–1181
- in fusion, 1209–1210
- as leptons, 1221
- as nonbaryonic dark matter, 1240
- from proton-antiproton annihilation, 1224t
- neutron diffraction, 1069
- neutron excess, 1169
- neutron generation time, in nuclear reactors, 1215
- neutron number, 1168
- selected nuclides, 1168t
- neutron rich fragments, 1197
- neutrons, 563, 1219
- accelerator studies, 748
- balance in nuclear reactors, 1202–1203, 1203
- charge, 571, 571t
- collision with protons, 1190
- control in nuclear reactors, 1202–1205, 1203
- discovery of, 1236
- and eightfold way, 1231t
- as fermions, 1220
- formation in early universe, 1241
- as hadrons, 1221
- magnetic dipole moment, 754
- and mass number, 572
- as matter wave, 1068
- spin angular momentum, 871
- neutron stars, 82, 352, 1190
- density of core, 360t
- escape speed, 341t
- magnetic field at surface, 738t
- newton (N), 88
- Newton, Isaac, 87, 330, 342, 991
- Newtonian form, of thin-lens formula, 956
- Newtonian mechanics, 87, 1071, 1218
- Newtonian physics, 1083
- newton per coulomb, 580
- Newton's first law, 87–91
- Newton's law of gravitation, 330–333, 331
- Kepler's laws follow from, 342
- Newton's laws, 87
- applying in problem solving, 100–105
- Newton's rings, 987, 987
- Newton's second law, 91–94
- angular form, 285–286
- applying in problem solving, 100–105
- for rotation, 260–262
- system of particles, 206–209, 207
- in terms of momentum, 210
- translational and rotational forms contrasted, 263t, 289t
- units in, 92t
- Newton's third law, 98
- NMR spectrum, 1121, 1121
- noble gases, 1126, 1186
- nodes, 431, 431, 433
- nonbaryonic dark matter, 1240
- nonconducting sheet
- Gauss' law, 617, 617
- nonconductors, 563
- nonconservative forces, 168
- noninertial frame, 89
- nonlaminar flow, 371
- nonpolar dielectrics, 672
- nonpolar molecules, 638
- nonquantized portion, of energy level diagram, 1092, 1092
- nonsteady flow, 371
- nonuniform displacement current, 885
- nonuniform displacement-current density, 885
- nonuniform electric field, 608, 884
- nonuniform electric flux, 884
- nonuniform magnetic field
- diamagnetic materials, 875
- and loop model for electrons, 873–874, 873, 874
- nonviscous flow, 371
- normal (optics), 904–906, 905, 906
- normal force, 96–98, 97
- normalizing, wave function, 1089
- north magnetic pole, 870
- north pole, magnets, 738–739, 739, 862, 862
- n*-type semiconductors, 1152, 1152–1153. *See also p-n junctions*
- nuclear angular momentum, 1172
- nuclear binding energy, 1171–1172, 1172, 1198, 1199
- and excitation energy, 1110
- selected nuclides, 1168t
- nuclear energy, 1195
- nuclear energy levels, 1172
- nuclear fission, 1172, 1196–1201, 1199
- nuclear force, 1173–1174
- nuclear fusion, *See thermonuclear fusion*
- nuclear magnetic moment, 1172
- nuclear magnetic resonance (NMR), 1121
- nuclear magnetic resonance (NMR) spectrum, 1121, 1121
- nuclear power plant, 1202, 1204
- nuclear radii, 1170
- nuclear reactions, 1043
- nuclear reactors, 1202–1205
- natural, 1206, 1206
- nuclear spin, 1172
- nuclear weapons, 1172
- nucleons, 1168, 1221
- magic nucleon numbers, 1186
- nucleus, 563, 1165–1168
- models, 1184–1186, 1185
- radioactive decay, 572, 1219–1220
- nuclides, 1167–1168, 1168t. *See also radioactive decay*
- magic nucleon numbers, 1185–1186
- organizing, 1169–1170, 1169
- nuclides chart, 1169–1170, 1169
- and radioactivity, 1181–1182, 1182
- number density of charge carriers, 742, 1143
- semiconductors, 1150–1152
- silicon and copper compared, 1143t
- number density of conduction electrons, 1145
- O**
- object, 926, 926–927
- object distance, 926
- objective
- compound microscope, 944, 944
- refracting telescope, 945, 945
- occupancy probability, 1147–1148, 1148
- occupied levels, 1122
- occupied state density, 1149, 1149–1150
- ohm (unit), 689
- ohmic losses, 850
- ohmmeter, 689
- ohm-meter, 689
- Ohm's law, 692, 692
- microscopic view of, 693, 693–694
- oil slick, interference patterns from, 973
- one-dimensional elastic collisions, 221, 221–223
- one-dimensional electron traps multiple electrons, 1122
- single electron, 1084–1096
- one-dimensional explosions, 216, 216
- one-dimensional inelastic collisions, 218, 218–221, 219
- one-dimensional infinite potential well, 1085
- one-dimensional motion, 13
- acceleration, 18–27
- average acceleration, 18
- average speed, 16
- average velocity, 14–16
- constant acceleration, 22, 22–25
- displacement, 13–14
- free-fall acceleration, 25–26
- graphical integration, 27, 27–28
- instantaneous acceleration, 18
- instantaneous velocity, 17–18
- position, 13, 13–14
- relative, 73, 73–74
- Schrödinger's equation, 1071
- speed, 18
- velocity, 17–18

- one-dimensional variable
 - force, 151–152, 152
 - one-half rule, for intensity of transmitted polarized light, 902
 - one-way processes, 536
 - open-tube manometer, 365, 365–366
 - optical fibers, 912, 1131
 - with junction lasers, 1158
 - optical instruments, 943–946
 - optical interference, 959. *See also* interference
 - and rainbows, 962, 962
 - optics, 889
 - orbital angular momentum, 872, 873, 1115–1117, 1116
 - and orbital quantum number, 1115t
 - orbital energy, 1098
 - orbital magnetic dipole moment, 872, 1115–1117, 1116
 - diamagnetic materials, 875–876
 - ferromagnetic materials, 875, 879
 - paramagnetic materials, 877–878
 - orbital magnetic quantum number, 873, 1101t, 1102, 1115t
 - orbital quantum number, 1101t, 1102, 1115t, 1143
 - orbital radius, 1097–1098
 - order numbers, diffraction gratings, 1006
 - order of magnitude, 4
 - origin, 13
 - oscillating circuit, 826
 - oscillation mode, 433–434
 - oscillations, 386. *See also* pendulums; simple harmonic motion; waves
 - forced, 402, 836–837, 837
 - free, 402
 - LC* circuits, 826–829, 827
 - out of phase
 - ac circuits, 842t
 - capacitive load, 838–839
 - inductive load, 840
 - simple harmonic motion, 390
 - sound waves, 451
 - thin-film interference, 975
 - waves, 425
 - overpressure, 366
 - oxygen, 523
 - distribution of molecular speeds at 300 K, 517
 - heats of transformation, 487t
 - molar specific heat and degrees of freedom, 524t
 - molar specific heat at constant volume, 520t
 - paramagnetism of liquid, 877
 - RMS speed at room temperature, 512t
- P**
- pair production, 572–573
 - pancake collapse, of tall building, 237
 - parallel-axis theorem, for calculating rotational inertia, 254–255, 255
 - parallel circuits
 - capacitors, 662–663, 663, 715t
 - inductors, 822
 - resistors, 714, 714–717, 715t
 - summary of relations, 715t
 - parallel components, of unpolarized light, 912
 - parallel currents, magnetic field between two, 770, 770–771
 - parallel-plate capacitors, 656, 656
 - calculating capacitance, 659
 - with dielectric, 671, 671–674, 672
 - displacement current, 866–868, 867
 - energy density, 668
 - induced magnetic field, 863–866
 - paramagnetic material, 875, 875, 877–878
 - paramagnetism, 875, 877, 877–878
 - parent nucleus, 572
 - partial derivatives, 419, 450, 895
 - partially occupied levels, 1122
 - partially polarized light, 901, 902
 - particle accelerators, 747–749, 1218, 1219, 1219
 - particle-antiparticle annihilation, 1222
 - particle detectors, 1219, 1219
 - particles, 13, 571
 - antiparticles, 1222, 1224
 - basic forces and messenger particles, 1235–1237
 - bosons, 1220–1221, 1221
 - circulating charged, 744–747, 745, 746
 - conservation of baryon number, 1229
 - conservation of lepton number, 1228
 - conservation of strangeness, 1231
 - decay processes, 1222–1227
 - fermions, 1220–1221, 1221
 - formation in early universe, 1241
 - hadrons, 1221, 1228–1229
 - leptons, 1221, 1227t, 1227–1228
 - magnetic force on, 736–738, 737
 - particle nature of matter, 1069–1070, 1069
 - quark model, 1232–1235, 1233
 - Standard Model, 1220–1222
 - particle systems
 - angular momentum, 288
 - center of mass, 201–206, 202, 205
 - linear momentum, 211
 - Newton's second law for, 206–209, 207
 - pascal (Pa), 361, 446, 900
 - Pascal's principle, 366, 366–367
 - Paschen series, 1100, 1101
 - path-dependent quantities, 490
 - path-independent quantities
 - conservative forces, 168–170, 169
 - gravitational potential energy, 340–341
 - work done by electrostatic force, 628
 - path length difference
 - double-slit interference, 965, 965–966, 971–972
 - single-slit diffraction, 993, 993, 994, 995
 - sound waves, 451
 - thin-film interference, 973
 - Pauli exclusion principle, 1121
 - bosons don't obey, 1220
 - and energy levels in crystalline solids, 1143
 - fermions obey, 1220
 - and Fermi speed, 1145
 - nucleons, 1185
 - and periodic table, 1125
 - pendulums
 - as angular simple harmonic oscillator, 394, 394–395
 - ballistic, 220, 220
 - conservation of mechanical energy, 173–175, 174
 - physical, 396–397, 397
 - simple, 395, 395–396
 - torsion, 394, 394
 - underwater swinging (damped), 400
 - perfect engines, 546, 546
 - perfect refrigerators, 549, 549
 - perihelion distance, 344
 - period
 - circulating charged particle, 744–747
 - LC* oscillations, 826
 - linear simple harmonic oscillator, 390
 - simple harmonic motion, 387, 388, 389
 - sound waves, 449
 - waves, 416, 416–417
 - periodic motion, 386
 - periodic table, 1057, 1113
 - building, 1124–1127
 - x rays and ordering of elements, 1127–1130
 - period of revolution, 71
 - permanent electric dipole moment, 638
 - permanent magnets, 736
 - permeability constant, 765
 - permittivity constant, 566
 - perpendicular components, of unpolarized light, 912
 - phase
 - alternating current, 842t
 - simple harmonic motion, 388, 388
 - waves, 415, 415
 - phase angle
 - alternating current, 842t
 - simple harmonic motion, 388, 388
 - phase change, 486
 - phase constant
 - alternating current, 842t, 848
 - series *RLC* circuits, 844–845, 845
 - simple harmonic motion, 388, 388
 - waves, 417, 417
 - phase difference
 - double-slit interference, 965, 969–970, 971–972
 - Michelson's interferometer, 980
 - optical interference, 961–962
 - and resulting interference type, 427t
 - single-slit diffraction, 995
 - sound waves, 451
 - thin-film interference, 975
 - waves, 425–426
 - phase-shifted
 - reflection phase shifts, 974, 974

- sound waves, 451
- waves, 425–426
- phasor diagram, 428–429, 429
- phasors, 428–430, 430
 - capacitive load, 838, 838, 839
 - double-slit interference, 971
 - inductive load, 840, 840, 841
 - resistive load, 838
 - series *RLC* circuits, 844–845, 845
 - single-slit diffraction, 995–999, 996, 998
- photodiode, 1158
- photoelectric current, 1059
- photoelectric effect, 1059–1062, 1061
- photoelectric equation, 1060–1061
- photoelectrons, 1059
- photomultiplier tube, 1067
- photon absorption, 1058, 1113
 - absorption lines, 1100, 1101
 - for electrons to gain energy, 1086
 - energy changes in hydrogen atom, 1098
 - lasers, 1132
- photon emission, 1058, 1113
 - for electrons to lose energy, 1086
 - emission lines, 1100, 1101
 - energy changes in hydrogen atom, 1098
 - lasers, 1132, 1132–1133
 - stimulated emission, 1132, 1133
- photons
 - as bosons, 1220
 - in early universe, 1241
 - energy of, 1058
 - momentum, 1062, 1062–1065, 1063
 - and photoelectric effect, 1059–1062
 - as quantum of light, 1057–1059
 - single-photon version of double-slit interference, 1065, 1065–1066
 - single-photon wide-angle version of double-slit interference, 1067, 1067
 - virtual, 1235–1236
- physical pendulum, 396–398, 398
- picofarad, 657
- piezoelectricity, 1075
- pinhole diffraction, 990
- pions, 1024, 1055, 1219
 - decay, 1225
 - and eightfold way, 1231t
- as hadrons, 1221
- as mesons, 1221
- from proton-antiproton annihilation, 1224t, 1222–1225
- pitch, 359
- pitot tube, 383
- planar symmetry, Gauss' law, 617, 617–618
- planar waves, 446
- Planck constant, 1058
- plane-concave lens, 957
- plane-convex lens, 957
- plane mirrors, 926–928, 927
- plane of incidence, 906
- plane of oscillation, polarized light, 901, 901
- plane of symmetry, center of mass of solid bodies with, 203
- plane-polarized waves, 901
- plane waves, 891
- plastics
 - as insulators, 563
 - rubbing rod with fur, 563
- plates, capacitors, 657, 657–658
- plate tectonics, 13
- plum pudding model, of atom, 1166
- pn* junction diode, 692, 697
- p-n* junctions, 1154–1156, 1155
 - junction lasers, 1158, 1158
 - junction rectifiers, 1156, 1156–1157, 1164
- light-emitting diodes (LEDs), 1157–1158, 1158
- point charges
 - Coulomb's law, 565, 565–572
 - in electric field, 592–593
 - electric field due to, 582, 582–583
 - electric potential due to, 635, 635
 - electric potential due to group of, 636, 636–637
 - electric potential energy of system, 642, 642–644
- point image, 926–927
- point of symmetry, center of mass of solid bodies with, 203
- point source, 446
 - isotropic, 454, 898
 - light, 898, 926
- polar dielectrics, 671
- polarity
 - of applied potential difference, 692
 - of Earth's magnetic field, reversals in, 870, 870
- polarization, 901, 901–904, 902, 903
 - by reflection, 912, 912–913
- polarized light, 901, 901–902
 - intensity of transmitted, 902, 902–904, 903
- polarized waves, 901, 901–904, 902
- polarizer, 903
- polarizing direction, 903, 903
- polarizing sheets, 902, 902–903, 903, 912
- polar molecules, 638
- Polaroid filters, 902
- pole faces, horseshoe magnet, 739
- polyatomic molecules, 520
 - degrees of freedom, 523, 523–524, 524t
 - molar specific heats at constant volume, 520t
- polycrystalline solids, 880
- population inversion, 1133, 1139
 - junction lasers, 1158
- porcelain
 - dielectric properties, 69t, 671
- position, 263t. *See also* angular position; displacement
 - one-dimensional motion, 13, 13–14
 - relating linear to angular, 250
 - simple harmonic motion, 388
 - two- and three-dimensional motion, 58–59, 59
- position vector, 58, 58
- positive charge, 672, 562
- positive charge carriers, 684
 - drift speed, 686
 - emf devices, 706
- positive direction, 13, 13
- positive ions, 563
- positive kaons, 1030
- positive terminal, batteries, 658, 658, 706
- positrons
 - antihydrogen, 1222
 - bubble chamber tracks, 572, 737
 - electron-positron annihilation, 572, 572, 1222
 - in fusion, 1207–1208
- potential, 629–631. *See also* electric potential
- potential barrier, 1074, 1074, 1200
- potential difference, 629–631
 - across real battery, 712–713
- capacitors, 659, 659–662
- capacitors in parallel, 662–663, 663
- capacitors in series, 664, 663–666, 842–844
- Hall, 741
- LC* oscillations, 826
- and Ohm's law, 692
- power, potential, and emf, 712–713
- resistors in parallel, 714–715
- resistors in series, 710, 710, 842–844
- RL* circuits, 807, 807–810
- single-loop circuits, 707, 707–709
 - between two points in circuit, 711, 711–713, 712
- potential energy. *See also* elastic potential energy; gravitational potential energy
 - and conservation of mechanical energy, 173, 173–175, 174
 - and conservation of total energy, 183–184
 - defined, 167
 - determining, 170–172
 - electric dipole, 594–595
 - satellites in orbit, 345, 345
 - simple harmonic motion, 392, 393
 - and work, 167, 167–170
 - yo-yo, 280–281
- potential energy barrier, 1074, 1074
- potential energy curves, 176–179, 177
- potential energy function, 176–178, 177
- potential method, of calculating current in single-loop circuits, 707–708
- potential well, 178
- potentiometer, 670
- pounds per square inch (psi), 361
- power, 155–156, 156, 185, 263t
 - in alternating current circuits, 847–848
 - in direct current circuits, 695–696
 - and emf in circuits, 712–713
 - in rotation, 262
 - traveling wave on stretched string, 421, 421–423
- power factor, 849
- power lines
 - transformers for, 850

- power transmission systems, 682
transformer requirements, 850
Poynting vector, 897–899, 898
precession, of gyroscope, 293–295, 295
precession rate, of gyroscope, 294
prefixes, for SI Units, 2t
pressure
fluids, 359–361
and ideal gas law, 508–511
measuring, 365, 365–366
and RMS speed of ideal gas, 511–513
scalar nature of, 38
as state property, 537
triple point of water, 478
work done by ideal gas at constant, 510
pressure amplitude
sound waves, 449, 449
pressure field, 580
pressure sensor, 360
pressurized-water nuclear reactor, 1204, 1204
primary coil, transformer, 850
primary loop, pressurized-water reactor, 1204, 1204
primary rainbow, 909, 922
primary rainbows and optical interference, 962, 962
primary winding, transformer, 850
principal quantum number, 1102, 1101t, 1115t, 1143
principle of conservation of mechanical energy, 173
principle of energy conservation, 140
principle of equivalence, 347
principle of superposition, 88
and electrostatic force, 566
for gravitation, 333–334
for waves, 425, 425
prism, 908, 908
angle of minimum deviation, 920
probability
and entropy, 552–553
probability density, 1071–1072
barrier tunneling, 1074–1075
finding, 1072
trapped electrons, 1088–1090, 1089
probability distribution function, 516
probability of detection
hydrogen electron, 1102–1103
trapped electrons, 1088–1089
probability wave
light as, 1066–1067
matter wave as, 1068
projectile, 64
elastic collisions in one dimension with moving target, 222–223
elastic collisions in one dimension with stationary target, 221–222
inelastic collisions in one dimension, 218
series of collisions, 213
single collision, 211–212
projectile motion, 64–70, 65, 66
proper frequency, 1040
proper length, 1032
proper period, 1042
proper time, 1028
proper wavelength, 1041
proton number, 1168
selected nuclides, 1168t
proton-proton (p-p) cycle, 1209–1210, 1209
protons, 563, 1219
accelerator studies, 748
and atomic number, 572
as baryons, 1221
charge, 571, 571t
decay rate, 1245
in equilibrium, 569
as fermions, 1220
formation in early universe, 1241
in fusion, 1207–1213
as hadrons, 1221
magnetic dipole moment, 754, 754t
mass energy, 1043t
and mass number, 572
as matter wave, 1068, 1069, 1083
spin angular momentum, 871
ultrarelativistic, 1047
in well, 652
proton synchrotrons, 749
p subshells, 1125, 1126
p-type semiconductors, 1153, 1153
pulleys, massless frictionless, 98, 98
Newton's laws applications, 100, 100–101
pulsars, 268
secondary time standard based on, 9
pulse, wave, 414, 414
- Q**
quadrupole moment, 600
quanta, 1057
quantization, 1057, 1083
electric charge, 571
energy in electron traps, 1084–1088
orbital angular momentum, 872
orbital energy, 1098
spin angular momentum, 871
quantum, 1057
quantum chromodynamics (QCD), 1236
quantum corrals, 1094, 1094
quantum dots, 1083, 1094, 1094
quantum electrodynamics (QED), 872, 1235
quantum jump, 1086
quantum mechanics, 87, 1057
quantum numbers, 1084, 1115t
charge, 1224
for hydrogen, 1102–1103
and Pauli exclusion principle, 1121
and periodic table, 1124–1126
quantum physics, 1057. *See also* electron traps; Pauli exclusion principle; photons; Schrödinger's equation
barrier tunneling, 1074, 1074–1076
and basic properties of atoms, 1112–1114, 1113
confinement principle, 1084
correspondence principle, 1089, 1106
Heisenberg's uncertainty principle, 1072, 1073, 1236
hydrogen wave function, 1099–1106
matter waves, 1083
nucleus, 1165
occupancy probability, 1147–1148, 1148
particles, 1219
and solid-state electronic devices, 1142
quantum states, 1083, 1114
degenerate, 1095
density of, 1146–1147, 1147
density of occupied, 1149, 1149–1150
hydrogen with $n = 2$, 1105, 1105–1106
quantum theory, 523, 525, 1057, 1083
- quantum transition, 1086
quantum tunneling, 1074, 1074–1076
quark family, 1233t
quark flavors, 1232, 1236–1237
quarks, 578, 654, 747, 1232–1235, 1233, 1233t
charge, 571
formation in early universe, 1241
quasars, 1053, 1237
gravitational lensing associated with, 348
quicksand, 385
- R**
R-value, 494
race cars
collision with wall, 214, 214
negative lift in Grand Prix cars, 126–127, 127
pit stop fuel dispenser fire, 723, 723–724
rad (unit), 1184
radar waves, 413
radial component
of linear acceleration, 251
of torque, 259
radial probability density, 1103, 1103, 1106
radians, 43, 242
radiated waves, 890
radiation (electromagnetic), 890
radiation dosage, 1184
radiation heat transfer, 496–497
radiation pressure, 899–900
radiation therapy, 1190
radioactive dating, 1183, 1183
radioactive decay, 572, 1169, 1219–1220
alpha decay, 1177–1179, 1178
beta decay, 1180–1181, 1181, 1234
muons, 1029
and nuclidic chart, 1181–1182, 1182
process, 1174–1177
radioactive elements, 1165
radioactive wastes, 1204, 1204
radionuclides, 1169
radiation dosage, 1184
radio waves, 413, 889, 890
Doppler effect, 461
radium, 1191
radius of curvature
spherical mirrors, 928, 928 – 929, 929
spherical refracting surfaces, 934, 934

- radon, 1165, 1191
rail gun, 770, 770–771, 790, 790
rainbows, 888, 908–909, 909, 922
and optical interference, 962, 962
ramp
rolling down, 278–279, 279
randomly polarized light, 901, 901–902
range, in projectile motion, 64, 68
rare earth elements, 1129
paramagnetism, 875
rattlesnake, thermal radiation sensors, 496
ray diagram, 932, 932
Rayleigh's criterion, 1000, 1000–1001
rays, 446, 446
locating extended object images by drawing, 939, 939
locating images by drawing, 932, 932
RBE (relative biology effectiveness) factor, 1184
RC circuits, 720, 720
capacitor charging, 720, 720–721
capacitor discharging, 720, 720, 722–724
real batteries, 707, 707
internal resistance, 711, 711
potential difference across, 711, 711–712
real emf devices, 707, 707
internal resistance, 711
real engines
efficiency, 546, 549–550
real fluids, 371
real focal point, 929, 929
real images, 924–925
spherical mirrors, 930
spherical refracting surfaces, 934, 934
thin lenses, 938, 938
real solenoid, 776
recessional speed parameter, Doppler shift, 1246
rechargeable batteries, 707
recharging batteries, 712
red shift, 1041
reference circle, 399
reference configuration, for potential energy, 171, 628
reference frames, 73–74
inertial, 89–90
noninertial, 89
reference line, 242, 242
reference particle, 399
reference point, for potential energy, 171
reference potential energy, 628
reflected light, 904
reflected radiation, 899
reflected ray, 904–905, 905
reflecting planes, 1012, 1012
reflection, 904–907, 905, 906
law of, 906
polarization by, 912, 912–913
of standing waves at boundary, 432, 432
total internal, 911, 911–912
reflection phase shifts, 974, 974–976
refracted light, 906
refracted ray, 906, 906
refracting telescope, 945, 945–946
refraction, 904–910, 906. *See also* index of refraction and chromatic dispersion, 907, 907–908, 908
law of, 906, 909–912, 909
refrigerators, 548, 548–549
relative biology effectiveness (RBE) factor, 1184
relative motion
in one dimension, 73–74, 73
in two dimensions, 74–75, 74
relativity, 1057, 1218
brief description, 1023
Doppler effect for light, 1040–1042, 1041
and energy, 1043–1047
general theory of, 347–348, 1022–1023, 1029
and kinetic energy, 1045, 1045–1047
of length, 1031, 1031–1034, 1037
Lorentz factor, 1028, 1028, 1029, 1043
Lorentz transformation, 1035, 1035–1036
and mass energy, 1043–1045, 1043t
measuring events, 1024–1025, 1025
and momentum, 1042
postulates, 1023
of simultaneity, 1025–1026, 1026, 1037
special theory of, 87, 893, 1022, 1029
speed of light postulate, 1023–1024
of time, 1027–1031, 1027, 1037, 1042
of total energy, 1044–1045
of velocities, 1039, 1039
relativity postulate, 1023
relaxed state, of spring, 149, 149
released energy, from fusion reaction, 1045
rem (unit), 1184
resistance, 689–691
alternating current, 842t
calculating from resistivity, 690, 690
Ohm's law, 692, 692–694
parallel circuits, 714, 714–719
RC circuits, 720–723
RL circuits, 807–810
RLC circuits, 833–834, 842–844
series circuits, 710, 710, 842–844
resistance rule, 709
resistivity, 689, 1142
calculating resistance from, 690, 690
Ohm's law, 693
selected materials at room temperature, 689t
semiconductors, 1151
silicon and copper compared, 696t, 696–697, 1143t
resistors, 689, 689–690
with ac generator, 836, 836–837
in multiloop circuits, 714, 714–719, 717
Ohm's law, 692, 692–694
in parallel, 714, 714–719
phase and amplitude relationships for ac circuits, 842t
power dissipation in ac circuits, 847–848
and power in circuits, 695–696
RC circuits, 720, 720–723
RL circuits, 807, 807–810
RLC circuits, 833, 833–834, 842–844, 843
in series, 710, 710, 842–844, 843
series RLC circuits, 842–844, 843
in single-loop circuits, 707, 707–710
work, energy, and emf, 706, 706–707
resolvability, 1000, 1000–1001
resolving power
diffraction gratings, 1009–1011, 1010
highest for microscope, 1080
refracting telescope, 945, 945
resolving vectors, 41
resonance
forced oscillations, 402, 837
series RLC circuits, 845–846, 846
and standing waves, 433, 433–435
resonance capture, of neutrons in nuclear reactors, 1202
resonance condition cyclotrons, 748
resonance curves
series RLC circuits, 845–846, 846
resonance peak, 402, 1121
resonant frequencies, 433, 433, 434
musical sounds, 457, 458
response time, nuclear reactor control rods, 1203
rest energy, 1043
rest frame, 1029
rest length, 1032
restoring force, 149
resultant, of vector addition, 39
resultant force, 92
resultant torque, 259
resultant wave, 425, 425
reverse saturation current, junction rectifiers, 1164
reversible processes, 537–541
right-handed coordinate system, 44, 44
right-hand rule, 247, 247
Ampere's law, 771, 771
angular quantities, 247, 247
displacement current, 866–867, 867
induced current, 795, 795
Lenz's law, 794, 794
magnetic dipole moment, 753–754, 754
magnetic field due to current, 765, 765
magnetic force, 737, 737
magnetism, 770
vector products, 50, 50, 769
rigid bodies
angular momentum of rotation about fixed axis, 288–290, 289
defined, 241
elasticity of real, 315
ring charge distributions, 586, 586–587, 590
RL circuits, 807, 807–810, 809
RLC circuits, 833, 833–834
series, 842–846, 843

- RMS (root-mean-square) current, in ac circuits, 847
- RMS (root-mean-square) speed, of ideal gas, 511, 511–513, 516
- selected substances, 512t
- rock climbing
- crimp hold, 324, 324
 - energy conservation in descent using rings, 183, 183
 - energy expended against gravitational force climbing Mount Everest, 199
 - friction coefficients between shoes and rock, 119
 - lie-back climb along fissure, 322, 322
- rockets, 224–226, 225
- roller-coaster headache, 252
- roller coasters
- maximum acceleration, 20
- rolling
- down ramp, 278, 278–280
 - forces of, 278–280, 279
 - friction during, 278, 278
 - kinetic energy of, 277, 277
 - as pure rotation, 276–277, 276
 - as translation and rotation combined, 275–277, 277
 - yo-yo, 280–281, 281
- room temperature, 476
- root-mean-square (RMS) current, in ac circuits, 847
- root-mean-square (RMS) speed, of ideal gas, 511, 511–513
- and distribution of molecular speeds, 516
 - selected substances, 512t
- rotation, 242. *See also* rolling; torque; terms beginning with “angular”
- angular momentum of rigid body rotating about fixed axis, 288–290, 289
 - conservation of angular momentum, 290–293, 291, 293
 - constant angular acceleration, 248–249
 - kinetic energy of, 253–254, 254
 - Newton’s second law for, 260–261
 - relating linear and angular variables, 250–253, 251
 - in rolling, 275–277, 276
- rotational equilibrium, 307
- rotational inertia, 253, 263t
- calculating, 254–258
- rotational kinetic energy, 253
- of rolling, 278
 - and work, 262–264
 - yo-yo, 280–281
- rotational symmetry, 582
- rotational variables, 241–246, 289t
- vector nature of, 246–247
- rotation axis, 242, 242
- Rotor (amusement park ride), 249
- Rowland ring, 879, 879
- rubber band, entropy change on stretching, 542
- rulers, 1
- rulings, diffraction grating, 1003, 1003
- Rutherford, Ernest, 1081, 1165
- R*-value, 494
- Rydberg constant, 1099
- S**
- satellites
- charge buildup in radiation belt, 623
 - geosynchronous orbit, 354
 - gravitational potential energy, 339
 - Kepler’s laws, 342–344
 - orbits and energy, 345, 345–346
- saturation magnetization, 877, 878, 879, 887
- scalar components, 44
- scalar fields, 580
- scalar product, 48
- scalars, 38
- multiplying vectors by, 47–48
- scanning tunneling microscope (STM), 1075–1076, 1075
- quantum corral, 1094, 1094
- scattering
- polarized light, 903
 - x rays, 1011, 1011
- schematic diagrams, 658
- Schrödinger’s equation, 1071–1072, 1083
- electron in finite well, 1091–1092
 - electron in one-dimensional infinite well, 1088–1091
 - electron in rectangular box, 1095
 - electron in rectangular corral, 1095
 - finding probability density, 1072
 - and hydrogen, 1099–1106
- hydrogen ground state, 1102–1103, 1103, 1115
- numerical solution for multicomponent atoms, 1124
- scientific notation, 2
- scintillation counter, 1248
- seat of emf, 705
- secondary coil, transformer, 850–851
- secondary loop, pressurized water reactor, 1204, 1204
- secondary maxima, diffraction patterns, 990, 990
- secondary rainbows, 909, 909, 922, 962
- secondary standards, 3
- secondary winding, transformer, 850–851
- second law of thermodynamics, 541–542
- second minima, interference patterns, 967
- second-order bright fringes, interference patterns, 966
- second-order dark fringes, interference patterns, 967
- second-order line, 1006
- second reflection point, 921, 921
- second side maxima, interference patterns, 966
- secular equilibrium, 1191
- seismic waves, 413
- self-induced emf, 806, 806
- self-induction, 806, 806–807
- mutual induction contrasted, 813
- semi-classical angle, 1116
- semiconducting devices, 696
- semiconductors, 563. *See also* *p-n* junctions; transistors
- doped, 1152–1153, 1153
 - electrical properties, 1150–1151, 1151
 - light-emitting diodes (LEDs), 1157–1158, 1158
 - nanocrystallites, 1093, 1093
 - resistivities of selected, 689t
 - unit cell, 1142
- semimajor axis, of orbits, 342–343, 343
- planets of Solar System, 344t
- series circuits
- capacitors, 663–666, 664, 715t, 842–844, 843
 - inductors, 842–844, 843
 - RC*, 720, 720–723
- resistors, 710, 710, 715t, 842–844, 843
- RLC*, 833, 833–834, 842–846, 843
 - summary of relations, 715t
- series limit, 1100, 1101
- shearing stress, 315, 315–316
- shear modulus, 316
- shells, 1106, 1115, 1115t
- and characteristic x-ray spectrum, 1128
 - and energy levels in crystalline solids, 1143
 - and periodic table, 1126–1127
- shell theorem, 332
- shock waves, 465, 465
- short wave radiation, 891
- side maxima
- diffraction patterns, 990, 990
 - interference patterns, 966
- sievert (unit), 1184
- sigma particle, 1219
- and eightfold way, 1231t
 - and strangeness, 1231
- sign
- acceleration, 20
 - displacement, 13–14
 - heat, 484
 - velocity, 20, 27
 - work, 143
- silicon
- effect of doping on Fermi energy, 1163
 - electric properties compared to copper, 696t, 696–697, 1143t, 1151
 - electron number density in conduction band, 1164
 - germanium compared to, 1162
 - in MOSFETs, 1159
 - properties of *n*- and *p*-doped compared, 1153t
 - resistivity of pure, 689t
 - as semiconductor, 563, 696–697, 1150–1151
 - temperature coefficient of resistivity, 1151
 - unit cell, 1142, 1142
- silicon, *n*-type resistivity, 689t
- silicon, *p*-type resistivity, 689t
- silk
- rubbing glass rod with, 562, 562–562, 572
- simple harmonic motion, 386–389, 387, 388
- acceleration, 389, 389
 - angular, 394, 394–395
 - damped, 400, 400–401, 401
 - energy, 392–394, 393

- force law for, 390–392
- pendulums, 395, 395–398, 397
- and uniform circular motion, 398–399, 399
- velocity, 388, 388, 389, 389
- waves produced by, 413–414
- simple harmonic oscillators. *See also* pendulums
- angular, 394, 394–395
- linear, 390, 390–392
- simple magnifying lens, 943, 943
- simple pendulum, 395, 395–396
- simultaneity
- consequences of Lorentz transformation equations, 1037
- relativity of, 1025–1026, 1026
- sine, 43, 43
- single-component forces, 88
- single-loop circuits, 836
- current calculation, 707, 707–708
- internal resistance, 709, 709
- resistances in series, 710, 710
- single-slit diffraction, 963–964, 964, 992, 992–995, 993
- intensity, 995–999, 996, 998
- sinusoidal waves, 414–417, 415, 417
- SI units, 2
- skateboarding, motion analyzed, 67
- slab
- rotational inertia, 255t
- sliding block, 100, 100–101
- sliding friction, 118, 118
- slit diffraction, 990
- slope, of line, 15–16, 16
- Snell's law, 906, 959
- soap bubbles, interference patterns from, 973, 976, 976
- sodium, 1126, 1186
- sodium chloride, 1126
- index of refraction, 906t
- x-ray diffraction, 1012, 1012
- sodium doublet, 1140
- sodium vapor lamp, 1058
- soft reflection, of traveling waves at boundary, 432
- solar system, 1243
- solar wind
- and comets, 917, 917
- solenoids, 774
- induced emf, 794–795
- inductance, 805–806
- magnetic energy density, 812
- magnetic field, 776, 776–777
- solid bodies
- center of mass, 203–206
- solids
- compressibility, 318
- electrical properties of crystalline, 1142, 1142–1143
- heat absorption, 485–488
- specific heats of selected, 485t
- speed of sound in, 447t
- thermal conductivity of selected, 494t
- thermal expansion, 481–483, 482
- solid state, 486
- solid-state electronic devices, 1142
- sonar, 445
- sonic boom, 465
- sound intensity, 452–456, 453
- sound level, 452–456
- selected sounds, 454t
- sound waves, 413–414
- beats, 459, 459–460
- Doppler effect, 461–464, 462, 463
- interference, 451, 451–452
- sources of musical, 456–459, 457, 458
- speed of, 446–448, 447t
- supersonic speed, 465, 465
- traveling waves, 448, 448–450
- south pole, magnets, 738–739, 739, 862, 862
- space charge, 1155
- space curvature, 347, 347
- space time, 347, 1057, 1241
- spacetime coordinates, 1024–1025
- spark. *See* electric spark
- special theory of relativity, 87, 893, 1023, 1029
- specific heat, 485. *See also* molar specific heat
- selected materials, 485t
- speckle, 969
- spectral emission lines, 1082
- spectrum, 1101
- speed. *See also* angular speed
- average in one-dimensional motion, 16
- one-dimensional motion, 18
- relating linear to angular, 250
- in rolling, 276–277, 277
- traveling waves, 417–420
- speed of light, 893
- as ultimate speed, 1023
- speed of light postulate, 1023–1024
- speed of sound, 446–448
- and RMS speed in gas, 513
- in various media, 447t
- speed parameter, in time dilation, 1028, 1028
- spherical aberration, 946
- spherical capacitors
- calculating capacitance, 661–662
- spherical conductors
- Coulomb's law for, 566, 566–570
- spherically symmetric charge distribution, 635
- Gauss' law for, 619, 619
- spherical mirrors, 928, 928–930, 929, 930
- focal points, 929–930, 930
- images from, 928, 928, 930, 930–933, 946, 946–947
- spherical refracting surfaces, 934, 934–935
- images from, 934, 934–935, 946, 946–947
- spherical shell
- Coulomb's law for, 566, 566–570
- rotational inertia, 255t
- spherical symmetry, Gauss' law, 619, 619–620
- spherical waves, 446
- spin, 1115, 1115t, 1219–1221, 1220
- isospin, 1245
- nuclides, 1168t, 1172
- spin angular momentum, 871–872, 1115, 1115t, 1116–1117, 1117
- and spin quantum number, 1115t
- spin-down electron state, 871, 1117, 1120, 1120
- spin-flipping, 883, 1121, 1121
- spin magnetic dipole moment, 871–872, 872, 1116–1117, 1117
- diamagnetic materials, 875–876
- ferromagnetic materials, 875, 879
- paramagnetic materials, 875, 877–878
- spin magnetic quantum number, 871, 1115, 1115t, 1219–1220
- spin quantum number, 1115, 1115t, 1219–1220
- spin-up electron state, 871, 1117, 1120, 1120
- spontaneous emission, 1132, 1132–1133
- spontaneous otoacoustic emission, 470
- spring constant, 149
- spring force, 149–150
- as conservative force, 168, 168
- work done by, 149, 149–151
- spring scale, 96, 96
- sprites, 585, 585
- s subshells, 1125, 1126
- stable equilibrium potential energy curves, 178
- stable static equilibrium, 305–306, 306
- stainless steel
- thermal conductivity, 494t
- standard kilogram, 6, 6
- standard meter bar, 3
- Standard Model, of fundamental particles, 1220–1222
- standards, 1
- standing waves, 431, 431–432, 1083
- reflections at boundary, 432, 432
- and resonance, 433, 433–435
- stars, 1057. *See also* black holes; neutron stars
- orbiting, 355
- Doppler shift, 1040
- formation in early universe, 1241
- fusion in, 1172, 1207, 1208–1210, 1209
- preponderance of matter over antimatter, 1222
- rotational speed as function of distance from galactic center, 1240, 1240
- S2 star at center of Milky Way galaxy, white dwarfs, 341t, 360t
- state, 486
- state function
- entropy as, 539–540
- state properties, 537–538
- static equilibrium, 305, 305–306, 306
- examples, 309–312, 313–311
- fluids, 362, 362
- indeterminate structures, 314, 314
- requirements of, 306–307
- static frictional force, 116–118, 117
- rolling wheel, 278
- statistical mechanics, 550–553
- steady flow, 371
- steady-state current, 682
- series RLC circuits, 842

- Stefan-Boltzmann constant, 496
- step-down transformer, 851
- step-up transformer, 851
- Stern-Gerlach experiment, 1118, 1118–1120
- stick-and-slip, 118
- stimulated emission, 1132–1133
- from comets, 1140, 1140
- Stirling engines, 546, 546
- Stirling's approximation, 553
- stopping potential, photoelectric effect, 1059–1061, 1061
- straight line charge distributions, 590
- straight-line motion, *See* one-dimensional motion
- strain, 315–318, 516
- strain gage, 316, 316
- strangeness, 1231
- and hypercharge, 1245
- strange particles, 1230–1231, 1231
- strange quark, 1232, 1233t, 1233
- streamlines
- in electric fields, 685
- in fluid flow, 371, 372
- stress, 315, 315
- stress-strain curves, 316, 316
- stress-strain test specimen, 316
- stretched strings, 446
- energy and power of traveling wave on, 421, 421–423
- harmonics, 434–435
- resonance, 433, 433–435
- standing waves, 431, 431–432, 432
- transverse and longitudinal waves on, 414, 414
- wave equation, 423–424
- wave speed on, 420–421, 420
- strike-slip, 57
- strokes, 543
- strong force, 115, 1173, 1221
- conservation of strangeness, 1231
- messenger particle, 1235–1236
- strong interaction, 1224
- strong nuclear force, 572
- subcritical state, nuclear reactors, 1203
- submarines
- Project Seafarer, 915
- sonar, 445
- subshells, 1106, 1115, 1115t
- and energy levels in crystalline solids, 1143
- and periodic table, 1125
- f* subshells, 1125
- substrate, 1159
- subtraction
- of vectors by components, 45–46
- of vectors geometrically, 39–40, 40
- Sun, 1243
- convection cells in, 496
- density at center of, 360t, 1217
- escape speed, 341t
- fusion in, 1172, 1207, 1209–1210, 1209
- monitoring charged particles from, 682
- neutrinos from, 1181
- period of revolution about galactic center, 354
- pressure at center of, 361t
- radiation from, 889
- randomly polarized light, 901
- speed distribution of photons in core, 517
- superconductivity, 697
- superconductors, 563, 697, 697–698
- supercooling, 557
- supercritical state, nuclear reactors, 1203
- superluminal jets, galaxies, 1054
- supermassive black holes, 330
- supernova, 82, 341t, 1210, 1210, 1242–1243
- supernova SN1987a, 1181, 1181, 1210, 1210
- supernumeraries, 962, 962
- superposition principle for forces, *See* principle of superposition
- surface charge density, 586, 586t
- S waves, 468
- symmetric lenses, 938–939
- symmetry
- cylindrical, Gauss' law, 615, 615–616
- importance in physics, 605
- line, plane, point: center of mass of bodies with, 203
- planar, Gauss' law, 617, 617–618
- spherical, Gauss' law, 619, 619–620
- system, 92, 483
- systems of particles, *See* particle systems
- T**
- tangent, 43, 43
- tangential component
- of linear acceleration, 250–251
- of torque, 258
- target
- collisions in two dimensions, 224, 224
- elastic collisions in one dimension with moving, 222–223, 223
- elastic collisions in one dimension with stationary, 221, 221
- inelastic collisions in one dimension, 218
- series of collisions, 213, 213
- single collision, 211–212
- tattoo inks, magnetic particles in, 861, 861
- tau neutrinos, 1227, 1227t
- tau particle, 1227, 1227t
- telescopes, 943
- refracting, 945, 945–946
- surveillance satellites, 1017
- television, 869
- magnet applications, 735
- television waves, 413
- temperature, 476
- defined, 478
- and heat, 483
- and ideal gas law, 508–511
- measuring, 478–479
- and RMS speed of ideal gas, 511–513
- scalar nature of, 38
- as state property, 537
- work done by ideal gas at constant, 509, 509–510
- and zeroth law of thermodynamics, 478, 477
- temperature coefficient of resistivity, 691, 1143
- selected materials, 689t
- semiconductors, 1151
- silicon and copper compared, 696t, 1143t
- temperature field, 580
- temperature scales, 476, 476, 479–481
- compared, 480
- temporal separation, of events, 1027
- tensile stress, 316, 316
- tension force, 97–98, 98
- and elasticity, 316–317
- and wave speed on stretched string, 421
- terminals, batteries, 658, 706
- terminal speed, 122, 122–123
- tesla (unit), 737
- test charge, 580–581, 581
- Tevatron, 1234
- thermal agitation
- ferromagnetic materials, 879
- paramagnetic materials, 877
- thermal capture, of neutrons in nuclear reactor, 1202
- thermal conduction, 494–495, 495
- thermal conductivity, 494–495
- selected materials, 494t
- thermal conductor, 494
- thermal efficiency
- Carnot engines, 545–546
- Stirling engines, 546
- thermal energy, 168, 475
- and conservation of total energy, 183
- thermal equilibrium, 477
- thermal expansion, 481–483, 482
- thermal insulator, 494
- thermal neutrons, 1196–1201, 1202
- thermal radiation, 496–497
- thermal reservoir, 488, 489
- thermal resistance to conduction, 494
- thermodynamic cycles, 489, 490, 492
- thermodynamic processes, 488–491, 489
- graphical summary, 528
- thermodynamics, 477. *See also* entropy; irreversible processes
- first law, 491–494
- second law, 541–542
- zeroth law, 477, 477
- thermometers, 477
- constant-volume gas, 478, 478–479
- liquid-in-glass, 482
- thermonuclear bomb, 1211
- thermonuclear fusion, 1045, 1172, 1207–1208
- controlled, 1211–1213
- in Sun and stars, 1207, 1209–1210, 1209
- thermopiles, 706
- thermoscope, 477, 477
- thin films
- interference, 973, 973–979
- thin-lens approximation, 948
- thin-lenses, 936–942

- images from, 938, 938–942, 939, 947, 947–948
 - two-lens systems, 939–941, 940
 - thin-lens formulas, 936
 - Newtonian and Gaussian forms contrasted, 956
 - third-law force pair, 98
 - three-dimensional electron traps, 1095, 1095–1096
 - three-dimensional motion
 - acceleration, 61, 61
 - position and displacement, 58, 58–60
 - velocity, 60, 60–62, 61
 - three-slit diffraction gratings, 1019
 - thrust, 226
 - thunderstorm sprites, 585, 585
 - time
 - directional nature of, 536
 - relativity of, 1027, 1027–1031, 1037
 - scalar nature of, 38
 - units, 5–6
 - time constant
 - LC* oscillations, 826
 - RC* circuits, 721, 721
 - RL* circuits, 809
 - time dilation, 1028
 - consequences of Lorentz transformation equations, 1037
 - tests of, 1029
 - and transverse Doppler effect, 1041, 1041–1042
 - time intervals, 5, 5t
 - between relativistic events, 1027, 1027–1031
 - time signals, 6
 - tokamak, 1212
 - ton, 11
 - top quark, 1233t, 1234
 - toroids
 - magnetic field, 777, 777–778
 - torque, 258–260, 289t
 - and angular momentum of system of particles, 288
 - and conservation of angular momentum, 290
 - on current loop, 752, 752–753
 - and gyroscope precession, 294, 295
 - internal and external, 288
 - net (resultant), 259
 - Newton's second law in angular form, 285
 - particle about fixed point, 281, 281–282
 - rolling down ramp, 278–279
 - torr, 361
 - torsion constant, torsion pendulum, 394
 - torsion pendulum, 394, 394
 - total internal reflection, 911, 911–912
 - tour jeté, 291, 291
 - tracer, for following fluid flow, 371, 371
 - trajectory, in projectile motion, 67
 - transformers, 850–852
 - autotransformer, 859, 859
 - energy transmission requirements, 850
 - ideal, 850, 850–852
 - impedance matching, 852
 - in *LC* oscillators, 891, 891
 - transient current series *RLC* circuits, 844
 - transistors, 697, 1059
 - field-effect-transistor (FET), 1159, 1159–1160
 - MOSFET (metal-oxide-semiconductor-field-effect transistor), 1159, 1159
 - transition elements
 - paramagnetism, 875
 - translation, 241. *See also* one-dimensional motion; two-dimensional motion
 - in rolling, 275–277, 276
 - translational equilibrium, 306
 - translational kinetic energy
 - ideal gases, 513–514
 - of rolling, 277
 - yo-yo, 280–281
 - translational variables, 289t
 - transmission coefficient, barrier tunneling, 1075
 - transparent material, 906
 - transuranic nuclides, 1204
 - transverse Doppler effect, 1041, 1041–1042
 - transverse motion, 414
 - transverse waves, 413–414, 414
 - electromagnetic waves, 891
 - traveling electromagnetic waves, 890–896, 893, 894
 - traveling waves, 414, 1083
 - energy and power, 421, 421–423
 - sound, 448, 448–450
 - speed, 417, 417–419
 - wave function, 1071–1072
 - triangular prism, 908, 908
 - angle of minimum deviation, 920
 - trig functions, 43, 43
 - triple point cell, 478, 478
 - triple point of water, 478–479
 - tritium, 1211, 1212
 - and valley of the nuclides, 1182
 - triton, 1211
 - tube length, compound microscope, 944
 - tunnel diode, 1074
 - tunneling, barrier, 1074, 1074–1076
 - turbulent flow, 371
 - turning points, in potential energy curves, 176–178, 177
 - turns
 - in coils, 752–753
 - in solenoids, 776
 - turns ratio, transformers, 852
 - two-dimensional collisions, 224, 224
 - two-dimensional electron traps, 1095, 1095–1096
 - two-dimensional explosions, 216, 216
 - two-dimensional motion
 - acceleration, 62–64, 63
 - position and displacement, 58–60, 58, 59
 - relative, 74–75, 74
 - velocity, 60–62, 60, 61
- U**
- ultimate strength, 316, 316
 - selected engineering materials, 317t
 - ultrarelativistic proton, 1047
 - ultrasound
 - bat navigation using, 464
 - blood flow speed measurement using, 473
 - ultrasound imaging, 445, 445
 - ultraviolet light, 413
 - ultraviolet radiation, 869
 - uncertainty principle, 1072, 1073, 1236
 - uniform charge distributions
 - electric field lines, 581, 581–582
 - types of, 590
 - uniform circular motion, 70–72, 71
 - centripetal force in, 124–128, 125
 - and simple harmonic motion, 398–399, 399
 - uniform displacement current, 885
 - uniform displacement-current density, 885
 - uniform electric field, 582, 582, 884
 - uniform electric flux, 884
 - unit cell, 1012, 1012
 - metals, insulators, and semiconductors, 1142–1143, 1142
 - United States Naval Observatory time signals, 6
 - units, 1–2
 - changing, 3
 - length, 3–4
 - mass, 6–8
 - time, 5–6
 - unit vectors, 44, 44
 - universe
 - Big Bang, 1237, 1240–1243, 1241
 - color-coded image of universe at 379,000 yrs old, 1242, 1242
 - cosmic background radiation, 1239, 1241, 1243, 1243
 - critical density for continued expansion, 1246
 - dark energy, 1243
 - dark matter, 1240
 - estimated age, 1238
 - expansion, 1238, 1243, 1242–1243, 1246
 - temperature of early, 477
 - unoccupied levels, 1122, 1144, 1185
 - unpolarized light, 901, 901
 - unstable equilibrium, 178
 - potential energy curves, 178
 - unstable static equilibrium, 306
 - up quark, 578, 655, 1232, 1233t, 1233, 1234
 - uranium
 - density of nucleus, 360t
 - electric field at surface of nucleus, 580t
 - mass energy, 1043t
 - uranium²²⁸
 - alpha decay, 1177
 - half-life, 1178, 1179t
 - uranium²³⁵, 1215
 - enriching fuel, 1202, 1216
 - fission, 1196–1201, 1199, 1214
 - fissionability, 1200t, 1201, 1206
 - in natural nuclear reactor, 1206
 - percentage in uranium ore, 1216
 - spontaneous fission, 1214
 - uranium²³⁶, 1197, 1200t

uranium₂₃₈, 572, 1174
 abundance in granite, 1192
 alpha decay, 1178, 1177–1179
 binding energy per nucleon, 1172
 fissionability, 1200t, 1201, 1206
 half-life, 1179, 1179t, 1192
 uranium₂₃₉, 1200t
 uranium enrichment, 1202, 1216

V

vacant levels, 1144
 valence band, 1151, 1151
 valence electrons, 1083, 1126, 1145
 valence number, 1152
 valley of nuclides, 1182, 1182
 vaporization, 486
 vapor state, 486
 variable capacitor, 678
 variable force
 work done by general variable, 151–153, 152
 work done by spring force, 149, 149–151
 variable-mass systems, rockets, 224–226, 225
 variable resistors, 727
 vector angles, 41, 41
 degrees and radians, 43
 measuring, 43
 vector-capable calculator, 44
 vector components, 41, 41–43, 42
 addition, 44–46
 vector equation, 39
 vector fields, 580
 vector product, 48, 50, 49–52
 vector quantities, 14, 38
 vectors, 580
 adding by components, 44–46
 adding geometrically, 39, 39–40, 40
 coupled, 1114
 cross product, 49–52, 51
 defined, 38, 38–40, 39
 dot product, 48, 48
 and laws of physics, 47
 multiplication, 47–52, 48
 resolving, 41
 scalar product, 48–49
 unit, 44, 44
 vector product, 47, 50, 50–51
 vector sum (resultant), 39, 39–40
 velocity, 263t. *See also* acceleration; angular velocity; forces; kinetic energy
 graphical integration in motion analysis, 27, 27–28
 line of sight, 355
 and Newton's first law, 87–91
 and Newton's second law, 91–94
 one-dimensional motion, 14–18
 projectile motion, 64–70
 relative motion in one dimension, 73–74
 relative motion in two dimensions, 74–75
 relativity of, 1039, 1039
 rockets, 224–226
 sign of, 20
 simple harmonic motion, 388, 388, 389, 389
 two- and three-dimensional motion, 60–62, 61, 62
 uniform circular motion, 70–72, 71
 as vector quantity, 38
 velocity amplitude
 forced oscillations, 402, 402
 simple harmonic motion, 389
 velocity field, 606, 606
 velocity vectors, 38
 venturi meter, 383
 vertical motion, in projectile motion, 66–67, 67
 virtual focal point, 929, 929
 virtual images
 defined, 924
 spherical mirrors, 930
 spherical refracting surfaces, 934, 934
 thin lenses, 938, 938
 virtual photons, 1235
 viscous drag force, 371
 visible light, 413, 890
 ultimate speed, 1023
 volt, 630
 voltage. *See also* potential difference
 ac circuits, 842t
 transformers, 850–852
 voltage law, Kirchhoff's, 708
 volt-ampere, 695
 voltmeters, 720, 720
 volume
 and ideal gas law, 508–511
 as state property, 537
 work done by ideal gas at constant, 510
 volume charge density, 576, 578, 586, 586t
 volume expansion, 482–483
 volume flow rate, 373, 605–606
 volume flux, 605

volume probability density, 1103, 1103, 1105, 1105, 1106

W

water
 boiling and freezing points in Celsius and Fahrenheit, 480t
 bulk modulus, 317, 446
 density, 360t
 dielectric properties, 669t
 diffraction of waves in tank, 963, 963
 as electric dipole, 594, 594, 638
 electric field within, 580t
 heats of transformation, 486, 487t
 heavy, 1217
 index of refraction, 906t
 in microwave cooking, 595
 as moderator for nuclear reactors, 1202
 as polar dielectric, 671
 polarization of light by reflection, 912
 pure, as insulator, 563
 RMS speed at room temperature, 512t
 specific heats, 485t
 speed of sound in, 447, 447t
 tap water, as conductor, 563
 thermal properties, 482–483
 triple point, 478
 water waves, 413
 watt (W), 2, 155–156
 Watt, James, 155
 wave equation, 423–424
 wave forms, 414, 414
 wavefronts, 445, 445
 wave function, 1071–1072. *See also* Schrödinger's equation
 electron traps, 1088–1091, 1089
 hydrogen ground state, 1102–1103, 1103
 normalizing, 1089
 wave interference, 425, 425–428, 426
 sound waves, 451, 451–452
 wavelength, 415, 415
 and index of refraction, 960–962
 sound waves, 449
 wave on stretched string, 420
 wavelength Doppler shift, 1040
 waves. *See also* seismic waves; sound waves; standing waves; stretched strings; traveling waves
 phasors, 428–430, 429
 principle of superposition for, 425, 425
 transverse and longitudinal, 413–414, 414
 types of, 413
 wavelength and frequency, 414–417
 wave shape, 415
 wave speed, 417, 417–420
 sound waves, 449
 on stretched string, 420, 420–421
 wave trains, 1131
 weak force, 1221
 messenger particle, 1235
 weak interaction, 1225
 weber (unit), 793
 weight, 95–96
 apparent weight in fluids, 369–370
 weightlessness, 124
 well depth, finite well electron traps, 1091
 whiplash injury, 28
 white dwarfs, 341t
 density of core, 360t
 white light
 chromatic dispersion, 908, 908
 Wien's law, 1247
 Wilkinson Microwave Anisotropy Probe (WMAP), 1242
 windings, solenoid, 776
 window glass
 thermal conductivity, 494t
 Wintergreen LifeSaver, blue flashes, 564
 WMAP (Wilkinson Microwave Anisotropy Probe), 1242
 W messenger particle, 1236
 work, 263t
 Carnot engines, 545
 and conservation of mechanical energy, 173–175
 and conservation of total energy, 183–186, 184
 defined, 142–143
 done by electrostatic force, 630–631
 done by external force with friction, 180–182
 done by external force without friction, 180
 done by gravitational force, 146, 146–148

done by ideal gas at constant pressure, 510
done by ideal gas at constant temperature, 509, 509–510
done by ideal gas at constant volume, 510
done by spring force, 149, 149–151
done by variable force, 151–155, 152
done in lifting and lowering objects, 147, 147–148
done on system by external force, 180–182, 181
first law of thermodynamics, 491–493
and heat, 484, 488–490
and kinetic energy, 143, 142–146
net, 144

path-dependent quantity, 490
path independence of conservative forces, 168–170, 169
and potential energy, 167, 167–170
and power, 155–156, 156
and rotational kinetic energy, 262–264
signs for, 143
work, energy, and emf, 706–707
work function, 1060
working substance, 543
work-kinetic energy theorem, 144–146, 263t
variable force, 153–155

X

x component, of vectors, 41–42, 41, 42

xenon
decay chain, 1197
xi-minus particle, 1230, 1231t, 1235
x-ray diffraction, 1011–1013, 1012
x rays, 413, 889, 890
characteristic x-ray spectrum, 1128–1129, 1129
continuous x-ray spectrum, 1127, 1127–1128
and ordering of elements, 1127–1130
radiation dosage, 1184

Y

y component, of vectors, 41–42, 41
yield strength, 316, 316
selected engineering materials, 317t

Young's double-slit interference experiment, 965, 965–967
single-photon, wide-angle version, 1067, 1067
single-photon version, 1065, 1066
Young's modulus, 316
selected engineering materials, 317t
yo-yo, 280–281, 281

Z

zero angular position, 242
zero-point energy, 1090
zeroth law of thermodynamics, 477, 477
zeroth-order line, 1006
Z messenger particle, 1236

Some Physical Constants*

Speed of light	c	$2.998 \times 10^8 \text{ m/s}$
Gravitational constant	G	$6.673 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2$
Avogadro constant	N_A	$6.022 \times 10^{23} \text{ mol}^{-1}$
Universal gas constant	R	$8.314 \text{ J/mol} \cdot \text{K}$
Mass–energy relation	c^2	$8.988 \times 10^{16} \text{ J/kg}$ 931.49 MeV/u
Permittivity constant	ϵ_0	$8.854 \times 10^{-12} \text{ F/m}$
Permeability constant	μ_0	$1.257 \times 10^{-6} \text{ H/m}$
Planck constant	h	$6.626 \times 10^{-34} \text{ J} \cdot \text{s}$ $4.136 \times 10^{-15} \text{ eV} \cdot \text{s}$
Boltzmann constant	k	$1.381 \times 10^{-23} \text{ J/K}$ $8.617 \times 10^{-5} \text{ eV/K}$
Elementary charge	e	$1.602 \times 10^{-19} \text{ C}$
Electron mass	m_e	$9.109 \times 10^{-31} \text{ kg}$
Proton mass	m_p	$1.673 \times 10^{-27} \text{ kg}$
Neutron mass	m_n	$1.675 \times 10^{-27} \text{ kg}$
Deuteron mass	m_d	$3.344 \times 10^{-27} \text{ kg}$
Bohr radius	a	$5.292 \times 10^{-11} \text{ m}$
Bohr magneton	μ_B	$9.274 \times 10^{-24} \text{ J/T}$ $5.788 \times 10^{-5} \text{ eV/T}$
Rydberg constant	R	$1.097\,373 \times 10^7 \text{ m}^{-1}$

*For a more complete list, showing also the best experimental values, see Appendix B.

The Greek Alphabet

Alpha	A	α	Iota	I	ι	Rho	P	ρ
Beta	B	β	Kappa	K	κ	Sigma	Σ	σ
Gamma	Γ	γ	Lambda	Λ	λ	Tau	T	τ
Delta	Δ	δ	Mu	M	μ	Upsilon	Υ	υ
Epsilon	E	ϵ	Nu	N	ν	Phi	Φ	ϕ, φ
Zeta	Z	ζ	Xi	Ξ	ξ	Chi	X	χ
Eta	H	η	Omicron	O	o	Psi	Ψ	ψ
Theta	Θ	θ	Pi	Π	π	Omega	Ω	ω

Some Conversion Factors*

Mass and Density

$$1 \text{ kg} = 1000 \text{ g} = 6.02 \times 10^{26} \text{ u}$$

$$1 \text{ slug} = 14.59 \text{ kg}$$

$$1 \text{ u} = 1.661 \times 10^{-27} \text{ kg}$$

$$1 \text{ kg/m}^3 = 10^{-3} \text{ g/cm}^3$$

Length and Volume

$$1 \text{ m} = 100 \text{ cm} = 39.4 \text{ in.} = 3.28 \text{ ft}$$

$$1 \text{ mi} = 1.61 \text{ km} = 5280 \text{ ft}$$

$$1 \text{ in.} = 2.54 \text{ cm}$$

$$1 \text{ nm} = 10^{-9} \text{ m} = 10 \text{ \AA}$$

$$1 \text{ pm} = 10^{-12} \text{ m} = 1000 \text{ fm}$$

$$1 \text{ light-year} = 9.461 \times 10^{15} \text{ m}$$

$$1 \text{ m}^3 = 1000 \text{ L} = 35.3 \text{ ft}^3 = 264 \text{ gal}$$

Time

$$1 \text{ d} = 86\,400 \text{ s}$$

$$1 \text{ y} = 365\frac{1}{4} \text{ d} = 3.16 \times 10^7 \text{ s}$$

Angular Measure

$$1 \text{ rad} = 57.3^\circ = 0.159 \text{ rev}$$

$$\pi \text{ rad} = 180^\circ = \frac{1}{2} \text{ rev}$$

Speed

$$1 \text{ m/s} = 3.28 \text{ ft/s} = 2.24 \text{ mi/h}$$

$$1 \text{ km/h} = 0.621 \text{ mi/h} = 0.278 \text{ m/s}$$

Force and Pressure

$$1 \text{ N} = 10^5 \text{ dyne} = 0.225 \text{ lb}$$

$$1 \text{ lb} = 4.45 \text{ N}$$

$$1 \text{ ton} = 2000 \text{ lb}$$

$$1 \text{ Pa} = 1 \text{ N/m}^2 = 10 \text{ dyne/cm}^2$$

$$= 1.45 \times 10^{-4} \text{ lb/in.}^2$$

$$1 \text{ atm} = 1.01 \times 10^5 \text{ Pa} = 14.7 \text{ lb/in.}^2$$

$$= 76.0 \text{ cm Hg}$$

Energy and Power

$$1 \text{ J} = 10^7 \text{ erg} = 0.2389 \text{ cal} = 0.738 \text{ ft} \cdot \text{lb}$$

$$1 \text{ kW} \cdot \text{h} = 3.6 \times 10^6 \text{ J}$$

$$1 \text{ cal} = 4.1868 \text{ J}$$

$$1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$$

$$1 \text{ horsepower} = 746 \text{ W} = 550 \text{ ft} \cdot \text{lb/s}$$

Magnetism

$$1 \text{ T} = 1 \text{ Wb/m}^2 = 10^4 \text{ gauss}$$

*See Appendix D for a more complete list.

