


User: model10



Special Edition

16.1 Copyright 1985-2019 StataCorp LLC
 StataCorp
 4905 Lakeway Drive
 College Station, Texas 77845 USA
 800-STATA-PC <https://www.stata.com>
 979-696-4600 stata@stata.com
 979-696-4601 (fax)

Stata license: 50-student lab perpetual
 Serial number: 401606339725
 Licensed to: Melanie M
 WLU

Notes:

1. Unicode is supported; see [help unicode advice](#).
2. Maximum number of variables is set to 5,000; see [help set maxvar](#).

Running C:\Program Files\Stata16\sysprofile.do ...

- 1 . import delimited "C:\Users\fari9860\OneDrive - Wilfrid Laurier University\MA489\~all_cog_var_pivot_a70.csv"
 (23 vars, 58,228 obs)
- 2 . mixed score ageminus50 smoke hispanic diabetes blood_pressure psych_problems is_depressed doesnt_drink i.education_level
 > tdweights] || id: ageminus50

Obtaining starting values by EM:

Performing gradient-based optimization:

Iteration 0: log pseudolikelihood = **-117159.1**
 Iteration 1: log pseudolikelihood = **-117156.47**
 Iteration 2: log pseudolikelihood = **-117156.46**

Computing standard errors:

Mixed-effects regression Number of obs = **58,228**
 Group variable: id Number of groups = **14,674**

Obs per group:
 min = **1**
 avg = **4.0**
 max = **9**

Log pseudolikelihood = **-117156.46** Wald chi2(18) = **10638.01**
 Prob > chi2 = **0.0000**

(Std. Err. adjusted for **14,674** clusters in id)

score	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
ageminus50	-.2317705	.0039881	-58.12	0.000	-.239587	-.2239539
smoke	-.2666512	.090468	-2.95	0.003	-.4439652	-.0893372
hispanic	-.7413381	.1072475	-6.91	0.000	-.9515393	-.5311368
diabetes	-.3594406	.0593707	-6.05	0.000	-.4758049	-.2430762
blood_pressure	-.1620823	.0523176	-3.10	0.002	-.2646229	-.0595417
psych_problems	-.5636726	.072361	-7.79	0.000	-.7054976	-.4218477
is_depressed	-.3902103	.0419723	-9.30	0.000	-.4724745	-.3079461
doesnt_drink	-.4399314	.0449108	-9.80	0.000	-.527955	-.3519078
education_level						
1	3.065664	.0782805	39.16	0.000	2.912237	3.219091
2	4.423632	.0968586	45.67	0.000	4.233792	4.613471
3	4.992152	.1166552	42.79	0.000	4.763512	5.220792

exercise_freq						
1	-.2028438	.0489593	-4.14	0.000	-.2988023	-.1068853
2	-1.04968	.0604381	-17.37	0.000	-1.168136	-.9312234
heavy_drinker	-.1505617	.1324181	-1.14	0.256	-.4100964	.1089729
marital_end	-.0212114	.049197	-0.43	0.666	-.1176358	.0752131
trouble_falling	.0463211	.053542	0.87	0.387	-.0586193	.1512616
wake_up_early	.0616458	.0508713	1.21	0.226	-.0380602	.1613518
underweight	-.4964715	.1228194	-4.04	0.000	-.7371931	-.2557498
_cons	18.56586	.1444889	128.49	0.000	18.28267	18.84905

Random-effects Parameters	Estimate	Robust Std. Err.	[95% Conf. Interval]	
id: Independent				
var(ageminus50)	.004275	.0003142	.0037014	.0049373
var(_cons)	4.472007	.2506166	4.006823	4.991199
var(Residual)	7.216741	.0798097	7.062	7.374873

Warning: Sampling weights were specified only at the first level in a multilevel model. If these weights are indicative

3 . mixed ageminus50 smoke hispanic diabetes blood_pressure psych_problems is_depressed doesnt_drink i.education_level i.

Obtaining starting values by EM:

Performing gradient-based optimization:

```
Iteration 0: log pseudolikelihood = -91485.081
Iteration 1: log pseudolikelihood = 501791.16 (not concave)
Iteration 2: log pseudolikelihood = 523489.73 (not concave)
Iteration 3: log pseudolikelihood = 540131.77 (not concave)
Iteration 4: log pseudolikelihood = 563998.09 (not concave)
Iteration 5: log pseudolikelihood = 585174.24 (not concave)
Iteration 6: log pseudolikelihood = 600884.11 (not concave)
Iteration 7: log pseudolikelihood = 632578.15 (not concave)
Iteration 8: log pseudolikelihood = 659451.35 (not concave)
Iteration 9: log pseudolikelihood = 691913.7 (not concave)
Iteration 10: log pseudolikelihood = 716291.5 (not concave)
Iteration 11: log pseudolikelihood = 738293.41 (not concave)
Iteration 12: log pseudolikelihood = 755108.62 (not concave)
Iteration 13: log pseudolikelihood = 778293.33 (not concave)
Iteration 14: log pseudolikelihood = 797320.39 (not concave)
Iteration 15: log pseudolikelihood = 811839.8 (not concave)
Iteration 16: log pseudolikelihood = 859158.75 (not concave)
```

—Break—

r(1);

4 . mixed score ageminus50 smoke hispanic diabetes blood_pressure psych_problems is_depressed doesnt_drink i.education_level
variable exercise_frew not found

r(111);

5 . mixed score ageminus50 smoke hispanic diabetes blood_pressure psych_problems is_depressed doesnt_drink i.education_level

Obtaining starting values by EM:

Performing gradient-based optimization:

Iteration 0: log pseudolikelihood = **-117161.25**

Iteration 1: log pseudolikelihood = **-117158.62**

Iteration 2: log pseudolikelihood = **-117158.61**

Computing standard errors:

Mixed-effects regression Number of obs = **58,228**
Group variable: id Number of groups = **14,674**

Obs per group:
 min = **1**
 avg = **4.0**
 max = **9**

Log pseudolikelihood = **-117158.61** Wald chi2(14) = **10613.63**
 Prob > chi2 = **0.0000**

(Std. Err. adjusted for **14,674** clusters in id)

score	Coef.	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
ageminus50	-.2321048	.0038473	-60.33	0.000	-.2396454	-.2245642
smoke	-.273446	.0905138	-3.02	0.003	-.4508498	-.0960422
hispanic	-.7433141	.1071624	-6.94	0.000	-.9533485	-.5332797
diabetes	-.3585268	.0593755	-6.04	0.000	-.4749007	-.2421529
blood_pressure	-.1625679	.0523008	-3.11	0.002	-.2650756	-.0600602
psych_problems	-.5596959	.0720202	-7.77	0.000	-.7008528	-.418539
is_depressed	-.3825966	.0416832	-9.18	0.000	-.4642943	-.300899
doesnt_drink	-.4308994	.0445658	-9.67	0.000	-.5182468	-.343552
education_level						
1	3.063536	.0782666	39.14	0.000	2.910136	3.216935
2	4.42178	.0967141	45.72	0.000	4.232223	4.611336
3	4.989374	.1163727	42.87	0.000	4.761287	5.21746
exercise_freq						
1	-.2031734	.0489425	-4.15	0.000	-.2990991	-.1072478
2	-1.05042	.060436	-17.38	0.000	-1.168873	-.9319677
underweight	-.4956166	.1228784	-4.03	0.000	-.7364539	-.2547794
_cons	18.57185	.1439716	129.00	0.000	18.28967	18.85403

Random-effects Parameters	Estimate	Robust Std. Err.	[95% Conf. Interval]	
id: Independent				
var(ageminus50)	.0042749	.0003141	.0037015	.004937
var(_cons)	4.472133	.250458	4.007227	4.990977
var(Residual)	7.21762	.0797998	7.062898	7.375732

Warning: Sampling weights were specified only at the first level in a multilevel model. If these weights are indicative

```
6 . predict r, resid
7 . predict yhat, xb
8 . scatter r yhat
9 . graph export "C:\Users\fari9860\OneDrive - Wilfrid Laurier University\MA489\model10revised.jpg", as(jpg) name("Graph")
   (file C:\Users\fari9860\OneDrive - Wilfrid Laurier University\MA489\model10revised.jpg written in JPEG format)
10 . qnorm r
11 . graph export "C:\Users\fari9860\OneDrive - Wilfrid Laurier University\MA489\model10revisedqnorm.jpg", as(jpg) name("Gr")
   (file C:\Users\fari9860\OneDrive - Wilfrid Laurier University\MA489\model10revisedqnorm.jpg written in JPEG format)
12 .
```