

Lattice paths and Lagrangian matroids

Edward D. Kim, *University of California, Davis*

AMS Spring 2010 Western Section Meeting
Special Session on Geometric Combinatorics

April 18, 2010

Joint work with:

Anna Gundert, *ETH Zürich*

Daria Schymura, *Freie Universität Berlin*

The playbill

The cast

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

The playbill


The cast (in order of appearance???)

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Lattice path matroids

Fix an $m \times n$ grid.


- Let P be a North/East path from $(0, 0)$ to (m, n) .
- Let Q be a North/East path that stays above P .
- Let \mathcal{B} be the collection of all North/East paths B between P and Q . Then \mathcal{B} is the set of bases of a matroid.



Lattice path matroids

Fix an $m \times n$ grid.


- Let P be a North/East path from $(0, 0)$ to (m, n) .
- Let Q be a North/East path that stays above P .
- Let \mathcal{B} be the collection of all North/East paths B between P and Q . Then \mathcal{B} is the set of bases of a matroid.



Lattice path matroids

Fix an $m \times n$ grid.


- Let P be a North/East path from $(0, 0)$ to (m, n) .
- Let Q be a North/East path that stays above P .
- Let \mathcal{B} be the collection of all North/East paths B between P and Q . Then \mathcal{B} is the set of bases of a matroid.



Lattice path matroids

Fix an $m \times n$ grid.


- Let P be a North/East path from $(0, 0)$ to (m, n) .
- Let Q be a North/East path that stays above P .
- Let \mathcal{B} be the collection of all North/East paths B between P and Q . Then \mathcal{B} is the set of bases of a matroid.



Lattice path matroids

Fix an $m \times n$ grid.

- Let P be a North/East path from $(0, 0)$ to (m, n) .
- Let Q be a North/East path that stays above P .
- Let \mathcal{B} be the collection of all North/East paths B between P and Q . Then \mathcal{B} is the set of bases of a matroid.



Lagrangian matroids

$$[n]^* = \{1^*, \dots, n^*\}, S = [n] \cup [n]^*$$

Definition

- $B \subset S$ is a **transversal** of S if $|B \cap \{i, i^*\}| = 1$ for all $i \in [n]$.
- For transversals X and Y of S , we say i is a **divergence** if $i \in X \triangle Y$.

Definition

Let \mathcal{B} be a set of transversals of S . \mathcal{B} is the set of bases of a **Lagrangian matroid** if the following *symmetric exchange axiom* holds:
For all $X, Y \in \mathcal{B}$ and each divergence i , there is a divergence j such that $X \triangle \{i, i^*, j, j^*\} \in \mathcal{B}$.

Lagrangian matroids

$$[n]^* = \{1^*, \dots, n^*\}, S = [n] \cup [n]^*$$

Definition

- $B \subset S$ is a **transversal** of S if $|B \cap \{i, i^*\}| = 1$ for all $i \in [n]$.
- For transversals X and Y of S , we say i is a **divergence** if $i \in X \triangle Y$.

Definition

Let \mathcal{B} be a set of transversals of S . \mathcal{B} is the set of bases of a **Lagrangian matroid** if the following *symmetric exchange axiom* holds:
For all $X, Y \in \mathcal{B}$ and each divergence i , there is a divergence j such that $X \triangle \{i, i^*, j, j^*\} \in \mathcal{B}$.

Lagrangian matroids

$$[n]^* = \{1^*, \dots, n^*\}, S = [n] \cup [n]^*$$

Definition


- $B \subset S$ is a **transversal** of S if $|B \cap \{i, i^*\}| = 1$ for all $i \in [n]$.
- For transversals X and Y of S , we say i is a **divergence** if $i \in X \triangle Y$.

Definition

Let \mathcal{B} be a set of transversals of S . \mathcal{B} is the set of bases of a **Lagrangian matroid** if the following *symmetric exchange axiom* holds:
For all $X, Y \in \mathcal{B}$ and each divergence i , there is a divergence j such that $X \triangle \{i, i^*, j, j^*\} \in \mathcal{B}$.


Lattice path Lagrangian matroids (LPLM)

- Let P be an up-down path of length n starting at 0.
- Let Q be an up-down path starting at 0 that stays above P .
- Denote by $\mathcal{B}[P, Q]$ the set of all paths between P and Q .
- $\mathcal{B}[P, Q]$ is the set of bases of a Lagrangian matroid. (Say, for instance, that $i^* \in X$ if the path B goes up on the i th step.)




Lattice path Lagrangian matroids (LPLM)

- Let P be an up-down path of length n starting at 0.
- Let Q be an up-down path starting at 0 that stays above P .
- Denote by $\mathcal{B}[P, Q]$ the set of all paths between P and Q .
- $\mathcal{B}[P, Q]$ is the set of bases of a Lagrangian matroid. (Say, for instance, that $i^* \in X$ if the path B goes up on the i th step.)




Lattice path Lagrangian matroids (LPLM)

- Let P be an up-down path of length n starting at 0.
- Let Q be an up-down path starting at 0 that stays above P .
- Denote by $\mathcal{B}[P, Q]$ the set of all paths between P and Q .
- $\mathcal{B}[P, Q]$ is the set of bases of a Lagrangian matroid. (Say, for instance, that $i^* \in X$ if the path B goes up on the i th step.)



Lattice path Lagrangian matroids (LPLM)


- Let P be an up-down path of length n starting at 0.
- Let Q be an up-down path starting at 0 that stays above P .
- Denote by $\mathcal{B}[P, Q]$ the set of all paths between P and Q .
- $\mathcal{B}[P, Q]$ is the set of bases of a Lagrangian matroid. (Say, for instance, that $i^* \in X$ if the path B goes up on the i th step.)



an example path $B \in \mathcal{B}[P, Q]$

Lattice path Lagrangian matroids (LPLM)

- Let P be an up-down path of length n starting at 0.
- Let Q be an up-down path starting at 0 that stays above P .
- Denote by $\mathcal{B}[P, Q]$ the set of all paths between P and Q .
- $\mathcal{B}[P, Q]$ is the set of bases of a Lagrangian matroid. (Say, for instance, that $i^* \in X$ if the path B goes up on the i th step.)



an example path $B \in \mathcal{B}[P, Q]$

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Theorem

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a Lagrangian matroid $\Leftrightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a matroid on $T \subset S$.

The plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Theorem

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a Lagrangian matroid $\Leftrightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a matroid on $T \subset S$.

The plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Theorem

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a Lagrangian matroid $\Leftrightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a matroid on $T \subset S$.

This plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Conjecture: de Mier

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a **lattice path** Lagrangian matroid $\Leftrightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a **lattice path** matroid on $T \subset S$.

The plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Theorem: Gundert, K., Schymura

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a **lattice path** Lagrangian matroid $\Rightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a **lattice path** matroid on $T \subset S$.

The plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Outline

lattice path matroids \subset matroids,
lattice path Lagrangian matroids \subset Lagrangian matroids.

Question: de Mier

What is the relationship between lattice path Lagrangian matroids and lattice path matroids?

Theorem: Gundert, K., Schymura

Let \mathcal{B} be a collection of transversals of $S = [n] \cup [n]^*$. For a transversal T of S , define $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$.

Then, \mathcal{B} is a **lattice path** Lagrangian matroid $\Rightarrow \mathcal{I}_T^{\mathcal{B}}$ is the set of independent sets of a **lattice path** matroid on $T \subset S$.


The plan...

Given a LPLM $\mathcal{B} = \mathcal{B}[P, Q]$ show that (i) $\mathcal{I}_T^{\mathcal{B}}$ is a LPM and
(ii) construct it!

Bases of $\mathcal{I}_T^{\mathcal{B}}$

$\mathcal{B} = \mathcal{B}[P, Q]$ a LPLM, T a transversal.


What are the bases of $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$?



Paths $B \in \mathcal{B}[P, Q]$ that agree maximally with T !

We call these **maximal** paths.

Bases of $\mathcal{I}_T^{\mathcal{B}}$


 $\mathcal{B} = \mathcal{B}[P, Q]$ a LPLM, T a transversal.

We call these **maximal** paths.

Bases of $\mathcal{I}_T^{\mathcal{B}}$

$\mathcal{B} = \mathcal{B}[P, Q]$ a LPLM, T a transversal.

What are the bases of $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$?




Paths $B \in \mathcal{B}[P, Q]$ that agree maximally with T !

We call these **maximal** paths.

Bases of $\mathcal{I}_T^{\mathcal{B}}$

$\mathcal{B} = \mathcal{B}[P, Q]$ a LPLM, T a transversal.

What are the bases of $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$?




Paths $B \in \mathcal{B}[P, Q]$ that agree maximally with T !

We call these **maximal** paths.

Bases of $\mathcal{I}_T^{\mathcal{B}}$

$\mathcal{B} = \mathcal{B}[P, Q]$ a LPLM, T a transversal.

What are the bases of $\mathcal{I}_T^{\mathcal{B}} = \{I : I \subseteq B \cap T \text{ for some } B \in \mathcal{B}\}$?




Paths $B \in \mathcal{B}[P, Q]$ that agree maximally with T !

We call these **maximal** paths.

P^{\max} and Q^{\max}

If two maximal paths cross,...




So there is a lowest maximal path P^{\max} and a highest maximal path Q^{\max} .

We only have to consider paths between P^{\max} and Q^{\max} !

P^{\max} and Q^{\max}


If two maximal paths cross,...



So there is a lowest maximal path P^{\max} and a highest maximal path Q^{\max} .

We only have to consider paths between P^{\max} and Q^{\max} !

All maximal paths end at the same point.




Suppose S, R maximal paths with $h_S(n) > h_R(n)$.

Among the disagreeing steps there have to be more with $R(j) = \searrow, S(j) = \nearrow$.

R cannot agree with T in each of these steps.

$\Rightarrow W$ has one more agreement than R .

All maximal paths end at the same point.




Suppose S, R maximal paths with $h_S(n) > h_R(n)$.

Among the disagreeing steps there have to be more with $R(j) = \searrow, S(j) = \nearrow$.

R cannot agree with T in each of these steps.

$\Rightarrow W$ has one more agreement than R .

All maximal paths end at the same point.




Suppose S, R maximal paths with $h_S(n) > h_R(n)$.

Among the disagreeing steps there have to be more with $R(j) = \searrow, S(j) = \nearrow$.

R cannot agree with T in each of these steps.

$\Rightarrow W$ has one more agreement than R .

All maximal paths end at the same point.




Suppose S, R maximal paths with $h_S(n) > h_R(n)$.

Among the disagreeing steps there have to be more with $R(j) = \searrow, S(j) = \nearrow$.

R cannot agree with T in each of these steps.


$\Rightarrow W$ has one more agreement than R .

Suppose all maximal paths agree in step i ...



...then it looks like this (i.e., no gaps)
 $\implies i$ is a “parallel step”

Suppose NOT all maximal paths agree in step i ...



...then the maximal paths form a zig-zag-shape!
 $\implies i$ is a “zig-zag step”

Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet before the last step.
- There are no parallel steps.

Then:

- All paths between P^{\max} and Q^{\max} are maximal. (Only zig-zag steps!)
- T is constant. W.l.o.g. $T = \nearrow \nearrow \dots \nearrow$.

Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet before the last step.
- There are no parallel steps.

Then:

- All paths between P^{\max} and Q^{\max} are maximal. (Only zig-zag steps!)
- T is constant. W.l.o.g. $T = \nearrow \nearrow \dots \nearrow$.

Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet before the last step.
- There are no parallel steps.

Then:

- All paths between P^{\max} and Q^{\max} are maximal. (Only zig-zag steps!)
- T is constant. W.l.o.g. $T = \nearrow \nearrow \dots \nearrow$.


Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet before the last step.
- There are no parallel steps.

Then:

- **All** paths between P^{\max} and Q^{\max} are maximal. (Only zig-zag steps!)
- T is constant. W.l.o.g. $T = \nearrow \nearrow \dots \nearrow$.




Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet before the last step.
- There are no parallel steps.

Then:

- **All** paths between P^{\max} and Q^{\max} are maximal. (Only zig-zag steps!)
- T is constant. W.l.o.g. $T = \nearrow \nearrow \dots \nearrow$.




Proof of the theorem

We can assume:

- P^{\max} and Q^{\max} do not meet
- There are no parallel

Then:

- All paths between steps!)
- T is cons



Algorithm

The maximally-agreeing paths correspond to bases of a lattice path matroid.

- **TASK:** Compute the lattice path matroid explicitly.

Plan of attack

Given T , P and Q of length n , iteratively build the lattice path matroid.

- If $n > 1$, then we take the lattice path matroid corresponding to
 - $T(1, \dots, n-1)$
 - $P(1, \dots, n-1)$
 - $Q(1, \dots, n-1)$

and modify it.

Algorithm

The maximally-agreeing paths correspond to bases of a lattice path matroid.

- **TASK:** Compute the lattice path matroid explicitly.

Plan of attack

Given T , P and Q of length n , iteratively build the lattice path matroid.

- If $n > 1$, then we take the lattice path matroid corresponding to
 - $T(1, \dots, n-1)$
 - $P(1, \dots, n-1)$
 - $Q(1, \dots, n-1)$

and modify it.

Algorithm

The maximally-agreeing paths correspond to bases of a lattice path matroid.

- **TASK:** Compute the lattice path matroid explicitly.

Plan of attack

Given T , P and Q of length n , iteratively build the lattice path matroid.

- If $n > 1$, then we take the lattice path matroid corresponding to
 - $T(1, \dots, n-1)$
 - $P(1, \dots, n-1)$
 - $Q(1, \dots, n-1)$

and modify it.

Algorithm

The maximally-agreeing paths correspond to bases of a lattice path matroid.

- **TASK:** Compute the lattice path matroid explicitly.

Plan of attack

Given T , P and Q of length n , iteratively build the lattice path matroid.


- If $n > 1$, then we take the lattice path matroid corresponding to
 - $T(1, \dots, n-1)$
 - $P(1, \dots, n-1)$
 - $Q(1, \dots, n-1)$and modify it.

Can assume $T(n) = \nearrow$

We can assume, without loss of generality, that $T(n) = \nearrow$.

Can assume $T(n) = \nearrow$


We can assume, without loss of generality, that $T(n) = \nearrow$.



$T(n) = \searrow$, so “flip” everything.

Can assume $T(n) = \nearrow$

We can assume, without loss of generality, that $T(n) = \nearrow$.




$T(n) = \nearrow$, paths agree on same steps.

Rank remains the same

Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



In the easy case, all paths could extend

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Now, cannot extend to agree with T

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



All maximal paths will still agree with T nine times

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Build space of potential new paths (one under P^{\max})

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



All new maximal paths agree with T exactly 9 times

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



This is the area of **all** potential new paths

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



New lowest path P^{\max}

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



Steps that are not allowed

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



New segments that are allowed

Rank remains the same


Harder case: from the endpoint of all maximal paths for $n = 10$, can not agree with T on 11th step.



All maximal paths for $n = 11$

Rank remains the same

Modify the lattice path matroid




	10	
9		10
7	9	9
5	7	7
3	5	5
2	3	3
1	2	2
8	1	
6		
4		

Maximal Lagrangian lattice paths for $n = 10$

Rank remains the same

Modify the lattice path matroid




Identify space of new paths

	10	
9		10
7	9	9
5	7	7
3	5	5
2	3	3
1	2	2
8	1	
6		
4		

Rank remains the same

Modify the lattice path matroid




Add last step for old paths

	10	11
9		10
7	9	9
5	7	7
3	5	5
2	3	3
1	2	2
8	1	
6		
4		

Rank remains the same

Modify the lattice path matroid




Add new lowest path

	10	11	
9		10	11
7	9	9	10
5	7	7	9
3	5	5	7
2	3	3	5
1	2	2	3
8	1	2	
6			
4			

Rank remains the same

Modify the lattice path matroid




Add new legal inside steps

	10	11	
9		10	11
7	9	9	10
5	7	7	9
3	5	5	7
2	3	3	5
1	2	2	3
8	1	2	
6			
4			

Rank remains the same


Modify the lattice path matroid



Highlight: zone of new paths

	10	11	
9		10	11
7	9	9	10
5	7	7	9
3	5	5	7
2	3	3	5
1	2	2	3
8	1	2	
6			
4			

Modify the lattice path matroid



Final picture

	10	11	
9		10	11
7	9	9	10
5	7	7	9
3	5	5	7
2	3	3	5
1	2	2	3
8	1	2	
6			
4			

Thank you!

Thank you for your attention!

Special thanks to: Margaret Bayer, Andrew Berget, Joe Bonin, Jesús De Loera, Anna de Mier, Matthias Köppe, and Jay Schweig. Research supported in part by NSF grant DMS-0608785, NSF VIGRE grants DMS-0135345 and DMS-0636297, the Centre de Recerca Matemàtica at the Universitat Autònoma de Barcelona, and the Deutsche Forschungsgemeinschaft Research Training Group on Methods for Discrete Structures.