

Triangulations, Order Polytopes, and Generalized Snake Posets

Andrés R. Vindas Meléndez
(UC Berkeley)

AMS Special Session: Algebraic, Geometric and Topological
Combinatorics
University of Texas at El Paso
15-September-2022


Team



Matias von Bell
(Inst. of Geom., TU Graz)



Ben Braun
(Univ. of Kentucky)



Derek Hanely
(Penn State Berend)



Khrystyna Serhiyenko
(Univ. of Kentucky)



Julie Vega
(Maret School)



Andrés R. Vindas Meléndez
(UC Berkeley)



Martha Yip
(Univ. of Kentucky)

Order Polytopes

Order Polytopes

Let P be a partially ordered set on the set of elements $[d] := \{1, \dots, d\}$.

Let P be a partially ordered set on the set of elements $[d] := \{1, \dots, d\}$.


The **order polytope** is defined as

$$\mathcal{O}(P) = \left\{ \mathbf{x} = (x_1, \dots, x_d) \in [0, 1]^d : x_i \leq x_j \text{ for } i <_P j \right\}.$$

Order Polytopes: An Example


Order Polytopes: An Example

Let P be the diamond poset




Order Polytopes: An Example

Let P be the diamond poset



The six upper order ideals of P are




Order Polytopes: An Example

Order Polytopes: An Example


Then $\mathcal{O}(P) = \{(x_1, x_2, x_3, x_4) \in [0, 1]^4 : x_4 \leq x_2 \leq x_1 \text{ and } x_4 \leq x_3 \leq x_1\}$.

Order Polytopes: An Example

Then $\mathcal{O}(P) = \{(x_1, x_2, x_3, x_4) \in [0, 1]^4 : x_4 \leq x_2 \leq x_1 \text{ and } x_4 \leq x_3 \leq x_1\}$.
From the upper order ideals of P ,



we get that $\mathcal{O}(P)$ is the convex hull of the points $(0, 0, 0, 0)$, $(1, 0, 0, 0)$, $(1, 1, 0, 0)$, $(1, 0, 1, 0)$, $(1, 1, 1, 0)$ and $(1, 1, 1, 1)$.



- The dimension of $\mathcal{O}(P)$ is the number of elements of P .

- The dimension of $\mathcal{O}(P)$ is the number of elements of P .
- The vertices of $\mathcal{O}(P)$ correspond to the filters of P , i.e., the upper order ideals.

- The dimension of $\mathcal{O}(P)$ is the number of elements of P .
- The vertices of $\mathcal{O}(P)$ correspond to the filters of P , i.e., the upper order ideals.
- Volume of $\mathcal{O}(P)$ is the number of linear extensions of P .

Generalized Snake Posets

Generalized Snake Posets

Definition

For $n \in \mathbb{Z}_{\geq 0}$, a *generalized snake word* is a word of the form $w = w_0 w_1 \cdots w_n$ where $w_0 = \varepsilon$ is the empty letter and w_i is in the alphabet $\{L, R\}$ for $i = 1, \dots, n$. The *length* of the word is n , which is the number of letters in $\{L, R\}$.

Generalized Snake Posets

Definition

For $n \in \mathbb{Z}_{\geq 0}$, a *generalized snake word* is a word of the form $w = w_0 w_1 \cdots w_n$ where $w_0 = \varepsilon$ is the empty letter and w_i is in the alphabet $\{L, R\}$ for $i = 1, \dots, n$. The *length* of the word is n , which is the number of letters in $\{L, R\}$.

Definition

Given a generalized snake word w , the *generalized snake poset* $P(w)$ is defined recursively.

Generalized Snake Posets


Definition

For $n \in \mathbb{Z}_{\geq 0}$, a *generalized snake word* is a word of the form $w = w_0 w_1 \cdots w_n$ where $w_0 = \varepsilon$ is the empty letter and w_i is in the alphabet $\{L, R\}$ for $i = 1, \dots, n$. The *length* of the word is n , which is the number of letters in $\{L, R\}$.

Definition

Given a generalized snake word w , the *generalized snake poset* $P(w)$ is defined recursively.

The snake poset $S_5 = P(\varepsilon L R L R L)$ and the ladder poset $\mathcal{L}_5 = P(\varepsilon L L L L L)$.



Volume of the Order Polytope of Generalized Snake Posets

Volume of the Order Polytope of Generalized Snake Posets

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

For $n \geq 0$, let $w = w_0 w_1 \cdots w_n$ be a generalized snake word. If $k \geq 0$ is the largest index such that $w_k \neq w_n$, then the normalized volume v_n of $\mathcal{O}(P(w))$ is given recursively by

$$v_n = \text{Cat}(n - k + 1)v_k + (\text{Cat}(n - k + 2) - 2 \cdot \text{Cat}(n - k + 1))v_{k-1}$$

with $v_{-1} = 1$ and $v_0 = 2$.

Volume of the Order Polytope of Generalized Snake Posets

Volume of the Order Polytope of Generalized Snake Posets

Corollary (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

The normalized volume of $\mathcal{O}(S_n)$ with $n \geq 0$ is given recursively by

$$v_n = 2v_{n-1} + v_{n-2},$$

with $v_{-1} = 1$ and $v_0 = 2$. These are the Pell numbers.

Volume of the Order Polytope of Generalized Snake Posets

Corollary (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

The normalized volume of $\mathcal{O}(S_n)$ with $n \geq 0$ is given recursively by

$$v_n = 2v_{n-1} + v_{n-2},$$

with $v_{-1} = 1$ and $v_0 = 2$. These are the Pell numbers.

Corollary (Benedetti et al. 2019, Mészáros–Morales 2019)

The normalized volume of $\mathcal{O}(\mathcal{L}_n)$ with $n \geq 0$ is given by

$$v_n = \text{Cat}(n + 2).$$

Volume of the Order Polytope of Generalized Snake Posets

Corollary (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

The normalized volume of $\mathcal{O}(S_n)$ with $n \geq 0$ is given recursively by

$$v_n = 2v_{n-1} + v_{n-2},$$

with $v_{-1} = 1$ and $v_0 = 2$. These are the Pell numbers.

Corollary (Benedetti et al. 2019, Mészáros–Morales 2019)

The normalized volume of $\mathcal{O}(\mathcal{L}_n)$ with $n \geq 0$ is given by

$$v_n = \text{Cat}(n + 2).$$

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

For any generalized snake word $w = w_0 w_1 \cdots w_n$ of length n ,

$$\text{vol } \mathcal{O}(S_n) \leq \text{vol } \mathcal{O}(P(w)) \leq \text{vol } \mathcal{O}(\mathcal{L}_n).$$

Triangulations

Triangulations

Consider a polytope $\mathcal{P} \subseteq \mathbb{R}^d$. A **triangulation** \mathcal{T} of \mathcal{P} is a subdivision of \mathcal{P} into d -simplices.

Triangulations

Consider a polytope $\mathcal{P} \subseteq \mathbb{R}^d$. A **triangulation** \mathcal{T} of \mathcal{P} is a subdivision of \mathcal{P} into d -simplices.

- A triangulation is **unimodular** if every simplex has normalized volume one.

Triangulations

Consider a polytope $\mathcal{P} \subseteq \mathbb{R}^d$. A **triangulation** \mathcal{T} of \mathcal{P} is a subdivision of \mathcal{P} into d -simplices.

- A triangulation is **unimodular** if every simplex has normalized volume one.
- A triangulation of $\mathcal{P} \subseteq \mathbb{R}^d$ is **regular** if it can be obtained by projecting the lower envelope of a lifting of \mathcal{P} from \mathbb{R}^{d+1} .

Triangulations

Consider a polytope $\mathcal{P} \subseteq \mathbb{R}^d$. A **triangulation** \mathcal{T} of \mathcal{P} is a subdivision of \mathcal{P} into d -simplices.

- A triangulation is **unimodular** if every simplex has normalized volume one.
- A triangulation of $\mathcal{P} \subseteq \mathbb{R}^d$ is **regular** if it can be obtained by projecting the lower envelope of a lifting of \mathcal{P} from \mathbb{R}^{d+1} .



Figure: From “Existence of Unimodular Triangulations” by Haase et al.




Figure: From “Triangulations” by De Loera et al.

Triangulations of Order Polytopes

Triangulations of Order Polytopes

Define a hyperplane $\mathcal{H}_{i,j} = \{x \in \mathbb{R}^d : x_i = x_j\}$ for $1 \leq i < j \leq d$.

Triangulations of Order Polytopes

Define a hyperplane $\mathcal{H}_{i,j} = \{x \in \mathbb{R}^d : x_i = x_j\}$ for $1 \leq i < j \leq d$. The set of all such hyperplanes induces a triangulation \mathcal{T} of $\mathcal{O}(P)$ known as the *canonical triangulation*, which has the following three fundamental properties:

Triangulations of Order Polytopes

Define a hyperplane $\mathcal{H}_{i,j} = \{x \in \mathbb{R}^d : x_i = x_j\}$ for $1 \leq i < j \leq d$. The set of all such hyperplanes induces a triangulation \mathcal{T} of $\mathcal{O}(P)$ known as the *canonical triangulation*, which has the following three fundamental properties:

- 1 \mathcal{T} is unimodular,

Triangulations of Order Polytopes

Define a hyperplane $\mathcal{H}_{i,j} = \{x \in \mathbb{R}^d : x_i = x_j\}$ for $1 \leq i < j \leq d$. The set of all such hyperplanes induces a triangulation \mathcal{T} of $\mathcal{O}(P)$ known as the *canonical triangulation*, which has the following three fundamental properties:

- 1 \mathcal{T} is unimodular,
- 2 the maximal simplices are in bijection with the linear extensions of P , so the normalized volume of the order polytope is

$$\text{vol}(\mathcal{O}(P)) = \# \text{ of linear extensions of } P, \text{ and}$$

Triangulations of Order Polytopes

Define a hyperplane $\mathcal{H}_{i,j} = \{x \in \mathbb{R}^d : x_i = x_j\}$ for $1 \leq i < j \leq d$. The set of all such hyperplanes induces a triangulation \mathcal{T} of $\mathcal{O}(P)$ known as the *canonical triangulation*, which has the following three fundamental properties:

- 1 \mathcal{T} is unimodular,
- 2 the maximal simplices are in bijection with the linear extensions of P , so the normalized volume of the order polytope is

$$\text{vol}(\mathcal{O}(P)) = \# \text{ of linear extensions of } P, \text{ and}$$

- 3 the simplex corresponding to a linear extension (a_1, \dots, a_d) of P is

$$\sigma_{a_1, \dots, a_d} = \left\{ x \in [0, 1]^d : x_{a_1} \leq x_{a_2} \leq \dots \leq x_{a_d} \right\},$$

with vertex set $\{0, e_{a_d}, e_{a_{d-1}} + e_{a_d}, \dots, e_{a_1} + \dots + e_{a_d} = 1\}$.

Triangulations of Order Polytopes

Triangulations of Order Polytopes

What has been studied?

Triangulations of Order Polytopes

What has been studied?

- Graded posets (Reiner and Welker 2005)

Triangulations of Order Polytopes

What has been studied?

- Graded posets (Reiner and Welker 2005)
- Non-unimodular triangulations related to graph-associahedra (Féray and Reiner 2012)

Triangulations of Order Polytopes

What has been studied?

- Graded posets (Reiner and Welker 2005)
- Non-unimodular triangulations related to graph-associahedra (Féray and Reiner 2012)
- Product of chains (Santos, Stump, and Welker 2017)

Triangulations of Order Polytopes

What has been studied?

- Graded posets (Reiner and Welker 2005)
- Non-unimodular triangulations related to graph-associahedra (Féray and Reiner 2012)
- Product of chains (Santos, Stump, and Welker 2017)
- s-Lecture hall order polytopes (Bränden and Solus 2019)

Triangulations of Order Polytopes

What has been studied?

- Graded posets (Reiner and Welker 2005)
- Non-unimodular triangulations related to graph-associahedra (Féray and Reiner 2012)
- Product of chains (Santos, Stump, and Welker 2017)
- s-Lecture hall order polytopes (Bränden and Solus 2019)
- disjoint union of chains

Flip Graph & Secondary Polytope

Flip Graph & Secondary Polytope


- The **flip graph** of a polytope \mathcal{P} is a graph where each triangulation corresponds to a vertex in the graph, and a flip from a triangulation to another corresponds to an edge.

Flip Graph & Secondary Polytope

- The **flip graph** of a polytope \mathcal{P} is a graph where each triangulation corresponds to a vertex in the graph, and a flip from a triangulation to another corresponds to an edge.
- The **secondary polytope** of \mathcal{P} is a polytope of dimension $n - d - 1$ (n = number of vertices of \mathcal{P} and $d = \dim(\mathcal{P})$) whose vertices correspond to regular triangulations of \mathcal{P} . The 1-skeleton of the secondary polytope is the subgraph of the flip graph induced by the regular triangulations of \mathcal{P} .

Flip Graph & Secondary Polytope

- The **flip graph** of a polytope \mathcal{P} is a graph where each triangulation corresponds to a vertex in the graph, and a flip from a triangulation to another corresponds to an edge.
- The **secondary polytope** of \mathcal{P} is a polytope of dimension $n - d - 1$ (n = number of vertices of \mathcal{P} and $d = \dim(\mathcal{P})$) whose vertices correspond to regular triangulations of \mathcal{P} . The 1-skeleton of the secondary polytope is the subgraph of the flip graph induced by the regular triangulations of \mathcal{P} .



The Order Polytope of Meet-irreducibles

The Order Polytope of Meet-irreducibles

The Order Polytope of Meet-irreducibles

- Define $\hat{P}(w)$ to be the generalized snake poset $P(w)$ with $\hat{0}$ and $\hat{1}$ adjoined.

The Order Polytope of Meet-irreducibles

- Define $\hat{P}(w)$ to be the generalized snake poset $P(w)$ with $\hat{0}$ and $\hat{1}$ adjoined.
- Let $Q_w = \text{Irr}_{\wedge}(\hat{P})$ denote the poset of meet-irreducibles of \hat{P} .

The Order Polytope of Meet-irreducibles

- Define $\hat{P}(w)$ to be the generalized snake poset $P(w)$ with $\hat{0}$ and $\hat{1}$ adjoined.
- Let $Q_w = \text{Irr}_{\wedge}(\hat{P})$ denote the poset of meet-irreducibles of \hat{P} .
- By the fundamental theorem of finite distributive lattices, $\hat{P} \cong J(Q_w)$, where $J(Q_w)$ is the lattice of filters of Q_w , ordered by reverse inclusion.

The Order Polytope of Meet-irreducibles

- Define $\hat{P}(w)$ to be the generalized snake poset $P(w)$ with $\hat{0}$ and $\hat{1}$ adjoined.
- Let $Q_w = \text{Irr}_{\wedge}(\hat{P})$ denote the poset of meet-irreducibles of \hat{P} .
- By the fundamental theorem of finite distributive lattices, $\hat{P} \cong J(Q_w)$, where $J(Q_w)$ is the lattice of filters of Q_w , ordered by reverse inclusion.




Figure: The lattice $\hat{P}(w)$ for $w = \varepsilon L^3 R^2 L^4 R^5 L^2$ (left) and its poset of meet-irreducibles $Q_w = \text{Irr}_{\wedge}(\hat{P})$.

The Order Polytope of Meet-irreducibles

The Order Polytope of Meet-irreducibles

Let \mathcal{V} denote the subset of words which do not contain the substring LRL or RLR .

The Order Polytope of Meet-irreducibles

Let \mathcal{V} denote the subset of words which do not contain the substring LRL or RLR .

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

For $w \in \mathcal{V}$, every vertex of the secondary polytope of $\mathcal{O}(Q_w)$ is a unimodular triangulation. Thus, every triangulation of $\mathcal{O}(Q_w)$ is unimodular.

The Order Polytope of Meet-irreducibles

Let \mathcal{V} denote the subset of words which do not contain the substring LRL or RLR .

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

For $w \in \mathcal{V}$, every vertex of the secondary polytope of $\mathcal{O}(Q_w)$ is a unimodular triangulation. Thus, every triangulation of $\mathcal{O}(Q_w)$ is unimodular.

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ have length k . The canonical triangulation of $\mathcal{O}(Q_w)$ admits exactly $k + 1$ flips.

The Order Polytope of Meet-irreducibles

The Order Polytope of Meet-irreducibles

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w = \varepsilon L^{n-1}$, and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The flip graph of triangulations of $\mathcal{O}(Q_w)$ is the Cayley graph of the symmetric group \mathfrak{S}_{n+1} with the simple transpositions as the generating set.

Definition

Given a ladder \mathcal{L}^i , define $\tau_i \in \mathfrak{S}_{|V_0|}$ to be the permutation of V_0 such that for $v \in V_0$,



$$\tau_i(v) = \begin{cases} x_{j-1}, & \text{if } v = x_j \text{ and } j \in [s] \text{ is even,} \\ x_{j+1}, & \text{if } v = x_j \text{ and } j \in [s] \text{ is odd,} \\ v, & \text{otherwise.} \end{cases}$$

Twists

Definition


Given a ladder \mathcal{L}^i , define $\tau_i \in \mathfrak{S}_{|V_0|}$ to be the permutation of V_0 such that for $v \in V_0$,

$$\tau_i(v) = \begin{cases} x_{j-1}, & \text{if } v = x_j \text{ and } j \in [s] \text{ is even,} \\ x_{j+1}, & \text{if } v = x_j \text{ and } j \in [s] \text{ is odd,} \\ v, & \text{otherwise.} \end{cases}$$



\mathcal{L}^i in \hat{P} containing boxes with labels w_p, \dots, w_q , where $w_p < w_{p+1} < \dots < w_q$. The left (right) represents the case where $w_q = L$ ($w_q = R$).

Twists



Definition

Let $\mathfrak{T}(w)$ denote the subgroup of $\mathfrak{S}_{|V_0|}$ generated by the set of the τ_i 's. We call $\mathfrak{T}(w)$ *the twist group of $\hat{P}(w)$* . Elements of $\mathfrak{T}(w)$ are called *twists* and the elements τ_i are called *elementary twists*.

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$, $Q_w = \text{Irr}_\wedge(\hat{P}(w))$, and let \mathcal{T} and $\tau(\mathcal{T})$ be two triangulations of $\mathcal{O}(Q_w)$ where τ is a twist. If $\mathcal{T} = \mathcal{T}_Z^+$ can be flipped at circuit Z and $\tau(\mathcal{T}_Z^+) = \tau(\mathcal{T}_Z^+)^+_{\tau(Z)}$, then $\tau(\mathcal{T}_Z^+)^-_{\tau(Z)} = \tau(\mathcal{T}_Z^-)$. In other words, the following diagram commutes.

$$\begin{array}{ccc}
 \mathcal{T}_Z^+ & \xrightarrow{\text{flip in } Z} & \mathcal{T}_Z^- \\
 \downarrow \text{twist} & & \downarrow \text{twist} \\
 \tau(\mathcal{T}_Z^+) = \tau(\mathcal{T}_Z^+)^+_{\tau(Z)} & \xrightarrow{\text{flip in } \tau(Z)} & \tau(\mathcal{T}_Z^+)^-_{\tau(Z)} = \tau(\mathcal{T}_Z^-)
 \end{array}$$

Twists

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$, $Q_w = \text{Irr}_\wedge(\hat{P}(w))$, and let \mathcal{T} and $\tau(\mathcal{T})$ be two triangulations of $\mathcal{O}(Q_w)$ where τ is a twist. If $\mathcal{T} = \mathcal{T}_Z^+$ can be flipped at circuit Z and $\tau(\mathcal{T}_Z^+) = \tau(\mathcal{T}_Z^+)^+_{\tau(Z)}$, then $\tau(\mathcal{T}_Z^+)^-_{\tau(Z)} = \tau(\mathcal{T}_Z^-)$. In other words, the following diagram commutes.

$$\begin{array}{ccc} \mathcal{T}_Z^+ & \xrightarrow{\text{flip in } Z} & \mathcal{T}_Z^- \\ \downarrow \text{twist} & & \downarrow \text{twist} \\ \tau(\mathcal{T}_Z^+) = \tau(\mathcal{T}_Z^+)^+_{\tau(Z)} & \xrightarrow{\text{flip in } \tau(Z)} & \tau(\mathcal{T}_Z^+)^-_{\tau(Z)} = \tau(\mathcal{T}_Z^-) \end{array}$$

Corollary (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$, $Q_w = \text{Irr}_\wedge(\hat{P}(w))$, and \mathcal{T} & $\tau(\mathcal{T})$ be two triangulations of $\mathcal{O}(Q_w)$. Then \mathcal{T} and $\tau(\mathcal{T})$ admit the same number of flips.

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Proof Idea:

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Proof Idea:

- Hasse diagram of Q_w is strongly planar, by work of Mészáros, Morales, and Striker, $\mathcal{O}(Q_w)$ is int. equiv. to a flow polytope \mathcal{F}_{G_Q} .

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Proof Idea:

- Hasse diagram of Q_w is strongly planar, by work of Mészáros, Morales, and Striker, $\mathcal{O}(Q_w)$ is int. equiv. to a flow polytope \mathcal{F}_{G_Q} .
- The canonical triangulation of $\mathcal{O}(Q_w)$ maps to Danilov-Karzonov-Koshevoy triangulations of \mathcal{F}_{G_Q} .

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Proof Idea:

- Hasse diagram of Q_w is strongly planar, by work of Mészáros, Morales, and Striker, $\mathcal{O}(Q_w)$ is int. equiv. to a flow polytope \mathcal{F}_{G_Q} .
- The canonical triangulation of $\mathcal{O}(Q_w)$ maps to Danilov-Karzonov-Koshevoy triangulations of \mathcal{F}_{G_Q} .
- DKK triangulations are regular \rightarrow canonical triangulation of $\mathcal{O}(Q_w)$ are regular.

Theorem (MB-BB-DH-KS-JV-ARVM-MY, 2022+)

Let $w \in \mathcal{V}$ and $Q_w = \text{Irr}_\wedge(\hat{P}(w))$. The canonical triangulation \mathcal{T}_w of $\mathcal{O}(Q_w)$ is a regular triangulation, and for any twist τ , $\tau(\mathcal{T}_w)$ is also a regular triangulation.

Proof Idea:

- Hasse diagram of Q_w is strongly planar, by work of Mészáros, Morales, and Striker, $\mathcal{O}(Q_w)$ is int. equiv. to a flow polytope \mathcal{F}_{G_Q} .
- The canonical triangulation of $\mathcal{O}(Q_w)$ maps to Danilov-Karzonov-Koshevoy triangulations of \mathcal{F}_{G_Q} .
- DKK triangulations are regular \rightarrow canonical triangulation of $\mathcal{O}(Q_w)$ are regular.
- The twist group $\mathfrak{T}(w)$ acts on the canonical triangulation of $\mathcal{O}(Q_w)$.
- Any twist τ , $\tau(\mathcal{T}_w)$ corresponds to a framed triangulation of $\mathcal{F}_{G_{Q_w}}$, by DKK we know are regular.

Conjectures

- (i) For $w \in \mathcal{V}$, the flip graph of regular triangulations for $\mathcal{O}(Q_w)$ is k -regular, where k is the dimension of the secondary polytope of $\mathcal{O}(Q_w)$.

- (i) For $w \in \mathcal{V}$, the flip graph of regular triangulations for $\mathcal{O}(Q_w)$ is k -regular, where k is the dimension of the secondary polytope of $\mathcal{O}(Q_w)$.
- (ii) If $w \in \mathcal{V}$, all triangulations of $\mathcal{O}(Q_w)$ are regular.

- (i) For $w \in \mathcal{V}$, the flip graph of regular triangulations for $\mathcal{O}(Q_w)$ is k -regular, where k is the dimension of the secondary polytope of $\mathcal{O}(Q_w)$.
- (ii) If $w \in \mathcal{V}$, all triangulations of $\mathcal{O}(Q_w)$ are regular.
- (iii) The number of regular triangulations of $\mathcal{O}(S_n)$ is $2^{n+1} \cdot \text{Cat}(2n+1)$.

- (i) For $w \in \mathcal{V}$, the flip graph of regular triangulations for $\mathcal{O}(Q_w)$ is k -regular, where k is the dimension of the secondary polytope of $\mathcal{O}(Q_w)$.
- (ii) If $w \in \mathcal{V}$, all triangulations of $\mathcal{O}(Q_w)$ are regular.
- (iii) The number of regular triangulations of $\mathcal{O}(S_n)$ is $2^{n+1} \cdot \text{Cat}(2n+1)$.

The End



¡Gracias!