

Higher integrality conditions and volumes of slices

Fu Liu

University of California, Davis

AMS sectional meeting, Albuquerque, NM

April 17, 2010

Basic definitions

A *(convex) polytope* P in the d -dimensional Euclidean space \mathbb{R}^d is the convex hull of finitely many points $V = \{v_1, v_2, \dots, v_n\} \subset \mathbb{R}^d$. In other words,

$$P = \text{conv}(V) = \{\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n : \text{all } \lambda_i \geq 0, \text{ and } \lambda_1 + \lambda_2 + \dots + \lambda_n = 1\}.$$

Basic definitions

A *(convex) polytope* P in the d -dimensional Euclidean space \mathbb{R}^d is the convex hull of finitely many points $V = \{v_1, v_2, \dots, v_n\} \subset \mathbb{R}^d$. In other words,

$$P = \text{conv}(V) = \{\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n : \text{all } \lambda_i \geq 0, \text{ and } \lambda_1 + \lambda_2 + \dots + \lambda_n = 1\}.$$

Throughout this talk, we assume P is *full-dimensional*, i.e., $\dim(P) = d$.

Basic definitions

A *(convex) polytope* P in the d -dimensional Euclidean space \mathbb{R}^d is the convex hull of finitely many points $V = \{v_1, v_2, \dots, v_n\} \subset \mathbb{R}^d$. In other words,

$$P = \text{conv}(V) = \{\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n : \text{all } \lambda_i \geq 0, \text{ and } \lambda_1 + \lambda_2 + \dots + \lambda_n = 1\}.$$

Throughout this talk, we assume P is *full-dimensional*, i.e., $\dim(P) = d$.

The d -dimensional *lattice* $\mathbb{Z}^d \subset \mathbb{R}^d$ is the collection of all points with integer coordinates. Any point in the lattice is called a *lattice point* or an *integral point*.

Basic definitions

A *(convex) polytope* P in the d -dimensional Euclidean space \mathbb{R}^d is the convex hull of finitely many points $V = \{v_1, v_2, \dots, v_n\} \subset \mathbb{R}^d$. In other words,

$$P = \text{conv}(V) = \{\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n : \text{all } \lambda_i \geq 0, \text{ and } \lambda_1 + \lambda_2 + \dots + \lambda_n = 1\}.$$

Throughout this talk, we assume P is *full-dimensional*, i.e., $\dim(P) = d$.

The d -dimensional *lattice* $\mathbb{Z}^d \subset \mathbb{R}^d$ is the collection of all points with integer coordinates. Any point in the lattice is called a *lattice point* or an *integral point*.

A polytope P is *integral* if its vertices are all lattice points.

Basic definitions

A *(convex) polytope* P in the d -dimensional Euclidean space \mathbb{R}^d is the convex hull of finitely many points $V = \{v_1, v_2, \dots, v_n\} \subset \mathbb{R}^d$. In other words,

$$P = \text{conv}(V) = \{\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n : \text{all } \lambda_i \geq 0, \text{ and } \lambda_1 + \lambda_2 + \dots + \lambda_n = 1\}.$$

Throughout this talk, we assume P is *full-dimensional*, i.e., $\dim(P) = d$.

The d -dimensional *lattice* $\mathbb{Z}^d \subset \mathbb{R}^d$ is the collection of all points with integer coordinates. Any point in the lattice is called a *lattice point* or an *integral point*.

A polytope P is *integral* if its vertices are all lattice points.

For any $0 \leq \ell \leq d$, let $\pi^{(\ell)} : \mathbb{R}^d \rightarrow \mathbb{R}^{d-\ell}$ be the projection which drops the last ℓ coordinates.

Lattice points of a polytope

Definition 1. For any polytope $P \subset \mathbb{R}^d$ and positive integer $m \in \mathbb{N}$, the *m th dilated polytope* of P is $mP = \{m\mathbf{x} : \mathbf{x} \in P\}$. We denote by

$$i(P, m) = |mP \cap \mathbb{Z}^d|$$

the number of lattice points in mP .


Lattice points of a polytope

Definition 1. For any polytope $P \subset \mathbb{R}^d$ and positive integer $m \in \mathbb{N}$, the *m th dilated polytope* of P is $mP = \{m\mathbf{x} : \mathbf{x} \in P\}$. We denote by

$$i(P, m) = |mP \cap \mathbb{Z}^d|$$

the number of lattice points in mP .

Example: For any d , P be the convex hull of the set $\{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : x_i = 0 \text{ or } 1\}$, i.e. P is the *unit cube* in \mathbb{R}^d . Then $i(P, m) = (m + 1)^d$.



$$i(P, 3) = (3 + 1)^2$$

Theorem of Ehrhart (on integral polytopes)

Theorem 2 (Ehrhart). *Let P be a d -dimensional integral polytope. Then $i(P, m)$ is a polynomial in m of degree d .*

Therefore, we call $i(P, m)$ the *Ehrhart polynomial* of P .

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

▮► The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

- ▮ The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .
- ▮ The second coefficient equals $1/2$ of the sum of the normalized volumes of each facet.

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

- ➡ The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .
- ➡ The second coefficient equals $1/2$ of the sum of the normalized volumes of each facet.
- ➡ The constant term of $i(P, m)$ is always 1.

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

- ➡ The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .
- ➡ The second coefficient equals $1/2$ of the sum of the normalized volumes of each facet.
- ➡ The constant term of $i(P, m)$ is always 1.
- ➡ No simple forms known for other coefficients for general polytopes.

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

- ▮▮▮▮ The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .
- ▮▮▮▮ The second coefficient equals $1/2$ of the sum of the normalized volumes of each facet.
- ▮▮▮▮ The constant term of $i(P, m)$ is always 1.
- ▮▮▮▮ No simple forms known for other coefficients for general polytopes.
 - It is **NOT** even true that all the coefficients are nonnegative.

Coefficients of Ehrhart polynomials

If P is an integral polytope, what can we say about the coefficients of its Ehrhart polynomial $i(P, m)$?

- ▢▢▢▢ The leading coefficient of $i(P, m)$ is the volume $\text{Vol}(P)$ of P .
- ▢▢▢▢ The second coefficient equals $1/2$ of the sum of the normalized volumes of each facet.
- ▢▢▢▢ The constant term of $i(P, m)$ is always 1.
- ▢▢▢▢ No simple forms known for other coefficients for general polytopes.
 - It is **NOT** even true that all the coefficients are nonnegative.

Question: When are the coefficients positive? In particular, when can they be described by volumes?

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Example: A 3-dimensional integral cyclic polytope:

$$\Rightarrow P = \text{conv}\{(1, 1, 1), (2, 4, 8), (3, 9, 27), (4, 16, 64)\} : \text{Vol}(P) = 2.$$

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Example: A 3-dimensional integral cyclic polytope:

$$\Rightarrow P = \text{conv}\{(1, 1, 1), (2, 4, 8), (3, 9, 27), (4, 16, 64)\} : \text{Vol}(P) = 2.$$

$$\Rightarrow \pi^{(1)}(P) = \text{conv}\{(1, 1), (2, 4), (3, 9), (4, 16)\} : \text{Vol}(\pi^{(1)}(P)) = 4.$$

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Example: A 3-dimensional integral cyclic polytope:

$$\Rightarrow P = \text{conv}\{(1, 1, 1), (2, 4, 8), (3, 9, 27), (4, 16, 64)\} : \text{Vol}(P) = 2.$$

$$\Rightarrow \pi^{(1)}(P) = \text{conv}\{(1, 1), (2, 4), (3, 9), (4, 16)\} : \text{Vol}(\pi^{(1)}(P)) = 4.$$

$$\Rightarrow \pi^{(2)}(P) = \text{conv}\{1, 2, 3, 4\} = [1, 4] : \text{Vol}(\pi^{(2)}(P)) = 3.$$

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Example: A 3-dimensional integral cyclic polytope:

$$\Rightarrow P = \text{conv}\{(1, 1, 1), (2, 4, 8), (3, 9, 27), (4, 16, 64)\} : \text{Vol}(P) = 2.$$

$$\Rightarrow \pi^{(1)}(P) = \text{conv}\{(1, 1), (2, 4), (3, 9), (4, 16)\} : \text{Vol}(\pi^{(1)}(P)) = 4.$$

$$\Rightarrow \pi^{(2)}(P) = \text{conv}\{1, 2, 3, 4\} = [1, 4] : \text{Vol}(\pi^{(2)}(P)) = 3.$$

$$\Rightarrow \pi^{(3)}(P) = \mathbb{R}^0 : \text{Vol}(\pi^{(3)}(P)) = 1.$$

Ehrhart polynomials of integral cyclic polytopes

Theorem 3. *Suppose P is a d -dimensional integral cyclic polytope. Then*

$$i(P, m) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Example: A 3-dimensional integral cyclic polytope:

$$\Rightarrow P = \text{conv}\{(1, 1, 1), (2, 4, 8), (3, 9, 27), (4, 16, 64)\} : \text{Vol}(P) = 2.$$

$$\Rightarrow \pi^{(1)}(P) = \text{conv}\{(1, 1), (2, 4), (3, 9), (4, 16)\} : \text{Vol}(\pi^{(1)}(P)) = 4.$$

$$\Rightarrow \pi^{(2)}(P) = \text{conv}\{1, 2, 3, 4\} = [1, 4] : \text{Vol}(\pi^{(2)}(P)) = 3.$$

$$\Rightarrow \pi^{(3)}(P) = \mathbb{R}^0 : \text{Vol}(\pi^{(3)}(P)) = 1.$$

\Rightarrow The theorem says:

$$i(P, m) = 2m^3 + 4m^2 + 3m + 1.$$

Higher integrality conditions for affine spaces

Definition 4. An ℓ -dimensional affine space $U \subset \mathbb{R}^d$ is *integral* if

$$\pi^{(d-\ell)}(U \cap \mathbb{Z}^d) = \mathbb{Z}^\ell.$$

Or equivalently, the projection $\pi^{(d-\ell)}$ induces a bijection between $U \cap \mathbb{Z}^d$ and \mathbb{Z}^ℓ .

Higher integrality conditions for affine spaces

Definition 4. An ℓ -dimensional affine space $U \subset \mathbb{R}^d$ is *integral* if

$$\pi^{(d-\ell)}(U \cap \mathbb{Z}^d) = \mathbb{Z}^\ell.$$

Or equivalently, the projection $\pi^{(d-\ell)}$ induces a bijection between $U \cap \mathbb{Z}^d$ and \mathbb{Z}^ℓ .

Example:

(i) $\ell = 0$: U is integral if and only if U is a lattice point.

Higher integrality conditions for affine spaces


Definition 4. An ℓ -dimensional affine space $U \subset \mathbb{R}^d$ is *integral* if

$$\pi^{(d-\ell)}(U \cap \mathbb{Z}^d) = \mathbb{Z}^\ell.$$

Or equivalently, the projection $\pi^{(d-\ell)}$ induces a bijection between $U \cap \mathbb{Z}^d$ and \mathbb{Z}^ℓ .

Example:

- (i) $\ell = 0$: U is integral if and only if U is a lattice point.
- (ii) $\ell = 1$: In \mathbb{R}^2 ,



Higher integrality conditions for affine spaces

Definition 4. An ℓ -dimensional affine space $U \subset \mathbb{R}^d$ is *integral* if


$$\pi^{(d-\ell)}(U \cap \mathbb{Z}^d) = \mathbb{Z}^\ell.$$

Or equivalently, the projection $\pi^{(d-\ell)}$ induces a bijection between $U \cap \mathbb{Z}^d$ and \mathbb{Z}^ℓ .

Example:

(i) $\ell = 0$: U is integral if and only if U is a lattice point.

(ii) $\ell = 1$: In \mathbb{R}^2 ,



In general, U is integral if and only if U contains a lattice point and $\text{dir}(U) = (1, z_2, \dots, z_d) \in \mathbb{Z}^d$.

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Example:

(i) $k = 0$: P is 0-integral if and only if P is integral.

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Example:

- (i) $k = 0$: P is 0-integral if and only if P is integral.
- (ii) $k = 1$: P is 1-integral if and only if P is integral and $\text{dir}(e) = (1, z_2, \dots, z_d) \in \mathbb{Z}^d$ for any edge e of P .

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Example:

- (i) $k = 0$: P is 0-integral if and only if P is integral.
- (ii) $k = 1$: P is 1-integral if and only if P is integral and $\text{dir}(e) = (1, z_2, \dots, z_d) \in \mathbb{Z}^d$ for any edge e of P .

Theorem 6 (Partial statement). *If P is k -integral, then for $0 \leq \ell \leq k$, the coefficient of m^ℓ in $i(P, m)$ is $\text{Vol}(\pi^{d-\ell}(P))$.*

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Example:

- (i) $k = 0$: P is 0-integral if and only if P is integral.
- (ii) $k = 1$: P is 1-integral if and only if P is integral and $\text{dir}(e) = (1, z_2, \dots, z_d) \in \mathbb{Z}^d$ for any edge e of P .

Theorem 6 (Partial statement). *If P is k -integral, then for $0 \leq \ell \leq k$, the coefficient of m^ℓ in $i(P, m)$ is $\text{Vol}(\pi^{d-\ell}(P))$.*

Remark 7. If $k = 0$, the theorem says that the constant term is 1 for integral polytopes.

If $k = d$, the theorem says that all the coefficients are volumes of projections.

Higher integrality conditions for polytopes

Definition 5. A polytope P is *k -integral* if for any $0 \leq \ell \leq k$, we have that $\text{aff}(F)$ is integral for any ℓ -dimensional face F of P .

We say P is *fully integral* if $k = d$.

Example:

- (i) $k = 0$: P is 0-integral if and only if P is integral.
- (ii) $k = 1$: P is 1-integral if and only if P is integral and $\text{dir}(e) = (1, z_2, \dots, z_d) \in \mathbb{Z}^d$ for any edge e of P .

Theorem 6 (Partial statement). *If P is k -integral, then for $0 \leq \ell \leq k$, the coefficient of m^ℓ in $i(P, m)$ is $\text{Vol}(\pi^{d-\ell}(P))$.*

Remark 7. If $k = 0$, the theorem says that the constant term is 1 for integral polytopes. If $k = d$, the theorem says that all the coefficients are volumes of projections.

Fact: An integral cyclic polytope is fully integral.

An example of 1-integral polytope

Example: $P = \text{conv}\{(0, 0, 0), (4, 0, 0), (3, 6, 0), (2, 2, 2)\}$. One can check that P is 1-integral.

$$i(P, m) = 8m^3 + 10m^2 + 4m + 1,$$

and $\text{Vol}(\pi^{(d-1)}(P)) = \text{Vol}([0, 4]) = 4$ and $\text{Vol}(\pi^{(d-0)}(P)) = \text{Vol}(\mathbb{R}^0) = 1$.

An example of 1-integral polytope

Example: $P = \text{conv}\{(0, 0, 0), (4, 0, 0), (3, 6, 0), (2, 2, 2)\}$. One can check that P is 1-integral.

$$i(P, m) = 8m^3 + 10m^2 + 4m + 1,$$

and $\text{Vol}(\pi^{(d-1)}(P)) = \text{Vol}([0, 4]) = 4$ and $\text{Vol}(\pi^{(d-0)}(P)) = \text{Vol}(\mathbb{R}^0) = 1$.

Question: Is there any interesting way to calculate the higher degree coefficients?

Slices of a polytope


Definition 8. For any $\mathbf{y} \in \pi^{(d-k)}(P)$, we define *the slice of P over \mathbf{y}* , denoted by $\pi_{d-k}(\mathbf{y}, P)$, to be the intersection of P with the inverse image of \mathbf{y} under $\pi^{(d-k)}$.

Slices of a polytope

Definition 8. For any $\mathbf{y} \in \pi^{(d-k)}(P)$, we define *the slice of P over \mathbf{y}* , denoted by $\pi_{d-k}(\mathbf{y}, P)$, to be the intersection of P with the inverse image of \mathbf{y} under $\pi^{(d-k)}$.

Example: $P = \text{conv}\{(0, 0, 0), (4, 0, 0), (3, 6, 0), (2, 2, 2)\}$.

Recall $\pi^{(d-1)}(P) = [0, 4]$ and $i(P, m) = 8m^3 + 10m^2 + 4m + 1$.




Slices of a polytope

Definition 8. For any $\mathbf{y} \in \pi^{(d-k)}(P)$, we define *the slice of P over \mathbf{y}* , denoted by $\pi_{d-k}(\mathbf{y}, P)$, to be the intersection of P with the inverse image of \mathbf{y} under $\pi^{(d-k)}$.

Example: $P = \text{conv}\{(0, 0, 0), (4, 0, 0), (3, 6, 0), (2, 2, 2)\}$.

Recall $\pi^{(d-1)}(P) = [0, 4]$ and $i(P, m) = 8m^3 + 10m^2 + 4m + 1$.



$i(\pi_2(0, P), m) = 1, i(\pi_2(1, P), m) = m^2 + 2m + 1, i(\pi_2(2, P), m) = 4m^2 + 4m + 1, i(\pi_2(3, P), m) = 3m^2 + 4m + 1$ and $i(\pi_2(4, P), m) = 1$. Their sum is

$$8m^2 + 10m + 5.$$

Main theorems

Theorem 9. *If P is k -integral, then the coefficient of m^ℓ in $i(P, m)$ is*

$$\begin{cases} \text{Vol}(\pi^{d-\ell}(P)) & \text{if } 0 \leq \ell \leq k, \\ \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{coefficient of } m^{\ell-k} \text{ in } i(\pi_{d-k}(\mathbf{y}, P), m) & \text{if } k+1 \leq \ell \leq d \end{cases}$$

Main theorems

Theorem 9. *If P is k -integral, then the coefficient of m^ℓ in $i(P, m)$ is*

$$\begin{cases} \text{Vol}(\pi^{d-\ell}(P)) & \text{if } 0 \leq \ell \leq k, \\ \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{coefficient of } m^{\ell-k} \text{ in } i(\pi_{d-k}(\mathbf{y}, P), m) & \text{if } k+1 \leq \ell \leq d \end{cases}$$

Theorem 10. *Suppose $k < d$. If P is k -integral, then*

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)),$$

where Vol_{d-k} is the volume with respect to the lattice \mathbb{Z}^{d-k} .

Main theorems

Theorem 9. *If P is k -integral, then the coefficient of m^ℓ in $i(P, m)$ is*

$$\begin{cases} \text{Vol}(\pi^{d-\ell}(P)) & \text{if } 0 \leq \ell \leq k, \\ \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{coefficient of } m^{\ell-k} \text{ in } i(\pi_{d-k}(\mathbf{y}, P), m) & \text{if } k+1 \leq \ell \leq d \end{cases}$$

Theorem 10. *Suppose $k < d$. If P is k -integral, then*

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)),$$


where Vol_{d-k} is the volume with respect to the lattice \mathbb{Z}^{d-k} .

Definition 11. We define the *k th S -volume of P* to be


$$\text{SVol}^k(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)).$$

Remarks


Remark 12.




$$\text{SVol}^0(P) = \text{Vol}(P)$$



$$\text{SVol}^1(P)$$







$$\text{SVol}^2(P)$$



$$\text{SVol}^d(P) = |P \cap \mathbb{Z}^d|$$

Remarks

Remark 12.



$$\text{SVol}^0(P) = \text{Vol}(P)$$

$$\text{SVol}^1(P)$$

$$\text{SVol}^2(P)$$

$$\text{SVol}^d(P) = |P \cap \mathbb{Z}^d|$$

Remark 13. Theorem 10 says if $k < d$ and P is k -integral, then $\text{Vol}(P) = \text{SVol}^k(P)$.

Note that P is ℓ -integral for any $\ell \leq k$, so we have

$$\text{Vol}(P) = \text{SVol}^0(P) = \text{SVol}^1(P) = \cdots = \text{SVol}^k(P).$$


Remarks

Remark 14. The condition k -integral in Theorem 10 can be relaxed to $(k - 1)$ -integral and *in k -general position*.

Remarks

Remark 14. The condition k -integral in Theorem 10 can be relaxed to $(k - 1)$ -integral and *in k -general position*.

Example: 1-dimensional affine space in \mathbb{R}^2 . integral vs in general position:




In general, U is in general position if and only if $\text{dir}(U) = (1, y_1, \dots, y_d) \in \mathbb{R}^d$, or equivalently U is not perpendicular to the x_1 -axis.

Remarks

Remark 14. The condition k -integral in Theorem 10 can be relaxed to $(k - 1)$ -integral and *in k -general position*.

Example: 1-dimensional affine space in \mathbb{R}^2 . integral vs in general position:



In general, U is in general position if and only if $\text{dir}(U) = (1, y_1, \dots, y_d) \in \mathbb{R}^d$, or equivalently U is not perpendicular to the x_1 -axis.

Theorem 9 can be reduced to Theorem 10.

Possible Applications

- ▮► Prove positivity conjectures of special families of polytopes: Birkhoff polytopes, matroid polytopes.

Possible Applications

- ➡ Prove positivity conjectures of special families of polytopes: Birkhoff polytopes, matroid polytopes.
- ➡ Calculate the lower degree coefficients of Ehrhart polynomial quickly.


Possible Applications

- ➡ Prove positivity conjectures of special families of polytopes: Birkhoff polytopes, matroid polytopes.
- ➡ Calculate the lower degree coefficients of Ehrhart polynomial quickly.
- ➡ Algorithm to calculate volumes of rational polytopes: for any rational polytope P , one can always choose a coordinate system such that P is in 1-general position. Choose $D \in \mathbb{N}$ such that DP is integral. Then

$$\text{Vol}(P) = \frac{1}{D} \sum_{\mathbf{y} \in \pi^{(d-1)}(P) \cap \frac{1}{D}\mathbb{Z}^1} \text{Vol}_{d-1}(\pi_{d-1}(\mathbf{y}, P)).$$


Reduction to volume formula

- i. If P is fully integral, then in particular P is $(d-1)$ -integral. For any $\mathbf{y} \in \pi^{(1)}(P) \cap \mathbb{Z}^{d-1}$, the slice $\pi_1(\mathbf{y}, P)$ is either a 1-dimensional integral polytope, or a lattice point. In either case, we have that $|\pi_1(\mathbf{y}, P) \cap \mathbb{Z}^d| = 1 + \text{Vol}_1(\pi_1(\mathbf{y}, P))$.



Reduction to volume formula

- i. If P is fully integral, then in particular P is $(d-1)$ -integral. For any $\mathbf{y} \in \pi^{(1)}(P) \cap \mathbb{Z}^{d-1}$, the slice $\pi_1(\mathbf{y}, P)$ is either a 1-dimensional integral polytope, or a lattice point. In either case, we have that $|\pi_1(\mathbf{y}, P) \cap \mathbb{Z}^d| = 1 + \text{Vol}_1(\pi_1(\mathbf{y}, P))$.



Then

$$\begin{aligned}
 |P \cap \mathbb{Z}^d| &= \sum_{\mathbf{y} \in \pi^{(1)}(P) \cap \mathbb{Z}^{d-1}} |\pi_1(\mathbf{y}, P) \cap \mathbb{Z}^d| \\
 &= \sum_{\mathbf{y} \in \pi^{(1)}(P) \cap \mathbb{Z}^{d-1}} (1 + \text{Vol}_1(\pi_1(\mathbf{y}, P))) = |\pi^{(1)}(P) \cap \mathbb{Z}^{d-1}| + \text{Vol}(P).
 \end{aligned}$$

Reduction to volume formula

However, $\pi^{(1)}(P)$ is fully integral. Hence,

$$|P \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)).$$

Note that P is k -integral $\Rightarrow mP$ is k -integral as well. Therefore,

$$i(P, m) = |mP \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(mP)) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

Reduction to volume formula

However, $\pi^{(1)}(P)$ is fully integral. Hence,

$$|P \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)).$$

Note that P is k -integral $\Rightarrow mP$ is k -integral as well. Therefore,

$$i(P, m) = |mP \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(mP)) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

- ii. If $k \leq d-1$, the projection $\pi^{(d-k)}(P)$ is fully integral and for any $\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k$, the slice $\pi_{d-k}(\mathbf{y}, P)$ is an integral polytope.

Reduction to volume formula

However, $\pi^{(1)}(P)$ is fully integral. Hence,

$$|P \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)).$$

Note that P is k -integral $\Rightarrow mP$ is k -integral as well. Therefore,

$$i(P, m) = |mP \cap \mathbb{Z}^d| = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(mP)) = \sum_{i=0}^d \text{Vol}(\pi^{(d-i)}(P)) m^i.$$

- ii. If $k \leq d-1$, the projection $\pi^{(d-k)}(P)$ is fully integral and for any $\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k$, the slice $\pi_{d-k}(\mathbf{y}, P)$ is an integral polytope.

We prove Theorem 9 by using the result on fully integral polytopes as well as a local formula relating the number of lattice points to volumes of faces for integral polytopes obtained by Morelli, McMullen, Pommersheim-Thomas, Berline-Vergne.

Sketch of the proof of Theorem 10

Recall the theorem: If P is k -integral, then

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)) = \text{SVol}^k(P).$$

Sketch of the proof of Theorem 10

Recall the theorem: If P is k -integral, then

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)) = \text{SVol}^k(P).$$

i. We reduce the problem to the case of $k = 1$.


Sketch of the proof of Theorem 10

Recall the theorem: If P is k -integral, then

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)) = \text{SVol}^k(P).$$

i. We reduce the problem to the case of $k = 1$.

Idea: If P is k -integral (with $k > 1$), then each slice $\pi_{d-1}(\mathbf{y}, P)$ contributing to $\text{SVol}^1(P)$ is $(k-1)$ -integral.




Sketch of the proof of Theorem 10

Recall the theorem: If P is k -integral, then

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)) = \text{SVol}^k(P).$$

i. We reduce the problem to the case of $k = 1$.

Idea: If P is k -integral (with $k > 1$), then each slice $\pi_{d-1}(\mathbf{y}, P)$ contributing to $\text{SVol}^1(P)$ is $(k - 1)$ -integral.



ii. We reduce the problem to the case of a simplex with $k = 1$.


Sketch of the proof of Theorem 10

Recall the theorem: If P is k -integral, then

$$\text{Vol}(P) = \sum_{\mathbf{y} \in \pi^{(d-k)}(P) \cap \mathbb{Z}^k} \text{Vol}_{d-k}(\pi_{d-k}(\mathbf{y}, P)) = \text{SVol}^k(P).$$

i. We reduce the problem to the case of $k = 1$.

Idea: If P is k -integral (with $k > 1$), then each slice $\pi_{d-1}(\mathbf{y}, P)$ contributing to $\text{SVol}^1(P)$ is $(k - 1)$ -integral.



ii. We reduce the problem to the case of a simplex with $k = 1$.

iii. Prove the case of a simplex with $k = 1$.