

Lecture 11: Bayes Nets: Inference

Shuai Li


John Hopcroft Center, Shanghai Jiao Tong University

<https://shuaili8.github.io>

<https://shuaili8.github.io/Teaching/CS410/index.html>

Part of slide credits: CMU AI & <http://ai.berkeley.edu>

Bayes Rule



Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

- Why is this at all helpful?
 - Lets us build one conditional from its reverse
 - Often one conditional is tricky but the other one is simple
 - Foundation of many systems (e.g. ASR, MT)
- In the running for most important AI equation!

That's my rule!



Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{cause}|\text{effect}) = \frac{P(\text{effect}|\text{cause})P(\text{cause})}{P(\text{effect})}$$

- Example:

- M: meningitis, S: stiff neck

$$\left. \begin{aligned} P(+m) &= 0.0001 \\ P(+s|+m) &= 0.8 \\ P(+s|-m) &= 0.01 \end{aligned} \right\} \text{Example givens}$$

$$P(+m|+s) = \frac{P(+s|+m)P(+m)}{P(+s)} = \frac{P(+s|+m)P(+m)}{P(+s|+m)P(+m) + P(+s|-m)P(-m)} = \frac{0.8 \times 0.0001}{0.8 \times 0.0001 + 0.01 \times 0.999}$$

- Note: posterior probability of meningitis still very small
- Note: you should still get stiff necks checked out! Why?

Quiz: Bayes' Rule

- Given:

$$P(W)$$

R	P
sun	0.8
rain	0.2

$$P(D|W)$$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

- What is $P(W \mid \text{dry})$?

Quiz: Bayes' Rule 2

- Given:

$P(W)$

R	P
sun	0.8
rain	0.2

$P(D|W)$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

- What is $P(W | \text{dry})$?

$$P(\text{sun} | \text{dry}) \propto P(\text{dry} | \text{sun})P(\text{sun}) = .9 * .8 = .72$$

$$P(\text{rain} | \text{dry}) \propto P(\text{dry} | \text{rain})P(\text{rain}) = .3 * .2 = .06$$

$$P(\text{sun} | \text{dry}) = 12/13$$

$$P(\text{rain} | \text{dry}) = 1/13$$

Ghostbusters, Revisited


- Let's say we have two distributions:
 - **Prior distribution** over ghost location: $P(G)$
 - Let's say this is uniform
 - Sensor reading model: $P(R | G)$
 - Given: we know what our sensors do
 - R = reading color measured at $(1,1)$
 - E.g. $P(R = \text{yellow} | G=(1,1)) = 0.1$
- We can calculate the **posterior distribution** $P(G|r)$ over ghost locations given a reading using Bayes' rule:

$$P(g|r) \propto P(r|g)P(g)$$


0.11	0.11	0.11
0.11	0.11	0.11
0.11	0.11	0.11

0.17	0.10	0.10
0.09	0.17	0.10
<0.01	0.09	0.17

Video of Demo Ghostbusters with Probability



Inference





Recall: Bayes' Net Representation

- A directed, acyclic graph, one node per random variable
- A conditional probability table (CPT) for each node
 - A collection of distributions over X , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$

- Bayes' nets implicitly encode joint distributions
 - As a product of local conditional distributions
 - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$



Inference

- Inference: calculating some useful quantity from a joint probability distribution




- Examples:

- Posterior probability

$$P(Q|E_1 = e_1, \dots, E_k = e_k)$$

- Most likely explanation:

$$\operatorname{argmax}_q P(Q = q|E_1 = e_1 \dots)$$



Queries

- What is the probability of *this* given what I know?

$$P(q \mid e) = \frac{P(q, e)}{P(e)} = \frac{\sum_{h_1} \sum_{h_2} P(q, h_1, h_2, e)}{P(e)}$$

- What are the probabilities of all the possible outcomes (given what I know)?

$$P(Q \mid e) = \frac{P(Q, e)}{P(e)} = \frac{\sum_{h_1} \sum_{h_2} P(Q, h_1, h_2, e)}{P(e)}$$

- Which outcome is the most likely outcome (given what I know)?

$$\begin{aligned} \operatorname{argmax}_{q \in Q} P(q \mid e) &= \operatorname{argmax}_{q \in Q} \frac{P(q, e)}{P(e)} \\ &= \operatorname{argmax}_{q \in Q} \frac{\sum_{h_1} \sum_{h_2} P(q, h_1, h_2, e)}{P(e)} \end{aligned}$$

Inference by Enumeration in Joint Distributions

- General case:


- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots, X_n \\ \text{All variables} \end{array} \right\}$

- We want:

** Works fine with multiple query variables, too*


$$P(Q|e_1 \dots e_k)$$

- Step 1: Select the entries consistent with the evidence



x	P(x)
-3	0.05
-1	0.25
0	0.07
1	0.2
5	0.01

- Step 2: Sum out H to get joint of Query and evidence



- Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$


$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots, X_n}$$

Inference by Enumeration in Bayes' Net

- Given unlimited time, inference in BNs is easy

$$\begin{aligned}P(B \mid +j, +m) &\propto_B P(B, +j, +m) \\&= \sum_{e,a} P(B, e, a, +j, +m) \\&= \sum_{e,a} P(B)P(e)P(a|B, e)P(+j|a)P(+m|a)\end{aligned}$$



$$\begin{aligned}=&P(B)P(+e)P(+a|B, +e)P(+j|+a)P(+m|+a) + P(B)P(+e)P(-a|B, +e)P(+j|-a)P(+m|-a) \\&P(B)P(-e)P(+a|B, -e)P(+j|+a)P(+m|+a) + P(B)P(-e)P(-a|B, -e)P(+j|-a)P(+m|-a)\end{aligned}$$

Example: Traffic Domain


- Random Variables

- R: Raining
- T: Traffic
- L: Late for class!

$$P(L) = ?$$

$$= \sum_{r,t} P(r, t, L)$$

$$= \sum_{r,t} P(r)P(t|r)P(L|t)$$



$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

Inference by Enumeration: Procedural Outline

- Track objects called **factors**
- Initial factors are local CPTs (one per node)

$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Any known values are selected
 - E.g. if we know $L = +\ell$, the initial factors are

$$P(R)$$

+r	0.1
-r	0.9


$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9


$$P(+\ell|T)$$

+t	+l	0.3
-t	+l	0.1


- Procedure: Join all factors, then sum out all hidden variables



Operation 1: Join Factors




- First basic operation: **joining factors**
- Combining factors:
 - **Just like a database join**
 - Get all factors over the joining variable
 - Build a new factor over the union of the variables involved
- Example: Join on R



- Computation for each entry: pointwise products $\forall r, t : P(r, t) = P(r) \cdot P(t|r)$

Example: Multiple Joins



$P(R)$

+r	0.1
-r	0.9

$P(T|R)$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$P(L|T)$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

Join R




$P(R, T)$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

$P(L|T)$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9


Join T



R, T, L

$P(R, T, L)$

+r	+t	+l	0.024
+r	+t	-l	0.056
+r	-t	+l	0.002
+r	-t	-l	0.018
-r	+t	+l	0.027
-r	+t	-l	0.063
-r	-t	+l	0.081
-r	-t	-l	0.729



Operation 2: Eliminate

- Second basic operation: **marginalization**
- Take a factor and sum out a variable
 - Shrinks a factor to a smaller one
 - A **projection** operation

- Example:

$$P(R, T)$$


+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

sum R



$$P(T)$$

+t	0.17
-t	0.83



Multiple Elimination

R, T, L

$P(R, T, L)$

+r	+t	+l	0.024
+r	+t	-l	0.056
+r	-t	+l	0.002
+r	-t	-l	0.018
-r	+t	+l	0.027
-r	+t	-l	0.063
-r	-t	+l	0.081
-r	-t	-l	0.729

Sum
out R



T, L

$P(T, L)$

+t	+l	0.051
+t	-l	0.119
-t	+l	0.083
-t	-l	0.747



Sum
out T



L

$P(L)$

+l	0.134
-l	0.866




Thus Far: Multiple Join, Multiple Eliminate (= Inference by Enumeration)

$$P(R)$$

$$P(T|R) \quad \longrightarrow \quad P(R, T, L) \quad \longrightarrow \quad P(L)$$

$$P(L|T)$$


Thus Far: Multiple Join, Multiple Eliminate (= Inference by Enumeration)



Inference by Enumeration in Bayes Net

- Reminder of inference by enumeration:
 - Any probability of interest can be computed by summing entries from the joint distribution
 - Entries from the joint distribution can be obtained from a BN by multiplying the corresponding conditional probabilities

$$\begin{aligned}P(B \mid j, m) &= \alpha P(B, j, m) \\&= \alpha \sum_{e,a} P(B, e, a, j, m) \\&= \alpha \sum_{e,a} P(B) P(e) P(a \mid B, e) P(j \mid a) P(m \mid a)\end{aligned}$$



- So inference in Bayes nets means computing sums of products of numbers: sounds easy!!
- Problem: sums of *exponentially many* products!

Can we do better?

- Consider

- $x_1y_1z_1 + x_1y_1z_2 + x_1y_2z_1 + x_1y_2z_2 + x_2y_1z_1 + x_2y_1z_2 + x_2y_2z_1 + x_2y_2z_2$
- 16 multiplies, 7 adds
- Lots of repeated subexpressions!

- Rewrite as

- $(x_1 + x_2)(y_1 + y_2)(z_1 + z_2)$
- 2 multiplies, 3 adds


$$\begin{aligned} \sum_e \sum_a P(B) P(e) P(a | B, e) P(j | a) P(m | a) \\ = P(B) P(+e) P(+a | B, +e) P(j | +a) P(m | +a) \\ + P(B) P(-e) P(+a | B, -e) P(j | +a) P(m | +a) \\ + P(B) P(+e) P(-a | B, +e) P(j | -a) P(m | -a) \\ + P(B) P(-e) P(-a | B, -e) P(j | -a) P(m | -a) \end{aligned}$$

- Lots of repeated subexpressions!


Variable Elimination

Inference by Enumeration vs. Variable Elimination

- Why is inference by enumeration so slow?
 - You join up the whole joint distribution before you sum out the hidden variables



- Idea: **interleave joining and marginalizing!**
 - Called “Variable Elimination”
 - Still NP-hard, but usually much faster than inference by enumeration



Inference Overview

- Given random variables Q, H, E (query, hidden, evidence)

- We know how to do inference on a joint distribution

$$\begin{aligned} P(q|e) &= \alpha P(q, e) \\ &= \alpha \sum_{h \in \{h_1, h_2\}} P(q, h, e) \end{aligned}$$

- We know Bayes nets can break down joint in to CPT factors

$$\begin{aligned} P(q|e) &= \alpha \sum_{h \in \{h_1, h_2\}} P(h) P(q|h) P(e|q) \\ &= \alpha [P(h_1) P(q|h_1) P(e|q) + P(h_2) P(q|h_2) P(e|q)] \end{aligned}$$




- But we can be more efficient

$$\begin{aligned} P(q|e) &= \alpha P(e|q) \sum_{h \in \{h_1, h_2\}} P(h) P(q|h) \\ &= \alpha P(e|q) [P(h_1) P(q|h_1) + P(h_2) P(q|h_2)] \\ &= \alpha P(e|q) P(q) \end{aligned}$$

- Now just extend to larger Bayes nets and a variety of queries

Enumeration
Variable Elimination

Traffic Domain



$$P(L) = ?$$

- Inference by Enumeration

$$= \sum_t \sum_r \underbrace{P(L|t)P(r)P(t|r)}_{\text{Join on } r}$$

$$\underbrace{\hspace{10em}}_{\text{Join on } t}$$

$$\underbrace{\hspace{10em}}_{\text{Eliminate } r}$$

$$\underbrace{\hspace{10em}}_{\text{Eliminate } t}$$

■ Variable Elimination


$$= \sum_t P(L|t) \underbrace{\sum_r P(r)P(t|r)}_{\text{Join on } r}$$

$$\underbrace{\hspace{10em}}_{\text{Eliminate } r}$$

$$\underbrace{\hspace{10em}}_{\text{Join on } t}$$



$$\underbrace{\hspace{10em}}_{\text{Eliminate } t}$$

Marginalizing Early (= Variable Elimination)




Answer Any Query from Bayes Net (Previous)

Bayes Net



$P(A)$ $P(B|A)$ $P(C|A)$ $P(D|C)$ $P(E|C)$

Joint







Query

$P(a | e)$

Next: Answer Any Query from Bayes Net

Bayes Net




$P(A)$ $P(B|A)$ $P(C|A)$ $P(D|C)$ $P(E|C)$

Query



$P(a | e)$

Marginalizing Early! (aka VE)



Evidence

- If evidence, start with factors that select that evidence

- No evidence uses these initial factors:

$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Computing $P(L|+r)$, the initial factors become:

$$P(+r)$$

+r	0.1
----	-----


$$P(T|+r)$$

+r	+t	0.8
+r	-t	0.2

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- We eliminate all vars other than query + evidence



Evidence II

- Result will be a selected joint of query and evidence
 - E.g. for $P(L \mid +r)$, we would end up with:

$$P(+r, L)$$

+r	+l	0.026
+r	-l	0.074


Normalize



$$P(L \mid +r)$$

+l	0.26
-l	0.74

- To get our answer, just normalize this!
- That's it!




Inference by Enumeration


- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots, X_n \\ \text{All variables} \end{array} \right\}$

- Step 1: Select the entries consistent with the evidence



- Step 2: Sum out H to get joint of Query and evidence



- Compute joint

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} P(Q, \underbrace{h_1 \dots h_r}_{\text{All variables}}, e_1 \dots e_k)$$

- Sum out hidden variables X_1, X_2, \dots, X_n

- We want:

** Works fine with multiple query variables, too*

$$P(Q|e_1 \dots e_k)$$

- Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$


$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

Variable Elimination

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots, X_n \\ \text{All variables} \end{array} \right\}$


- Step 1: Select the entries consistent with the evidence



x	P(x)
-3	0.05
-1	0.25
0	0.07
1	0.2
5	0.01

2 0.15

- Step 2: Sum out H to get joint of Query and evidence



$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} P(Q, \underbrace{h_1 \dots h_r}_{\text{Hidden Variables}}, e_1 \dots e_k)$$

- Interleave joining and summing out X_1, X_2, \dots, X_n

- We want:

** Works fine with multiple query variables, too*

$$P(Q|e_1 \dots e_k)$$

- Step 3: Normalize


$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$



$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$


General Variable Elimination

- Query: $P(Q|E_1 = e_1, \dots, E_k = e_k)$
- Start with initial factors:
 - Local CPTs (but instantiated by evidence)
- While there are still hidden variables (not Q or evidence):
 - Pick a hidden variable H
 - Join all factors mentioning H
 - Eliminate (sum out) H
- Join all remaining factors and normalize



x	P(x)
-3	0.05
-1	0.25
0	0.07
1	0.2
5	0.01




$$\times \frac{1}{Z}$$


Variable Elimination

```
function VariableElimination( $Q$ ,  $e$ ,  $bn$ ) returns a distribution over  $Q$   
   $factors \leftarrow []$   
  for each  $var$  in ORDER( $bn.vars$ ) do  
     $factors \leftarrow [MAKE-FACTOR(var,  $e$ ) |  $factors$ ]  
    if  $var$  is a hidden variable then  
       $factors \leftarrow SUM-OUT(var, factors)$   
  return NORMALIZE(POINTWISE-PRODUCT( $factors$ ))$ 
```

Example

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------



$$P(B|j, m) \propto P(B, j, m)$$

$$= \sum_{e, a} P(B, j, m, e, a)$$

$$= \sum_{e, a} P(B)P(e)P(a|B, e)P(j|a)P(m|a)$$

$$= \sum_e P(B)P(e) \sum_a P(a|B, e)P(j|a)P(m|a)$$

$$= \sum_e P(B)P(e)f_1(j, m|B, e)$$

$$= P(B) \sum_e P(e)f_1(j, m|B, e)$$

$$= P(B)f_2(j, m|B)$$

marginal can be obtained from joint by summing out

use Bayes' net joint distribution expression

use $x*(y+z) = xy + xz$

joining on a, and then summing out gives f_1

use $x*(y+z) = xy + xz$


joining on e, and then summing out gives f_2

All we are doing is exploiting $uwy + uwz + uxy + uxz + vwy + vwz + vxy + vxz = (u+v)(w+x)(y+z)$ to improve computational efficiency!

Example (cont'd)

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------




Choose A


$$P(A|B, E)$$

$$P(j|A)$$

$$P(m|A)$$



$$P(j, m, A|B, E)$$



$$P(j, m|B, E)$$

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------

Example (cont'd)

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------


Choose E

$$\begin{array}{c} P(E) \\ P(j, m|B, E) \end{array} \xrightarrow{\times} P(j, m, E|B) \xrightarrow{\Sigma} P(j, m|B)$$

$P(B)$	$P(j, m B)$
--------	-------------

Finish with B

$$\begin{array}{c} P(B) \\ P(j, m|B) \end{array} \xrightarrow{\times} P(j, m, B) \xrightarrow{\text{Normalize}} P(B|j, m)$$



Another Variable Elimination Example

Query: $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$P(Z), P(X_1|Z), P(X_2|Z), P(X_3|Z), P(y_1|X_1), P(y_2|X_2), P(y_3|X_3)$$

Eliminate X_1 , this introduces the factor $f_1(y_1|Z) = \sum_{x_1} P(x_1|Z)P(y_1|x_1)$, and we are left with:

$$P(Z), P(X_2|Z), P(X_3|Z), P(y_2|X_2), P(y_3|X_3), f_1(y_1|Z)$$

Eliminate X_2 , this introduces the factor $f_2(y_2|Z) = \sum_{x_2} P(x_2|Z)P(y_2|x_2)$, and we are left with:

$$P(Z), P(X_3|Z), P(y_3|X_3), f_1(y_1|Z), f_2(y_2|Z)$$


Eliminate Z , this introduces the factor $f_3(y_1, y_2, X_3) = \sum_z P(z)P(X_3|z)f_1(y_1|Z)f_2(y_2|Z)$, and we are left with:

$$P(y_3|X_3), f_3(y_1, y_2, X_3)$$

No hidden variables left. Join the remaining factors to get:

$$f_4(y_1, y_2, y_3, X_3) = P(y_3|X_3), f_3(y_1, y_2, X_3)$$


Normalizing over X_3 gives $P(X_3|y_1, y_2, y_3) = f_4(y_1, y_2, y_3, X_3) / \sum_{x_3} f_4(y_1, y_2, y_3, x_3)$



Computational complexity critically depends on the largest factor being generated in this process. Size of factor = number of entries in table. In example above (assuming binary) all factors generated are of size 2 --- as they all only have one variable (Z , Z , and X_3 respectively).

Variable Elimination Ordering

- For the query $P(X_n | y_1, \dots, y_n)$ work through the following two different orderings as done in previous slide: Z, X_1, \dots, X_{n-1} and X_1, \dots, X_{n-1}, Z . What is the size of the maximum factor generated for each of the orderings?



- Answer: 2^n versus 2 (assuming binary)
- In general: the ordering can greatly affect efficiency


Detail of size 4

- Elimination order: C, B, A, Z

- $P(D) = \alpha \sum_{z,a,b,c} P(D|z) P(z) P(a|z) P(b|z) P(c|z)$
- $= \alpha \sum_z P(D|z) P(z) \sum_a P(a|z) \sum_b P(b|z) \sum_c P(c|z)$
- Largest factor has 2 variables (D,Z)

- Elimination order: Z, C, B, A

- $P(D) = \alpha \sum_{a,b,c,z} P(a|z) P(b|z) P(c|z) P(D|z) P(z)$
- $= \alpha \sum_a \sum_b \sum_c \sum_z P(a|z) P(b|z) P(c|z) P(D|z) P(z)$
- Largest factor has 4 variables (A,B,C,D)



- In general, with n leaves, factor of size 2^n

VE: Computational and Space Complexity

- The computational and space complexity of variable elimination is determined by the largest factor
- The elimination ordering can greatly affect the size of the largest factor.
 - E.g., previous slide's example 2^n vs. 2
- Does there always exist an ordering that only results in small factors?
 - No!

Worst Case Complexity?

- CSP:

$$(x_1 \vee x_2 \vee \neg x_3) \wedge (\neg x_1 \vee x_3 \vee \neg x_4) \wedge (x_2 \vee \neg x_2 \vee x_4) \wedge (\neg x_3 \vee \neg x_4 \vee \neg x_5) \wedge (x_2 \vee x_5 \vee x_7) \wedge (x_4 \vee x_5 \vee x_6) \wedge (\neg x_5 \vee x_6 \vee \neg x_7) \wedge (\neg x_5 \vee \neg x_6 \vee x_7)$$

$$P(X_i = 0) = P(X_i = 1) = 0.5$$

$$Y_1 = X_1 \vee X_2 \vee \neg X_3$$

...

$$Y_8 = \neg X_5 \vee X_6 \vee X_7$$

$$Y_{1,2} = Y_1 \wedge Y_2$$


...

$$Y_{7,8} = Y_7 \wedge Y_8$$

$$Y_{1,2,3,4} = Y_{1,2} \wedge Y_{3,4}$$

$$Y_{5,6,7,8} = Y_{5,6} \wedge Y_{7,8}$$

$$Z = Y_{1,2,3,4} \wedge Y_{5,6,7,8}$$



- If we can answer $P(z)$ equal to zero or not, we answered whether the 3-SAT problem has a solution
- Hence inference in Bayes' nets is NP-hard. No known efficient probabilistic inference in general


“Easy” Structures: Polytrees

- A polytree is a directed graph with no undirected cycles
- For poly-trees you can always find an ordering that is efficient
 - Try it!!
- Cut-set conditioning for Bayes' net inference
 - Choose set of variables such that if removed only a polytree remains
 - Exercise: Think about how the specifics would work out!

Variable Elimination: The basic ideas

- Move summations inwards as far as possible

$$\begin{aligned} P(B \mid j, m) &= \alpha \sum_e \sum_a P(B) P(e) P(a \mid B, e) P(j \mid a) P(m \mid a) \\ &= \alpha P(B) \sum_e P(e) \sum_a P(a \mid B, e) P(j \mid a) P(m \mid a) \end{aligned}$$




- Do the calculation from the inside out

- I.e., sum over a first, then sum over e
- Problem: $P(a \mid B, e)$ isn't a single number, it's a bunch of different numbers depending on the values of B and e
- Solution: use arrays of numbers (of various dimensions) with appropriate operations on them; these are called **factors**


Operation 1: Pointwise product

- First basic operation: pointwise product of factors (similar to a database join, **not** matrix multiply!)
 - New factor has **union** of variables of the two original factors
 - Each entry is the product of the corresponding entries from the original factors
- Example: $P(A) \times P(J|A) = P(A,J)$



$P(A)$		\times	$P(J A)$			$=$	$P(A,J)$		
A \ J			true	false	A \ J		true	false	
true	0.1		true	0.9	0.1		true	0.09	0.01
false	0.9		false	0.05	0.95		false	0.045	0.855

Example: Making larger factors



- Example: $P(J/A) \times P(M/A) = P(J,M/A)$

$P(J/A)$

$A \setminus J$	true	false
true	0.99	0.01
false	0.145	0.855

\times

$P(M/A)$

$A \setminus M$	true	false
true	0.97	0.03
false	0.019	0.891


$=$

$P(J,M/A)$

	$J \setminus M$		
	true	false	
$J \setminus M$	true	false	
true			18
false		.0003	

A=false
A=true

Example: Making larger factors 2



- Example: $f_1(U,V) \times f_2(V,W) \times f_3(W,X) = f_4(U,V,W,X)$
- Sizes: $[10,10] \times [10,10] \times [10,10] = [10,10,10,10]$
- i.e., 300 numbers blows up to 10,000 numbers!
- Factor blowup can make VE very expensive


Operation 2: Summing out a variable

- Second basic operation: **summing out** (or eliminating) a variable from a factor
 - Shrinks a factor to a smaller one
- Example: $\sum_j P(A, J) = P(A, j) + P(A, \neg j) = P(A)$

$P(A, J)$				
$A \setminus J$	true	false		
true	0.09	0.01		
false	0.045	0.855		


Sum out J

$P(A)$	
true	0.1
false	0.9



Summing out from a product of factors

- Project the factors each way first, then sum the products
- Example: $\sum_a P(a|B,e) P(j|a) P(m|a)$
 $= P(a|B,e) P(j|a) P(m|a) + P(\neg a|B,e) P(j|\neg a) P(m|\neg a)$
 $= P(a,j,m|B,e) + P(\neg a,j,m|B,e)$
 $= P(j,m|B,e)$



Summary

- Bayes

Shuai Li

<https://shuaili8.github.io>

Questions?