

Finite Element Method for Quantum Mechanics

Jerome Troy

May 13, 2021

What are We Going to do?

- Photonics - building electric circuitry for photons rather than electrons
- Why - photons travel at light speed (electrons are snails)
- Photons don't interact with each other (easier to deal with)
- Can play tricks with resonance to do computations




Figure: Figure taken from S. T. Fard et. al.

An Overview of Quantum Mechanics

- **Heisenberg Uncertainty Principle** : “ You cannot simultaneously know the position and momentum of a quantum particle”
- Classical rules $\mathbf{F} = m\mathbf{a}$ are changed to quantum rules: **Schödinger Equation**

$$i\hbar \frac{\partial \Psi}{\partial t} = \hat{H}[\Psi]$$

- \hbar - Planck's Constant (divided by 2π), \hat{H} : *Hermitian* operator which produces energy
- Qol: Ψ - wavefunction
- Given operator \hat{O} , giving property ω , the **Probability Density Function** for ω , ρ_ω :

$$\rho_\omega(\mathbf{x}, t) = \Psi^*(\mathbf{x}, t) \hat{O} [\Psi(\mathbf{x}, t)]$$

A Simple Case

- Particles interacting with an external potential $V(\mathbf{x})$ (time-independent, $\mathbf{x} \in \mathbb{R}^2$)
- Particles do not interact with each other (linear PDE)
- Energy: Kinetic + Potential
- Kinetic: $\frac{1}{2m}\hat{p}^2$, mass m , *momentum operator* $\hat{\mathbf{p}} = -i\hbar\nabla$
- Schödinger equation for a single particle:

$$i\hbar\frac{\partial\Psi}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\Psi + V(\mathbf{x})\Psi$$


- Nondimensionalize

$$\implies i\frac{\partial\Psi}{\partial t} = -\nabla^2\Psi + \nu V(\mathbf{x})\Psi$$

- An interesting question: **Can we understand tunnelling?**

Quantum Tunnelling

- Particle “trapped” (in the classical regime) in a potential well can escape!
- Consider the following example
- Input boundary: Γ_{in} , Other blue boundaries: Γ_{out} , all other boundaries: Γ_D



FEM and Variational Form

- $\Psi \in H^1(\Omega) \times C(\mathbb{R}_+)$
- Inner products (bra-ket notation):

$$\langle \phi | \psi \rangle = \int_{\Omega} \phi^* \psi \, dx, \quad \langle \phi | \mathcal{O} | \psi \rangle = \int_{\Omega} \phi^* \mathcal{O}(\psi) \, dx.$$

- Variational form: $\Phi \in H_{D,\text{in}}^1(\Omega)$ of the Schrödinger equation

$$i \langle \Phi | \partial_t | \Psi \rangle = \langle \nabla \Phi | \nabla \Psi \rangle + \langle \Phi | V | \Psi \rangle + \int_{\Gamma_{\text{out}}} \Phi^* \frac{\partial \Psi}{\partial n} \, d\sigma$$

- Φ zero on outer boundary and input boundary, nonzero on other (blue boundaries), wave can escape this way
- Enforce open boundary on Γ_{out} :

$$\frac{\partial \Psi}{\partial t} = i \frac{\partial \Psi}{\partial n} \implies \int_{\Gamma_{\text{out}}} \Phi^* \frac{\partial \Psi}{\partial n} \, d\sigma = i \frac{\partial}{\partial t} \int_{\Gamma_{\text{out}}} \Phi^* \Psi \, d\sigma.$$

Finite Dimensional FEM

- Boundary wrapper:

$$B(\Phi, \Psi) = \int_{\Gamma_{\text{out}}} \Phi^* \Psi \, d\sigma.$$

- Build triangulation \mathcal{T}_h on Ω , let $\phi_j(\mathbf{x})$ be a set of basis functions for $H_{D,\text{in}}^1(\Omega)$

$$\begin{aligned} \Rightarrow i \sum_{k=1}^N \langle \phi_j | \phi_k \rangle \lambda'_k(t) = \\ \sum_{k=1}^N [\langle \nabla \phi_j | \nabla \phi_k \rangle + \langle \phi_j | V | \phi_k \rangle + B(\phi_j, \phi_k)] \lambda_k(t) \end{aligned}$$

- ϕ_j are real functions in $H^1(\Omega)$, λ_k are complex functions
- Mass matrix: $\langle \phi_j | \phi_k \rangle$, Stiffness matrix: $\langle \nabla \phi_j | \nabla \phi_k \rangle$, “Potential” matrix: $\langle \phi_j | V | \phi_k \rangle$, “Boundary” matrix: $B(\phi_j, \phi_k)$

Doing this in FEniCS

Build mesh:


```
origin = Point(0, 0)
farcorner = Point(domainwidth, domainlength)
mesh = RectangleMesh(origin, farcorner, nx, ny)
```

Variational forms

```
H1 = FunctionSpace(mesh, "CG", 1)
Psi = TrialFunction(H1)
Phi = TestFunction(H1)
mass = assemble(Phi * Psi * dx)
stiffness = assemble(
    dot(grad(Phi), grad(Psi)) * dx)
boundary = assemble(Phi * Psi *
    [boundaryindicator] * ds)
potential = assemble(Phi * V * Psi * dx)
```


Time Stepping and Solution


- Crank-Nicolson method used for time stepping
- Domain : $h_{\min} = 0.1$
- $T_{\max} = 5, n_t = 1000 \Rightarrow \Delta t = 0.005$



Transmission Coefficient

- Computation:

$$C_{\text{trans}} = \frac{\langle \Psi | \mathbb{1}_{\Omega_1} | \Psi \rangle}{\langle \Psi | \mathbb{1}_{\Omega_2} | \Psi \rangle}.$$



Future Improvements

- Better boundary condition
- Proper B/C involves fractional derivatives

$$\partial_x \Psi = \sqrt{-i\partial_t - \partial_{yy}} \Psi \quad x \in \Gamma_{\text{out}}.$$

- Test various geometries
- Goal is to get optimal resonance for specified input frequency

$$(\omega_{\text{in}}, R_{\text{ring}}) \mapsto C_{\text{trans}}.$$

- Apply Maxwell equations - photons are electromagnetic wave packets

References I



Anton Arnold.

Open boundary conditions for wave propagation problems on unbounded domains.

2007.



Sahba Talebi Fard, Samantha M. Grist, Valentina Donzella, Shon A. Schmidt, Jonas Flueckiger, Xu Wang, Wei Shi, Andrew Millspaugh, Mitchell Webb, Daniel M. Ratner, Karen C. Cheung, and Lukas Chrostowski.

Label-free silicon photonic biosensors for use in clinical diagnostics.
International Society for Optical Engineering, 2013.



David J. Griffiths.

Introduction to Quantum Mechanics.

Pearson Education, Inc., New Jersey, 2005.

References II



Anders Logg, Kent-Andre Mardal, Garth N. Wells, et al.
Automated Solution of Differential Equations by the Finite Element Method.
Springer, 2012.



Larry Wasserman.
All of Statistics: A Concise Course in Statistical Inference.
Springer, New York, 2005.



Adam L. Washurn, L. Cary Gunn, and Ryan C. Bailey.
Label-free quantitation of a cancer biomarker in complex media using silicon photonic microring resonators.
Analytical Chemistry, 2009.

References III



Tong-Biao Wang, Xie-Wen Wen, Cheng-Ping Yin, and He-Zhou Wang.

The transmission characteristics of surface plasmon polaritons in ring resonator.

The Optical Society, 2009.