

Planetary magnetospheres

1. The Sun

- 1.1. 2-D heliosphere
- 1.2. 3-D structure
- 1.3. Transient processes – Solar flares and CMEs

2. Solar wind interaction with planets

- 2.1. Types of interaction
- 2.2. No atmosphere – No magnetic field
- 2.3. Role of the atmosphere
- 2.4. Role of the intrinsic magnetic field

3. Magnetospheric processes

- 3.1. Planetary effects
- 3.2. Radio emissions
- 3.3. Variations with planet rotation
- 3.4. Influence of nearby bodies


1. The Sun

1.1. 2-D heliosphere

Observations from the ecliptic plane → 2-D structure

Magnetic lines of force leaving the Sun form giant magnetosphere: heliosphere.

Entrained by Sun rotation - Studied with MHD (Magneto-Hydro-Dynamics)



Encarnaz & Bibring, 2002

Magnetic pressure

$$p = \frac{B^2}{2\mu_0}$$


Interplanetary field: average 4 nT

Solar wind: 99% H^+ and He^{++} ions (3.5-4.5%)

Some particles reach velocities of up to 700 km/s

1.2. 3-D structure

We need to measure away from the ecliptic plane:



Interplanetary magnetic field organised into sectors of opposite magnetic polarity:
“solar ballerina” model of Alfvén (1977)




Fig. 1.7. The solar ballerina proposed by H. Alfvén (1977) – schematic picture of the structure of the inner heliosphere near solar minimum. The Sun's poles are covered with large coronal holes of opposite magnetic field polarity. Co-rotating solar wind streams of opposite magnetic polarity with a heliospheric current sheet (HCS) in between them dominate the flow pattern. The situation shown here is typical for odd cycles.


Blondel & Mason, 2006

Two states of solar wind:

- low-speed wind (~ 400 km/s), combining jets of different velocities, close to ecliptic;
- high-speed wind (700-750 km/s), homogeneous, at high (solar) latitudes;

Separated by sharp boundary

1.3. Transient processes – Solar flares and CMEs



Blondel & Mason, 2006

Fig. 1.10. Three EUV-images of the solar corona taken by SoHO/EIT at 195 Å in 1996, 1998 and 1999. Solar maximum was in 2000. Note the strong change in coronal structure, the increase of the number of bright shining coronal regions with stronger underlying photospheric magnetic fields and the appearances of dark coronal regions with underlying open magnetic field structure at lower latitudes. Courtesy: SOHO/EIT consortium.

Solar flares

Sudden flash of light on the Sun – Can disrupt radio signals

Analogy: muzzle flash

Coronal Mass Ejections (CMEs)


Occur at active regions – New material (particles) preceded by shock wave

Analogy: cannonball

1.3. Transient processes - CMEs

High-speed jets interact with slower ambient solar wind.

Creates compression/rarefaction zones as well as shock waves.



LATEST RESULTS

NASA Parker Solar Probe (Astrophys. J., Feb. 2020)

> 0.25 AU from Sun

Unanticipated “switchbacks” in the solar wind

the Sun's magnetic field abruptly doubles-back on itself

Evidence for a “dust-free zone” — first hypothesized in 1929, but never successfully detected —
beginning an estimated 3.5 million miles from the Sun,


The first measurement of rotation in the solar wind — before it begins to flow radially, or straight out from the Sun, as it does near Earth,

Detections of never-before-seen particle events

so small that all trace of them is wiped out before they reach Earth.

ESA/NASA Solar Orbiter (17 Feb. 2020):

> 0.28 AU from Sun



Deployment of magnetic boom:

$< 10^{-3} B_{\text{earth}}$

First mission to image Sun's poles
(in 2025–2029)

Planetary magnetospheres

1. The Sun

- 1.1. 2-D heliosphere
- 1.2. 3-D structure
- 1.3. Transient processes – Solar flares and CMEs


2. Solar wind interaction with planets

- 2.1. Types of interaction
- 2.2. No atmosphere – No magnetic field
- 2.3. Role of the atmosphere
- 2.4. Role of the intrinsic magnetic field

3. Magnetospheric processes

- 3.1. Planetary effects
- 3.2. Radio emissions
- 3.3. Variations with planet rotation
- 3.4. Influence of nearby bodies

Interaction of solar wind with planet depends on its properties



solar wind directly arrives on planetary body

left: encounters atmosphere

right: encounters external magnetic field

Encrenaz and Bibring (2002)

Distance at which planetary field can stand off solar wind pressure: *Chapman-Ferraro distance*


$$R = R_{\text{planet}} \left(\frac{2 B_{\text{surface}}^2}{\mu_0 m_p \rho v_{\text{solar_wind}}^2} \right)^{\frac{1}{6}}$$

density of solar wind components



2.2. No atmosphere – No magnetic field


First case: **insulator, absorbing ions of the solar wind**. Example: the Moon.



Encrenaz and Bibring (2002)

Downstream: plasma-free cavity


Progressively filled by ion/e⁻ diffusion, further down




Solar wind experiments (Apollo 11, 12, 14, 15, 16)

Second case: **central body is good conductor. Magnetic lines bent/stretched over it.**

Example: Europa and Callisto (Galileo measurements)
Presence of liquid conducting layer (ocean?)



Encarnaz and Bibring (2002)



Europa



Callisto

*Moment of inertia
confirm possible
liquid ocean*

2.3. Role of the atmosphere

Dense atmosphere – No magnetic field


Solar wind interacts directly with atmosphere.

Examples: Venus, Mars, comets


Also:

Titan (+ interaction with Saturn's magnetic field)


Io (+ interaction with Jupiter's magnetosphere)



Horizontal entrainment
of the ionospheric plasma



Encrenaz and Bibring (2002)



Widely different characteristics of oxygen and hydrogen ion escape from Venus

R. Jarvinen,¹ E. Kallio,¹ S. Dyadechkin,¹ P. Janhunen,¹ and I. Sillanpää²







Figure 1. Flow vectors of the planetary (top) O⁺ and (bottom) H⁺ ions in the regions where their bulk fluxes are high ($n_i v_i > 10^{10} \text{ s}^{-1} \text{ m}^{-2}$). Three different projections onto the (listed from the left) xz, xy and yz planes are shown. Vectors are normalized and their orientation and coloring illustrate the velocity. The background coloring shows the densities at the $y = 0$, $z = 0$ and $x = -2R_V$ planes. The arrows in the lower left corners show the orientation of the velocity (V_{SW}), electric field (E_{SW}) and magnetic field (B_{SW}) of the solar wind.



2.4. Role of intrinsic magnetic field


Magnetic field + atmosphere. Magnetic field creates “cavity” in solar wind.



Examples: Mercury, Earth, Jupiter, Saturn,
Uranus, Neptune







1) Bow shock

2) Magnetosheath

3) Magnetopause

4) Magnetosphere

5) Northern tail lobe

6) Southern tail lobe

7) Plasmasphere

Asymmetric:

10 R_{Earth} sunward

> 200 R_{Earth} downstream

Planetary magnetospheres

1. The Sun

- 1.1. 2-D heliosphere
- 1.2. 3-D structure
- 1.3. Transient processes – Solar flares and CMEs

2. Solar wind interaction with planets

- 2.1. Types of interaction
- 2.2. No atmosphere – No magnetic field
- 2.3. Role of the atmosphere
- 2.4. Role of the intrinsic magnetic field

3. Magnetospheric processes

- 3.1. Planetary effects
- 3.2. Radio emissions
- 3.3. Variations with planet rotation
- 3.4. Influence of nearby bodies