

This is Table 6a

t_v	$\sum_{i=1}^{27} dN_i(t_v)$	$\sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v)$	$n_v = \sum_{i=1}^{66} Y_i(t_v)$	$n_{v1} = \sum_{i=1}^{66} I(8 \leq r_i) Y_i(t_v)$	$U_8(t_v)$
12	2	0	$16 + 13 = 29$	$10 + 10 = 20$	-1.37931
13	2	0	$17 + 14 = 31$	$13 + 11 = 24$	-1.54839
15	3	3	$22 + 23 = 45$	$20 + 20 = 40$	0.33333
16	4	3	$20 + 34 = 54$	$18 + 31 = 49$	-0.62963
17	2	1	$16 + 39 = 55$	$15 + 36 = 51$	-0.85454
18	6	6	$14 + 39 = 53$	$14 + 36 = 50$	0.33962
19	1	1	$8 + 39 = 47$	$8 + 36 = 44$	0.06383
20	2	2	$7 + 39 = 46$	$7 + 36 = 43$	0.13043
21	2	2	$5 + 39 = 44$	$5 + 36 = 41$	0.13636
22	1	1	$3 + 39 = 42$	$3 + 36 = 39$	0.07143
23	2	2	$2 + 39 = 41$	$2 + 36 = 38$	0.14634

Your tables "sampled Y" and "sampled IY" are the same.

1) In table "sampled Y" compute the sum of entries in columns titled 12- 23.

2) Denote the sum of entries in columns 12-23 by $Y^1(12) - Y^1(23)$, where "1" stands for the first run or first imputation.

3) Add $Y^1(12)$ to $n_1 = 29$, add $Y^1(13)$ to $n_2 = 31$, ..., add $Y^1(23)$ to $n_{11} = 41$

4) In table "sampled IY", denote the sum of entries in columns 12- 23 by $IY^1(12) - IY^1(23)$

5) add $IY^1(12)$ to $n_{11} = 20$, add $IY^1(13)$ to $n_{21} = 24$, ..., and add $IY^1(23)$ to $n_{11,1} = 38$.

6) Compute $U_8^1(t_v)$, $1 \leq v \leq 11$ using the formula above Table 6a.

7) Add $U_8^1(t_v)$, $1 \leq v \leq 11$ to get $U_8^1 = \sum_{v=1}^{11} U_8^1(t_v)$

8) Repeat above for 99 more imputations to have $U_8^1, U_8^2, \dots, U_8^{100}$. The unstandardized test statistic is

$$U_8^I = \frac{\sum_{i=1}^{100} U_8^i}{100},$$

where superscript I says that the statistic is based on imputation.