

Adapting TP statistic to DIC model for Aids data

In DIC model the only choice for states j, l , and m , is $j = 1, l = 2$, and $m = 3$. The only null hypothesis for testing the Markov property against general alternative is tiis

$$H_0 : \alpha_{23}(t|X(s) = 1) = \alpha_{23}(t|X(s) = 2), t > s.$$

Proper and improper MI rectangles

We divide MI rectangles obtained from M observation rectangles into two subsets according to whether the time of 2 to 3 transition is finite or is right censored in state 2:

$$\{R_u \equiv (l_u, r_u] \times (w_u, z_u], z_u < \infty, 1 \leq u \leq U_1\}$$

and

$$R_u \equiv (l_u, r_u] \times (w_u, \infty], 1 \leq u \leq U_2\}.$$

We refer to an MI rectangle in the first subset as proper and in the second as improper. Similarly an observation rectangle (OR) is proper or improper depending on whether its time of 2 to 3 transition is finite or right censored in state 2

1. Reduction step

If $(l_u, r_u] \times (w_u, z_u]$ is the MI rectangle for a proper OR, we reduce it to the proper pair (r_u, z_u) , thus assuming that 1 to 2 and 2 to 3 transitions occur at the end points of their respective intervals.

If $(l_u, r_u] \times (w_u, \infty]$ is the MI rectangle for an improper OR we reduce it to the improper pair (r_u, w_u) thus assuming that 1 to 2 transition occurs at the endpoint of its interval and keep the censoring time w_u in state 2 as is.

2. Specializing to Aids data. This dataset has $M = 96$ ORs of which $M_p = 27$ are proper and $M_{ip} = 69$ are improper rectangles. There are 25 MI rectangles, of which $m_p = 17$ are proper and $m_{ip} = 8$ are improper. The 17 proper MI rectangles are listed in Table 1 below and 8 improper MI rectangles are listed in Table 4 below. Among 17 proper MI rectangles, there are $V = 11$ unique times of the $2 \rightarrow 3$ transition. We denote these times by $t_1 < t_2 < \dots < t_V$. They are listed in Table 3.

The Aids data are such that each proper OR rectangle contains only one of 17 proper MI rectangles. For each proper OR we have to know which proper MI it contains, see Table 2 below.

In Aids data, each improper OR is of the form

$$(L_m, R_m] \times (23, \infty],$$

where $(L_m, R_m]$ is an interval in which $1 \rightarrow 2$ transition occurs after which a patient is right censored in state 2 by time 23, which is the end of study time. In particular if improper OR is of the form

$$(L_m, L_m + 1] \times (23, \infty] \quad (1)$$

it coincides with one of the improper MI rectangles. There are 30 improper ORs that satisfy this condition, see Table 5. There are 9 other improper ORs which contain only one improper MI rectangle, see Table 6. We refer to the improper OR which contains only one improper MI as type 1 improper OR. We refer to the remaining $69 - 39 = 30$ improper ORs which contain more than one improper MI rectangle as type 2 improper ORs.

Computation of the TP statistic for all observations using $s = 8$.

The logrank statistic is

$$\begin{aligned} \hat{U}_8 &= \sum_{v=1}^{11} U_8(t_v) \\ &= \sum_{v=1}^{11} \left\{ \sum_{i=1}^{27} I(X(8) = 1) dN_i(t_v) - \left(\sum_{i=1}^{96} I(X(8) = 1) Y_i(t_v) \right) \frac{\sum_{i=1}^{27} dN_i(t_v)}{\sum_{i=1}^{96} Y_i(t_v)} \right\}, \end{aligned} \quad (2)$$

where the integral with respect to t was replaced by the sum over t'_v s. $N(t)$ is a counting process for $2 \rightarrow 3$ transition by time t , and $Y(t)$ is the number at risk for 2 to 3 transition at time t -. Here $s = 8$ was chosen so that for some proper reduced observations, $X(8) = 1$, and for the remaining reduced observations, $X(8) = 2$. Otherwise, (2) would be equal to zero.

For a given t_v

$$\sum_{i=1}^{27} I(X(8) = 1) dN_i(t_v) = \sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v), \quad (3)$$

the number of subjects who were in state 1 at time 8 and made $2 \rightarrow 3$ transition at time t_v , while

$$\sum_{i=1}^{27} dN_i(t_v) \quad (4)$$

is the total number of subjects who made $2 \rightarrow 3$ transition at time t_v . Next

$$\begin{aligned} \sum_{i=1}^{96} Y_i(t_v) &= \sum_{i=1}^{27} I(r_i < t_v \leq z_i) + \sum_{i=28}^{66} I(r_i < t_v) \\ &+ \sum_{i=67}^{96} [I(R_i < t_v) + I(R_i \geq t_v) \frac{\sum_{k=1}^{K_i} I(r_{i,k} < t_v) P(MI_{i,k})}{P(OR_i)}] \end{aligned}$$

is the total number of patients who are at risk for $2 \rightarrow 3$ transition at time t_v . The first part of above expression is the contribution from 27 proper ORs to the risk set at time t_v , the second is the contribution of 39 type 1 improper ORs, and the last one is the contribution of type 2 improper ORs, where K_i is the number of improper MI rectangles contained in type 2 improper OR_i and $r_{i,k}$ is the right endpoint of the k 'th improper MI rectangle belonging to improper OR_i .

Finally, the number of patients at risk of the $2 \rightarrow 3$ transition at t_v and who were in state 1 at time $s = 8$ is

$$\begin{aligned} \sum_{i=1}^{96} I(8 \leq r_i) Y_i(t_v) &= \sum_{i=1}^{27} I(8 \leq r_i < t_v \leq z_i) \\ &+ \sum_{i=28}^{66} [I(8 \leq r_i < t_v)] \quad (5) \\ &+ \sum_{i=67}^{96} [I(8 \leq R_i < t_v) + I(R_i \geq t_v) \frac{\sum_{k=1}^{K_i} I(8 \leq r_{i,k} < t_v) P(MI_{i,k})}{P(OR_i)}]. \end{aligned}$$

The four quantities (3)-(5) are computed using the information in Tables 1-10 in Section 4.

3 The standardized test statistic

Under H_0 , the standardized $\hat{U}_8$, namely

$$\tilde{U}_8 = \frac{\hat{U}_8}{\sqrt{V(\hat{U}_8)}},$$

has asymptotically standard normal distribution, see Titman and Putter (2022), who also give the variance of $\hat{U}_8$ as

$$V(\hat{U}_8) = \sum_{v:t_v \geq 8} \frac{n_{v1}(n_v - n_{v1})}{(n_v)^2}, \quad (6)$$

where

$$n_v = \sum_{i=1}^{96} Y_i(t_v), n_{v1} = \sum_{i=1}^{96} I(8 \leq r_i) Y_i(t_v).$$

4. Data

Table 1 Proper MI rectangles with their probabilities

	Proper MI	p
1	$(5, 7) \times (11, 12)$	0.0208
2	$(5, 7) \times (13, 13)$	0.0208
3	$(1, 7) \times (16, 16)$	0.0104
4	$(3, 7) \times (17, 17)$	0.0104
5	$(7, 9) \times (19, 19)$	0.0104
6	$(9, 10) \times (15, 15)$	0.0156
7	$(10, 11) \times (14, 15)$	0.0156
8	$(10, 11) \times (15, 16)$	0.0312
9	$(9, 11) \times (17, 18)$	0.0312
10	$(10, 12) \times (16, 17)$	0.0104
11	$(9, 12) \times (22, 22)$	0.0104
12	$(10, 12) \times (23, 23)$	0.0104
13	$(12, 13) \times (20, 20)$	0.0104
14	$(13, 14) \times (17, 18)$	0.0312
15	$(13, 14) \times (19, 20)$	0.0104
16	$(13, 14) \times (20, 21)$	0.0208
17	$(14, 15) \times (23, 23)$	0.0104

Table 2 Proper OR(first column), with its proper MI rectangle(2nd column) and in reduced form (third column) There are 27 proper ORs listed below.

Proper OR	Proper MI	Reduced (r,z)
1(5, 7) $\times$ (11, 12)	(5, 7) $\times$ (11, 12)	(7, 12)
2(1, 10) $\times$ (11, 12)	(5, 7) $\times$ (11, 12)	(7, 12)
3(5, 8) $\times$ (13, 13)	(5, 7) $\times$ (13, 13)	(7, 13)
4(1, 7) $\times$ (12, 13)	(5, 7) $\times$ (13, 13)	(7, 13)
5(9, 13) $\times$ (14, 15)	(10, 11) $\times$ (14, 15)	(11, 15)
6(8, 10) $\times$ (15, 15)	(9, 10) $\times$ (15, 15)	(10, 15)
7(10, 11) $\times$ (14, 15)	(10, 11) $\times$ (14, 15)	(11, 15)
8(10, 12) $\times$ (15, 16)	(10, 11) $\times$ (15, 16)	(11, 16)
9(10, 14) $\times$ (15, 16)	(10, 11) $\times$ (15, 16)	(11, 16)
10(1, 7) $\times$ (16, 16)	(1, 7) $\times$ (16, 16)	(7, 16)
11(10, 11) $\times$ (15, 16)	(10, 11) $\times$ (15, 16)	(11, 16)
12(10, 12) $\times$ (16, 17)	(10, 12) $\times$ (16, 17)	(12, 17)
13(3, 7) $\times$ (17, 17)	(3, 7) $\times$ (17, 17)	(7, 17)
14(9, 13) $\times$ (17, 18)	(9, 11) $\times$ (17, 18)	(11, 18)
15(13, 14) $\times$ (17, 18)	(13, 14) $\times$ (17, 18)	(14, 18)
16(13, 15) $\times$ (17, 18)	(13, 14) $\times$ (17, 18)	(14, 18)
17(9, 12) $\times$ (17, 18)	(9, 11) $\times$ (17, 18)	(11, 18)
18(13, 15) $\times$ (17, 18)	(13, 14) $\times$ (17, 18)	(14, 18)
19(9, 11) $\times$ (17, 18)	(9, 11) $\times$ (17, 18)	(11, 18)
20(7, 9) $\times$ (19, 19)	(7, 9) $\times$ (19, 19)	(9, 19)
21(12, 13) $\times$ (20, 20)	(12, 13) $\times$ (20, 20)	(13, 20)
22(13, 14) $\times$ (19, 20)	(13, 14) $\times$ (19, 20)	(14, 20)
23(1, 15) $\times$ (20, 21)	(13, 14) $\times$ (20, 21)	(14, 21)
24 (13, 14) $\times$ (20, 21)	(13, 14) $\times$ (20, 21)	(14, 21)
25(9, 12) $\times$ (22, 22)	(9, 12) $\times$ (22, 22)	(12, 22)
26(10, 12) $\times$ (23, 23)	(10, 12) $\times$ (23, 23)	(12, 23)
27(14, 15) $\times$ (23, 23)	(14, 15) $\times$ (23, 23)	(15, 23)

Table 3 (For proper ORs) First column lists 11 unique times of $2 \rightarrow 3$ transition, second the number of patients that made the $2 \rightarrow 3$ transition at t_v and were in state 1 at $s = 8$, third, the number of patients who are at risk for $2 \rightarrow 3$ at t_v , the last one gives the number of patients at risk at t_v who were in state 1 at $s=8$.

t_v	$\sum_{i=1}^{27} dN_i(t_v)$	$\sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v)$	$\sum_{i=1}^{27} Y_i(t_v)$	$\sum_{i=1}^{27} I(8 \leq r_i) Y_i(t_v)$
12	2	0	16	10
13	2	0	17	13
15	3	3	22	20
16	4	3	20	18
17	2	1	16	15
18	6	6	14	14
19	1	1	8	8
20	2	2	7	7
21	2	2	5	5
22	1	1	3	3
23	2	2	2	2

Note that multiplicities (entries in second column) add up to 27, which is the number of proper ORs. The entries in in Table 3 are obtained from Table 2 in the following way.

$$\begin{aligned}
\sum_{i=1}^{27} dN_i(t_v) &= \sum_{i=1}^{27} I(z_i = t_v), \\
\sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v) &= \sum_{i=1}^{27} I(8 \leq r_i < z_i = t_v), \\
\sum_{i=1}^{27} Y_i(t_v) &= \sum_{i=1}^{27} I(z_i \geq t_v), \\
\sum_{i=1}^{27} I(8 \leq r_i) Y_i(t_v) &= \sum_{i=1}^{27} I(8 \leq r_i, z_i \geq t_v).
\end{aligned}$$

Table 4. Improper MI rectangles with their probabilities and in reduced form

	Improper MI rectangle (l,r)	probability	reduced (r, 23)
1	$(5, 7] \times (23, \infty)$	0.0437	(7, 23)
2	$(8, 9] \times (23, \infty)$	0.0235	(9, 23)
3	$(9, 10] \times (23, \infty)$	0.0639	(10, 23)
4	$(10, 11] \times (23, \infty)$	0.1649	(11, 23)
5	$(11, 12] \times (23, \infty)$	0.0379	(12, 23)
6	$(12, 13] \times (23, \infty)$	0.1682	(13, 23)
7	$(14, 15] \times (23, \infty)$	0.1548	(15, 23)
8	$(15, 16] \times (23, \infty)$	0.0619	(16, 23)

Table 5. Type 1 improper ORs which are of the form $OR_m = (L_m, L_m + 1] \times (23, \infty]$ with their probabilities and in reduced form.

Improper OR=Improper MI	multiplicity	reduced (r, 23)	p
$(9, 10) \times (23, \infty)$	2	(10, 23)	0.0639
$(10, 11) \times (23, \infty)$	7	(11, 23)	0.1649
$(11, 12) \times (23, \infty)$	1	(12, 23)	0.0379
$(12, 13) \times (23, \infty)$	7	(13, 23)	0.1682
$(14, 15) \times (23, \infty)$	8	(15, 23)	0.1548
$(15, 16) \times (23, \infty)$	5	(16, 23)	0.0619
total	30		

In "reduced" column, the entry (10, 23) means that a patient made $1 \rightarrow 2$ transition at $r = 10$ and stayed in state 2 until the end of the study denoted by 23

Table 6. Type 1 improper ORs which are not of the form $OR_m = (L_m, L_m + 1] \times (23, \infty]$ with their probabilities and in reduced form.

Improper OR	Improper MI	multiplicity	reduced $(r, 23)$
$(1, 7) \times (23, \infty)$	$(5, 7) \times (23, \infty)$	1	$(7, 23)$
$(5, 7) \times (23, \infty)$	$(5, 7) \times (23, \infty)$	2	$(7, 23)$
$(7, 9) \times (23, \infty)$	$(8, 9) \times (23, \infty)$	1	$(9, 23)$
$(12, 14) \times (23, \infty)$	$(12, 13) \times (23, \infty)$	2	$(13, 23)$
$(13, 15) \times (23, \infty)$	$(14, 15) \times (23, \infty)$	3	$(15, 23)$
total		9	

We now compute the test statistic based on 66 ORs which contain only one MI rectangle assuming that $s = 8$. This statistic is

$$\hat{U}_8 = \sum_{v=1}^{11} U_8(t_v),$$

where

$$U_8(t_v) = \sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v) - \left[\sum_{i=1}^{66} I(8 \leq r_i) Y_i(t_v) \right] \frac{\sum_{i=1}^{27} dN_i(t_v)}{\sum_{i=1}^{66} Y_i(t_v)}.$$

and is to be computed based on **Table 6a** below.

Table 6a The first two columns come from Table 3. Columns 3 and 4 use information from the same columns in Table 3 (first number). The second number is obtained. by inspection of Tables 5 and 6.

t_v	$\sum_{i=1}^{27} dN_i(t_v)$	$\sum_{i=1}^{27} I(8 \leq r_i) dN_i(t_v)$	$n_v = \sum_{i=1}^{66} Y_i(t_v)$	$n_{v1} = \sum_{i=1}^{66} I(8 \leq r_i) Y_i(t_v)$	$U_8(t_v)$
12	2	0	16 + 13 = 29	10 + 10 = 20	-1.37931
13	2	0	17 + 14 = 31	13 + 11 = 24	-1.54839
15	3	3	22 + 23 = 45	20 + 20 = 40	0.33333
16	4	3	20 + 34 = 54	18 + 31 = 49	-0.62963
17	2	1	16 + 39 = 55	15 + 36 = 51	-0.85454
18	6	6	14 + 39 = 53	14 + 36 = 50	0.33962
19	1	1	8 + 39 = 47	8 + 36 = 44	0.06383
20	2	2	7 + 39 = 46	7 + 36 = 43	0.13043
21	2	2	5 + 39 = 44	5 + 36 = 41	0.13636
22	1	1	3 + 39 = 42	3 + 36 = 39	0.07143
23	2	2	2 + 39 = 41	2 + 36 = 38	0.14634

Adding up the entries in the last column gives $\hat{U}_8 = -3.19052$

To compute the variance of the test statistic $\hat{U}_8$ we use **Table 6**.

Table 6b The second and third columns are defined in **Table 6a**. The last column is obtained from expression for variance of the test statistic $\hat{U}_8$ given in section 3 by (6).

t_v	n_v	n_{v1}	$n_{v1}(n_v - n_{v1})/n_v^2$
12	29	20	0.214031
13	31	24	0.174818
15	45	40	0.0987654
16	54	49	0.0840192
17	55	51	0.0674380
18	53	50	0.0533998
19	47	44	0.0597555
20	46	43	0.0609641
21	44	41	0.0635331
22	42	39	0.0663265
23	41	38	0.0678167

Adding up the entries in the last column gives $V(\hat{U}_8) = 1.01087$. Then the standardized test statistic is

$$\frac{\hat{U}_8}{\sqrt{V(\hat{U}_8)}} = \frac{-3.19052}{\sqrt{1.01087}} = -3.1733,$$

which leads to the rejection of null hypothesis.

We next include type 2 improper ORs for the computation of the test statistic (there are such 30 such rectangles and they contain more than one improper MI rectangle). These ORs are listed in Table 7

Table 7 The 30 type 2 improper ORs together with their multiplicity and probabilities. The probabilities are obtained from Table 4 by adding the probabilities of improper MI rectangles that belong to a given improper OR as shown in the table. There are 18 different improper ORs in the table. The

multiplicity of each OR is shown in column 2.

Imp OR (L,R]	multiplicity	Total probability of improper OR of type 2
1 (1, 11]	1	$0.0437 + 0.0235 + 0.0639 + 0.1649 = 0.296$
2 (1, 12]	2	$0.0437 + 0.0235 + 0.0639 + 0.1649 + 0.0379 = 0.3339$
3 (1, 13)	2	$0.3339 + 0.1682 = 0.5021$
4 (1, 14)	3	0.5021
5 (1, 15)	2	$0.5021 + 0.1548 = 0.6569$
6 (1, 16)	1	$0.6569 + 0.0619 = 0.7188$
7 (3, 14)	1	0.5021
8 (3, 15)	1	0.6569
9 (7, 10)	1	$0.0235 + 0.0639 = 0.0874$
10 (7, 15)	1(15)	$0.0874 + 0.1649 + 0.0379 + 0.1682 + 0.1548 = 0.6132$
11(8, 10)	1	$0.0235 + 0.0639 = 0.0874$
12(8, 15)	1	0.6132
13 (14, 16)	2	$0.1548 + 0.0619 = 0.2167$
14 (9, 12)	3	$0.0639 + 0.1649 + 0.0379 = 0.2667$
15(10, 12)	2	$0.2667 - 0.0639 = 0.2028$
16(10, 15)	1	$0.2028 + 0.1682 + 0.1548 = 0.5258$
17(11, 13)	4	$0.0379 + 0.1682 = 0.2061$
18(13, 16)	1	$0.1548 + 0.0619 = 0.2167$

. The contribution of the observations in Table 7 to the computation of the test statistic. The observation may contribute "1" to the risk at time t_v . (column 3 in Table 6a) and if it does it may also contribute to column 4 in Table 6a. These contributions are obtained through imputation.

Imputation

The goal of an imputation is to assess if the i 'th type 2 improper OR contributes to $\sum_{i=67}^{99} Y_i(t_v)$ and/or to $\sum_{i=67}^{99} I(8 < r_i)Y_i(t_v)$.

Consider the first OR (OR1) in Table 7. namely $(1, 11]$. Going back to Table 4, we see that OR1 contains four MI rectangles

$$\text{OR1} = (1, 11] = (5, 7] \cup (8, 9] \cup (9, 10] \cup (10, 11].$$

Table **I1a**: for $Y_1(t_v)$, $1 \leq v \leq 11$ shows that for each MI rectangle $Y_1(v) = 1$ for $1 \leq v \leq 11$ so imputation is not needed.

outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.147635	1	1	1	1	1	1	1	1	1	1	1
(8, 9]	0.079392	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.215878	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.557095	1	1	1	1	1	1	1	1	1	1	1

Table **I1b** for $I(8 \leq r_i)Y_1(v)$ shows the contribution of the first OR to the modified risk set at t_v , $1 \leq v \leq 11$.

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.147635	0	0	0	0	0	0	0	0	0	0	0
(8, 9]	0.079392	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.215878	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.557095	1	1	1	1	1	1	1	1	1	1	1

Table I1 The final table to be used in the simulation. Here 0 stands for $(5, 7]$ and 1 for the remaining MI's. It indicates that we need to draw from the distribution with values in column 1, probabilities in column 2. The contribution to the modified risk and risk sets is described in columns 3 and 4 respectively..

draw	probability	$I(8 \leq r_i)Y_1(t_v)$	$Y_1(t_v)$
0	0.147635	$0, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$,
1	0.852465	$1, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$

OR2=(1, 11] = (5, 7] $\cup$ (8, 9] $\cup$ (9, 10] $\cup$ (10, 11] $\cup$ (11, 12] **There are two OR2s.**

Table I2a for $Y_2(v)$, ($1 \leq v \leq 11$)

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.130878	1	1	1	1	1	1	1	1	1	1	1
(8, 9]	0.070380	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.191375	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.493860	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.113507	0	1	1	1	1	1	1	1	1	1	1

Table I2a indicates that you have to draw from the distribution with two values as described in Table I2a. below Column 1 In Table I2a states two values. column 2 adjusted probabilities and column 3. describes the contribution to the risk set at t_v , ($1 \leq v \leq 11$) of OR2

Table I2b for $I(8 \leq r_i)Y_2(v)$, ($1 \leq v \leq 11$)

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.130878	0	0	0	0	0	0	0	0	0	0	0
(8, 9]	0.070380	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.191375	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.493860	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.113507	0	1	1	1	1	1	1	1	1	1	1

Combining information from Tables I2a and Table I2b we get

Table I2: 0 represents (5,7], 2 represents (11,12] and 1 remaining MI's

draw	AdjProb	$I(8 \leq r_i)Y_2(v)$	$Y_2(v)$
0	0.130878	0, ($1 \leq v \leq 11$)	1, ($1 \leq v \leq 11$)
1		1, ($1 \leq v \leq 11$)	1, ($1 \leq v \leq 11$)
2	0.113507	$\left\{ \begin{array}{l} 0, (v=1) \\ 1, (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, (v=1) \\ 1, (2 \leq v \leq 11) \end{array} \right\}$

We have to draw **twice** from the distribution in Table I2 because there are two ORs 2 to also get $Y_3(v)$ and $I(8 \leq r_i)Y_3(v)$, ($1 \leq v \leq 11$).

$\text{OR3}=(1,13] = (5, 7] \cup (8, 9] \cup (9, 10] \cup (10, 11] \cup (11, 12] \cup (12, 13]$ **There are two OR3s**

Table I3a for $Y_4(v)$ and $Y_5(v)$, $(1 \leq v \leq 11)$

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.0870345	1	1	1	1	1	1	1	1	1	1	1
(8, 9]	0.0468034	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.127265	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.328421	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.075483	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.334993	0	0	1	1	1	1	1	1	1	1	1

We build Table I3a to show the contribution of the 4th (5th) OR to risk sets at times t_v $(1 \leq v \leq 11)$ which depends on the chosen MI.

Table I3b for $I(8 \leq r_i)Y_4(v)$ and $I(8 \leq r_i)Y_5(v)$, $(1 \leq v \leq 11)$

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.0870345	0	0	0	0	0	0	0	0	0	0	0
(8, 9]	0.0468034	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.127265	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.328421	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.075483	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.334993	0	0	1	1	1	1	1	1	1	1	1

Table I3 represents (5,7] by 0, (11, 12] by 2 and (12, 13] by 3 and remaining MI's by 1

draw	Prob	$Y_4(v)$	$I(8 \leq r_i)Y_4(v)$
0	0.0870345	1 $(1 \leq v \leq 11)$	0 $(1 \leq v \leq 11)$
1		1 $(1 \leq v \leq 11)$	1 $(1 \leq v \leq 11)$
2	0.075483	$\left\{ \begin{array}{l} 0 (v = 1) \\ 1 (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0 (v = 1) \\ 1 (2 \leq v \leq 11) \end{array} \right\}$
3	0.334993	$\left\{ \begin{array}{l} 0 (v = 1, 2) \\ 1 (3 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0 (v = 1, 2) \\ 1 (3 \leq v \leq 11) \end{array} \right\}$

We have to draw **twice** from the distribution in Table I3 because there are two ORs 3 to also get $Y_5(v)$ and $I(8 \leq r_i)Y_5(v)$, $(1 \leq v \leq 11)$

OR4=(1,14]=(5, 7] $\cup$ (8, 9] $\cup$ (9, 10] $\cup$ (10, 11] $\cup$ (11, 12] $\cup$ (12, 13] **There are three OR4s**

Table **I4**=Table**I3**.

Since there are three OR4s we need to draw 3 times from the distribution in Table I3 to get $Y_j(v)$ and $I(8 \leq r_i)Y_j(v)$, ($1 \leq v \leq 11$) where $6 \leq j \leq 8$.

Table I3 represents (5,7] by 0, (11, 12] by 2 and (12, 13] by 3 and remaining MI's by 1

draw	Prob	$Y_4(v)$	$I(8 \leq r_i)Y_4(v)$
0	0.0870345	1 ($1 \leq v \leq 11$)	0 ($1 \leq v \leq 11$)
1		1 ($1 \leq v \leq 11$)	1 ($1 \leq v \leq 11$)
2	0.075483	$\left\{ \begin{array}{l} 0 \ (v = 1) \\ 1 \ (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0 \ (v = 1) \\ 1 \ (2 \leq v \leq 11) \end{array} \right\}$
3	0.334993	$\left\{ \begin{array}{l} 0 \ (v = 1, 2) \\ 1 \ (3 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0 \ (v = 1, 2) \\ 1 \ (3 \leq v \leq 11) \end{array} \right\}$

$$\text{OR5}=(1,15]=(5,7] \cup (8,9] \cup (9,10] \cup (10,11] \cup (11,12] \cup (12,13] \cup (14,15]$$

There are two OR5

Table **I5a** for $Y_9(v)$ and $Y_{10}(v)$

outcome	prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.066525	1	1	1	1	1	1	1	1	1	1	1
(8, 9]	0.035774	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.097275	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.251028	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.057695	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.256051	0	0	1	1	1	1	1	1	1	1	1
(14, 15]	0.235652	0	0	0	1	1	1	1	1	1	1	1

Table **I5b** for $I(8 \leq r_i)Y_9(v)$ and $I(8 \leq r_i)Y_{10}(v)$

outcome	prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.066525	0	0	0	0	0	0	0	0	0	0	0
(8, 9]	0.035774	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.097275	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.251028	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.057695	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.256051	0	0	1	1	1	1	1	1	1	1	1
(14, 15]	0.235652	0	0	0	1	1	1	1	1	1	1	1

Combining we get

Table I5

draw	prob	$Y_9(t_v)$	$I(8 \leq r_i)Y_9(v)$
0	0.066525	$1, (1 \leq v \leq 11)$	$0, (1 \leq v \leq 11)$
1		$1, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$
2	0.057695	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
3	0.256051	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
4	0.235652	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$

$\text{OR6} = (1,16] = (5,7] \cup (8,9] \cup (9,10] \cup (10,11] \cup (11,12] \cup (12,13] \cup (14,15] \cup (15,16]$ There is one OR6.

Table **I6a**

outcome	prob	12	13	15	16	17	18	19	20	21	22	23
(5, 7]	0.060796	1	1	1	1	1	1	1	1	1	1	1
(8, 9]	0.032693	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.088898	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.22941	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.052727	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.234001	0	0	1	1	1	1	1	1	1	1	1
(14, 15]	0.215359	0	0	0	1	1	1	1	1	1	1	1
(15, 16]	0.086116	0	0	0	0	1	1	1	1	1	1	1

Table I6b is the same as table I6a but with the first row changed to zeros

Table I6) 0 represents (5,7], 2 (11,12], 3 (12,13], 4 (14,15], 5 (15,16] and 1 the remaining MIs

draw	adjProb	$Y_{11}(v)$	$I(8 \leq r_i)Y_{11}(v)$
0	0.060796	$1, (1 \leq v \leq 11)$	$0, (1 \leq v \leq 11)$
1		$1, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$
2	0.052727	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
3	0.234001	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
4	0.215359	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$
5	0.086116	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 4) \\ 1, (5 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 4) \\ 1, (5 \leq v \leq 11) \end{array} \right\}$

$\text{OR7}=(3,14]=(5,7] \cup (8,9] \cup (9,10] \cup (10,11] \cup (11,12] \cup (12,13]$ There is one **OR7, which is the 12th observation**

Table **I7**=Table **I4**=Table**I3**

$\text{OR8}=(3,15]=(5,7] \cup (8,9] \cup (9,10] \cup (10,11] \cup (11,12] \cup (12,13] \cup (14,15]$.

There is one **OR8, which is the 13th observation**

Table I8=Table I5

Table I5

draw	prob	$Y_{13}(t_v)$	$I(8 \leq r_i)Y_{13}(v)$
0	0.066525	$1, (1 \leq v \leq 11)$	$0, (1 \leq v \leq 11)$
1		$1, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$
2	0.057695	$\left\{ \begin{array}{c} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{c} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
3	0.256051	$\left\{ \begin{array}{c} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{c} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
4	0.235652	$\left\{ \begin{array}{c} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$	$\left\{ \begin{array}{c} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$

OR**9**=(7, 10) = (8, 9] $\cup$ (9, 10) There is one OR9, which the 14th observation

Table **I9a=TableI9b**

outcome		12	13	15	16	17	18	19	20	21	22	23
(8, 9]	0.268879	1	1	1	1	1	1	1	1	1	1	1
(9, 10)	0.731121	1	1	1	1	1	1	1	1	1	1	1

We do not need to draw.

Table I9

$Y_{14}(v)$	$I(8 \leq r_i)Y_{14}(v)$
$1, (1 \leq v \leq 11)$	$1, (1 \leq v \leq 11)$

OR10=(7, 15] = (8, 9]∪(9, 10] ∪ (10, 11] ∪ (11, 12] ∪ (12, 13] ∪ (14, 15]
Table I10a=Table IbTable I10

outcome	Prob	12	13	15	16	17	18	19	20	21	22	23
(8, 9]	0.038324	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.104207	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.268917	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.061807	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.274299	0	0	1	1	1	1	1	1	1	1	1
(14, 15]	0.252446	0	0	0	1	1	1	1	1	1	1	1

Table I10

outcome	AdjProb	$Y_{15}(v) = I(8 \leq r_i)Y_{15}(v)$
0		$1, 1 \leq v \leq 11$
1	0.061807	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
2	0.274299	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
3	0.252446	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$

$$\text{OR}\mathbf{11} = (8, 10] = (8, 9] \cup (9, 10]$$

Table **I11**

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(8, 9]	0.268879	1	1	1	1	1	1	1	1	1	1	1
(9, 10]	0.731121	1	1	1	1	1	1	1	1	1	1	1

We do not need to draw

Table **I9**

$$\begin{array}{cc} Y_{16}(v) & I(8 \leq r_i)Y_{16}(v) \\ 1, (1 \leq v \leq 11) & 1, (1 \leq v \leq 11) \end{array}$$

OR12 = $(8, 15] = (8, 9] \cup (9, 10] \cup (10, 11] \cup (11, 12] \cup (12, 13] \cup (14, 15]$

Table **I12** = Table **I10**.

Table I10

outcome	AdjProb	$Y_{17}(v) = I(8 \leq r_i)Y_{17}(v)$
0		$1, 1 \leq v \leq 11$
1	0.061807	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
2	0.274299	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
3	0.252446	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$

OR**13**=(14, 16] = (14, 15] $\cup$ (15, 16] There are two OR13s
Table **I13**

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(14, 15]	0.714352	0	0	0	1	1	1	1	1	1	1	1
(15, 16]	0.285648	0	0	0	0	1	1	1	1	1	1	1

Outcome	CProb	$Y_{18}(v) = I(8 \leq r_i)Y_{18}(v)$
(14, 15]	0.714352	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$
(15, 16]	0.714352	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 4) \\ 1, (5 \leq v \leq 11) \end{array} \right\}$

and the same for observations 19 and 20.

OR14=(9,12]=(9, 10] $\cup$ (10, 11] $\cup$ (11, 12]

Table I14

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(9, 10]	0.239595	1	1	1	1	1	1	1	1	1	1	1
(10, 11]	0.618298	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.142107	0	1	1	1	1	1	1	1	1	1	1

Draw	AdjProb	$Y(v) = I(8 \leq r_i)Y(v)$
0	0.857893	$1, (1 \leq v \leq 11)$
1	0.142107	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$

OR**15**=(10, 12] = (10, 11] $\cup$ (11, 12]

Table **I15**

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(10, 11]	0.813116	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.186884	0	1	1	1	1	1	1	1	1	1	1

Draw	CProb	$Y_{20}(v) = I(8 \leq r_i)Y_{20}(v)$
(10, 11]	0.813116	$1, (1 \leq v \leq 11)$
(11, 12]	0.186884	$\left\{ \begin{array}{c} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$

Draw	CProb	$Y_{21}(v) = I(8 \leq r_i)Y_{21}(v)$
(10, 11]	0.813116	$1, (1 \leq v \leq 11)$
(11, 12]	0.186884	$\left\{ \begin{array}{c} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$

OR**16**=(10,15]=(10, 11] $\cup$ (11, 12] $\cup$ (12, 13] $\cup$ (14, 15]

Table **I16**

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(10, 11]	0.313617	1	1	1	1	1	1	1	1	1	1	1
(11, 12]	0.072081	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.319893	0	0	1	1	1	1	1	1	1	1	1
(14, 15]	0.294409	0	0	0	1	1	1	1	1	1	1	1

draw	CProb	$Y(v) = I(8 \leq r_i)Y(v)$
(10, 11]	0.313617	$1, (1 \leq v \leq 11)$
(11, 12]	0.072081	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
(12, 13]	0.319893	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$
(14, 15]	0.294409	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$

OR**17**=(11, 13] = (11, 12] $\cup$ (12, 13]

Table **I17**

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(11, 12]	0.183891	0	1	1	1	1	1	1	1	1	1	1
(12, 13]	0.816109	0	0	1	1	1	1	1	1	1	1	1

<i>Draw</i>	CProb	$Y(v) = I(8 \leq r_i)Y(v)$
(11, 12]	0.183891	$\left\{ \begin{array}{l} 0, v = 1 \\ 1, (2 \leq v \leq 11) \end{array} \right\}$
(12, 13]	0.816109	$\left\{ \begin{array}{l} 0, v = 1, 2 \\ 1, (3 \leq v \leq 11) \end{array} \right\}$

OR**18**=(13, 16] = (14, 15] $\cup$ (15, 16]

Table I18

Outcome	CProb	12	13	15	16	17	18	19	20	21	22	23
(14, 15]	0.714352	0	0	0	1	1	1	1	1	1	1	1
(15, 16]	0.285648	0	0	0	0	1	1	1	1	1	1	1

Draw	CProb	$Y(t_v) = I(8 \leq r_i)Y(t_v)$
(14, 15]	0.714352	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 3) \\ 1, (4 \leq v \leq 11) \end{array} \right\}$
(15, 16]	0.285648	$\left\{ \begin{array}{l} 0, (1 \leq v \leq 4) \\ 1, (5 \leq v \leq 11) \end{array} \right\}$