



Review

Gene-regulatory networks controlling inflorescence and flower development in *Arabidopsis thaliana**



Christopher Ralf Wils, Kerstin Kaufmann*

Institute for Biochemistry and Biology, University of Potsdam, 14476 Potsdam, Germany

ARTICLE INFO

Article history:

Received 22 April 2016

Received in revised form 21 July 2016

Accepted 22 July 2016

Available online 31 July 2016

Keywords:

Inflorescence

Flower development

Floral meristem

Gene-regulatory network

Transcription factor

ABSTRACT

Reproductive development in plants is controlled by complex and intricate gene-regulatory networks of transcription factors. These networks integrate the information from endogenous, hormonal and environmental regulatory pathways. Many of the key players have been identified in *Arabidopsis* and other flowering plant species, and their interactions and molecular modes of action are being elucidated. An emerging theme is that there is extensive crosstalk between different pathways, which can be accomplished at the molecular level by modulation of transcription factor activity or of their downstream targets. In this review, we aim to summarize current knowledge on transcription factors and epigenetic regulators that control basic developmental programs during inflorescence and flower morphogenesis in the model plant *Arabidopsis thaliana*.

This article is part of a Special Issue entitled: Plant Gene Regulatory Mechanisms and Networks, edited by Dr. Erich Grotewold and Dr. Nathan Springer.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Successful reproduction in plants requires the coordinated transition from vegetative growth to reproductive development. The floral

transition in angiosperms is triggered by environmental factors, such as photoperiod, temperature and nutrient conditions, as well as by endogenous cues [1]. The exact environmental conditions that initiate flowering are variable between species and show a high degree of adaptation to local habitats of natural populations (see, e.g. [2–4]). External signals are typically perceived in leaves and roots, and long-distance transport of signaling compounds to the shoot meristem then triggers its switch from vegetative to inflorescence meristem identity [1]. These changes in meristem identities result in alteration of internode growth, branching and the identities of organs that are produced by these meristems. In the course of flower development, the floral meristem becomes terminated and consumed during organ outgrowth.

While floral transition and inflorescence development are highly variable among different species of flowering plants, the generic flower bauplan with distinct perianth, male and female reproductive organ types is evolutionarily conserved. Angiosperm flowers are typically characterized by sterile perianth organs in the outer whorls, and by male (stamens) and female (carpels) reproductive organs in the inner whorls, respectively. The model plant species *Arabidopsis thaliana* (*Arabidopsis*, thale cress) has a differentiated perianth consisting of sepals and petals that is typical for eudicots. The specification of floral organ identities is controlled by floral homeotic genes, which have been identified to encode transcription factors (TFs) about 25 years ago [5–7]. According to the classical ABC model [8] that is based on phenotypes of homeotic mutants in *Arabidopsis* and *Antirrhinum majus*, floral organ identities are specified by 3 functional gene classes that act together in a combinatorial fashion: class A genes specify the identities of sepals, and together with class B genes determine petal identity. Class

Abbreviations: TF, transcription factor; MADS, MCM1, AGAMOUS, DEFICIENS, SRF; SEP, SEPALATA; ChIP-seq, chromatin immunoprecipitation followed by deep sequencing; PcG, Polycomb group; H3K27me3, histone 3 lysine 27 tri-methylation; PRC, polycomb repressive complex; trxG, trithorax group; SAND domain, Sp100, AIRE-1, NucP41/75, DEAF-1 domain; ULT, ULTRAPETALA; miRNA, microRNA; AP2, APETALA2; FLC, FLOWERING LOCUS C; SOC1, SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1; FT, FLOWERING LOCUS T; FD, FLOWERING LOCUS D; IM, inflorescence meristem; AGL24, AGAMOUS-LIKE 24; FM, floral meristem; LFY, LEAFY; AP1, APETALA1; CUC2, CUP-SHAPED COTYLEDON 2; NAC, NAM, ATAF, CUC; BP, BREVIPEDICELLUS; KNAT, KNOTTED1-LIKE HOMEODOMAIN GENE from *Arabidopsis thaliana*; BLH, BEL1-LIKE HOMEODOMAIN; PNY, PENNYWISE; BOP, BLADE-ON-PETIOLE; ATH1, ARABIDOPSIS THALIANA HOMEODOMAIN GENE 1; PNF, POUNDFOOLISH; CAL, CAULIFLOWER; PIN1, PIN-FORMED 1; MP, MONOPTEROS; ARF, AUXIN RESPONSE FACTOR; ANT, AINTEGUMENTA; AIL6, AINTEGUMENTA-LIKE 6; SWI/SNF, SWITCH/SUCROSE NONFERMENTING; LMI, LATE MERISTEM IDENTITY; GA, gibberellic acid; SPL, SQUAMOSA PROMOTER BINDING PROTEIN-LIKE; ELA1, EUI-LIKE P450 A1; TOE, TARGET OF EAT; WUS, WUSCHEL; SVP, SHORT VEGETATIVE PHASE; SEU, SEUSS; AG, AGAMOUS; PI, PISTILLATA; AP3, APETALA 3; TALE-HD, three amino acid loop extension homeodomain; BEL1, BELL1; BRM, BRAHMA; SYD, SPLAYED; UFO, UNUSUAL FLORAL ORGANS; HEC, HECATE; CRC, CRABCLAW; IAA, INDOLE-3-ACETIC ACID INDUCIBLE; GA2OX, GIBBERELLIN 2-OXIDASE; JA, jasmonic acid; GNC, GATA; NITRATE INDUCIBLE, CARBON METABOLISM INVOLVED; GNL, GNC-LIKE; KNU, KNUCKLES; PAN, PERIANTHIA; AGO, ARGONAUTE; SQN, SQUINT; UIF, ULTRAPETALA1 INTERACTING FACTOR1; SUP, SUPERMAN; Cas9, CRISPR-associated protein 9.

* This article is part of a Special Issue entitled: Plant Gene Regulatory Mechanisms and Networks, edited by Dr. Erich Grotewold and Dr. Nathan Springer.

* Corresponding author.

E-mail address: kerstin.kaufmann@uni-potsdam.de (K. Kaufmann).

B genes together with class C genes specify stamens. Class C genes alone determine carpel identity. The model was later extended by class E genes which are important for specification of all types of floral organs [9,10], while the 'A class' is debated and a new definition combining class A and E functions has been proposed as to set a floral 'ground state' [11]. With the exception of the classical A class, the functions of homeotic genes are widely conserved across angiosperms, and some evolutionary changes in basic flower morphology can be explained by modifications in the ABC(E) model [12].

Reproductive meristem specification and organ differentiation require orchestrated changes in expression of many genes. The characterization of the underlying gene-regulatory networks and molecular mechanisms has made tremendous progress in recent years, in particular in model plant species such as *Arabidopsis* and rice. In this review, we attempt to provide an overview on gene-regulatory factors and processes controlling reproductive development in the best studied model plant *Arabidopsis*. We do not attempt to comprehensively cover the floral transition process, which has recently been extensively reviewed elsewhere [1,2,13,14].

2. Gene-regulatory networks: what are the 'players'?

Developmental gene regulation in plants, like in other eukaryotic organisms, is controlled by the interplay of TFs, epigenetic mechanisms and posttranscriptional processes. The *Arabidopsis* genome encodes at least 2300 transcriptional regulators, at least 360 of which have annotated roles in development [15–17]. For example, at least 17 MADS (MCM1, AGAMOUS, DEFICIENS, SRF)-box genes and 23 homeobox TF genes have clear roles in flower development (Table 1). Some TF families appear to have predominant roles in individual processes, such as Myb-domain TFs which have been mainly described to act in stamen development (Table 1). Forward and reverse genetic approaches in the past 25 years have boosted our knowledge on the functions of developmental TF genes. It has also become clear that genetic redundancy is a major confounding factor in elucidating developmental roles of TFs. For example, style and stigma development in the gynoecium is controlled by four largely redundantly acting *NGATHA* genes that belong to the B3 TF family [18,19]. Another example is the four redundantly acting *SEPALLATA* (*SEP1–4*) MADS-box TFs which together control floral meristem and organ specification by constituting the so-called 'E class' [9,10,20]. Classical genetic analyses have described reproductive development as hierarchically organized genetic network with sequentially acting genes. Linked with genome-wide TF target gene identification, it has now become clear that gene-regulatory networks are inherently complex, with multiple feedback and feedforward loops not all of which have been fully understood to date [21]. Genome-wide target gene identification has been enabled by technologies such as chromatin immunoprecipitation followed by deep sequencing (ChIP-seq) or hybridization to tiling arrays (ChIP-chip) [21] together with gene perturbation and genome-wide expression analyses [22].

Next to TFs, also major epigenetic regulators have been shown to act in plant reproductive development. In general, epigenetic regulation is required for cellular memory of gene expression states across cell divisions. Switches in stage- and tissue-specific gene expression programs during development, and developmental plasticity in response to environmental signals require changes in epigenetic states of gene bodies and promoters [21,23,24]. The maintenance of gene repression across cell divisions, termed 'epigenetic memory' is mediated by Polycomb group (PcG) factors, which deposit repressive histone 3 lysine 27 trimethylation (H3K27me3) marks (see [25,26] for recent review). PcG mediated regulation is not limited to flower development, but affects thousands of genes (see, e.g. [27–29]), and multiple mechanisms contribute to the recruitment of PcG proteins to their target gene promoters [30–34]. During vegetative development, floral homeotic genes are epigenetically repressed by PcG protein complexes. Gene regulation by Polycomb Repressive Complex (PRC) 1 and PRC2 is achieved by

depositing H3K27me3 and histone 2A ubiquitination marks at target genes and subsequent chromatin compaction (for review, see [25]). Mutations in PRC components often show decreased H3K27me3 levels and precocious expression of floral homeotic genes in the vegetative phase [35,36]. Proteins that counteract the repressive function of PcG proteins are considered as trithorax group (trxG) factors. Some trxG proteins display histone methyl transferase activity, others assist in that function or are involved in chromatin remodeling (reviewed by [24,26]). For example, the SAND (Sp100, AIRE-1, NucP41/75, DEAF-1) domain proteins ULTRAPETALA (ULT) 1 and ULT2, are considered to function as trxG proteins influencing histone 3 lysine 4 tri-methylation [37–40]. Furthermore ULT1 acts as an anti-repressor, similarly counteracting PcG action on thousands of target genes including floral homeotic genes [41]. ULT1 and ULT2 have dual roles in floral stem cell accumulation and the establishment of apical-basal polarity in the gynoecium [38,40].

Next to TFs and other proteins, the role of regulatory RNAs in the control of developmental gene expression has also been widely appreciated now [42]. Most well-known are microRNAs (miRNAs), which are in their mature state approximately 21-nucleotides in length. MiRNAs originate from stem-loop regions of long primary transcripts and are processed by a Dicer-like enzyme [42,43]. They either function in degradation of complementary target mRNAs or in translational inhibition [43]. An example for a well-known family of miRNAs with roles in flower development is the miR172 family, which regulates several APETALA2 (AP2) class TFs thereby controlling flower initiation, patterning and floral stem cell activity [44–48]. Linked with the availability of large-scale sequencing approaches, long non-coding RNAs have been identified systematically more recently [49,50]. The best known example is the long non-coding antisense RNA *COOLAIR*, which regulates the epigenetic status of the central repressor of flowering, *FLOWERING LOCUS C* (*FLC*) [4,51,52]. However, roles of long non-coding RNAs after the transition to flowering are still largely unexplored.

3. Gene-regulatory networks controlling inflorescence development

Inflorescence development starts with the activation of the MADS-box TF *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1* (*SOC1*) in the shoot apex of *Arabidopsis* [53,54], mainly by the 'florigen' *FLOWERING LOCUS T* (FT) and its interaction partner FD. In the IM, *SOC1* constitutes a major hub in controlling the expression of many other genes involved in floral transition, meristem identity and other functions [55,56]. *SOC1* acts in concert with *AGAMOUS-LIKE 24* (*AGL24*) and *FRUITFULL*, which also encode MADS-domain TFs, and triple mutants show a drastic late-flowering 'perennial' phenotype with floral reversion events and prolonged indeterminate growth [57]. FT/FD furthermore activate the expression of the floral meristem (FM) identity genes *LEAFY* (*LFY*) and *APETALA1* (*API*) gene in the laterally emerging meristems [58–60].

Besides TFs that control IM identity, other transcriptional regulators, in concert with phytohormone signaling pathways, control inflorescence morphology, e.g. by affecting branching or phyllotaxis (reviewed in [61–63]). For example, the boundary gene *CUP-SHAPED COTYLEDON 2* (*CUC2*), which encodes a NAC (NAM, ATAF, CUC) family TF, plays a role in elongation of stem internodes and therefore in establishing the spiral inflorescence phyllotaxis [64]. Inflorescence architecture is furthermore controlled by a network of homeobox TFs. The *KNOTTED1-LIKE HOMEBOX GENE FROM ARABIDOPSIS THALIANA* (*KNAT*) TF *BREVIPEDICELLUS* (BP) and the interacting *BEL1-LIKE HOMEODOMAIN* (BLH) protein *PENNYWISE* (PNY) have partially redundant roles in internode and radial patterning during inflorescence growth (see [65] for recent review). Their role is indicated by the enhanced double mutant phenotype with highly irregular internode length and increased branching [66]. BP and PNY act by restricting the expression of boundary genes such as *KNAT6*, *BLADE-ON-PETIOLE* (*BOP*) 1 and *BOP2*, and *ARABIDOPSIS THALIANA HOMEBOX GENE1* (*ATH1*) [67,68]. PNY acts redundantly with another *BELL1* family member, *POUNDFOOLISH* (PNF) to promote flowering and to establish the FM, since *pny pnf* double mutants

Table 1Examples of transcription factors with known roles in flower development in *Arabidopsis*.

| TF-family | Gene | Identifier | Function | Selected recent references/reviews |
|------------|--|---|---|------------------------------------|
| AP2/ERF | DRNL | AT1G24590 | Stamen development, organ founder cells | [179,183] |
| | AP2 | AT4G36920 | FM establishment and floral patterning | [44,46,123,182,184] |
| | ANT, AIL5, AIL6 | AT4G37750, AT5G57390, AT5G10510 | Organ morphology and growth, FM initiation, ovule development, flower primordium initiation | [185] |
| B3 ARF | SHN1, SHN2 | AT1G15360, AT5G25390 | Organ morphology | [217] |
| | TOE3 | AT5G67180 | Determinacy | [158] |
| | ARF5/MP | AT1G19850 | Flower primordium formation, ovule development | [77,78,187] |
| | ARF6, ARF8 | AT1G30330, AT5G37020 | Organ maturation, carpel development | [135–137] |
| | ARF3/ETT, ARF4 | AT2G33860, AT5G60450 | Gynoecium development, determinacy, organ polarity | [170,171,188] |
| B3 RAV | ARF2 | AT5G62000 | Organ size | [138,180,189] |
| | NGA1, NGA2, NGA3, NGA4 | AT2G46870, AT3G61970, AT1G01030, AT4G01500 | Style development | [19,188] |
| bHLH | MYC2, MYC3, MYC4, MYC5 | AT1G32640, AT5G46760, AT4G17880, AT5G46830 | Stamen development | [176] |
| | BPE | AT1G59640 | Petal growth | [133–135] |
| | bHLH010, bHLH089, bHLH091 | AT2G31220, AT1G06170, AT2G31210 | Anther development | [190] |
| | DYT1, AMS | AT4G21330, AT2G16910 | Anther development | [190,191] |
| | SPT, ALC | AT4G36930, AT5G67110 | Gynoecium development | [192] |
| BPC | HEC1, HEC2, HEC3 | AT5G67060, AT3G50330, AT5G09750 | Gynoecium development | [192,193] |
| | BPC1, BPC2, BPC3, BPC4, BPC6 | AT2G01930, AT1G14685, AT1G68120, AT2G21240, AT5G42520 | Organ number, morphology, meristem size | [175] |
| bZIP | TGA9, TGA10 | AT1G08320, AT5G06839 | Stamen development | [194,195] |
| bZIP | PAN | AT1G68640 | Organ number, determinacy | [153,154,194] |
| C2C2-GATA | GATA15, HAN, HANL2, GNC, GNL | AT3G06740, AT3G50870, AT4G36620, AT5G56860, AT4G26150 | Stamen growth, organ number, boundary formation | [138,196–198] |
| C2C2-YABBY | INO | AT1G23420 | Ovule development | [199] |
| | CRC | AT1G69180 | Gynoecium development, determinacy | [163,172,199] |
| | FIL | AT2G45190 | Organ number, size, morphology, boundary formation, FM identity | [186] |
| C2H2 | JAG, NUB | AT1G68480, AT1G13400 | Organ growth and patterning | [127] |
| | SUP | AT3G23130 | Boundary formation, floral patterning, determinacy, ovule development, petal and stamen development | [173,174] |
| | RBE | AT5G06070 | Petal development, boundary formation | [200] |
| GARP | KNU | AT5G14010 | Gynoecium development, determinacy | [30,34,152,178] |
| | UIF1 | AT4G37180 | Organ number, organ boundary | [167] |
| | KAN1, KAN2, KAN4/AT5 | AT5G16560, AT1G32240, AT5G42630 | Gynoecium development, organ polarity | [170,201] |
| HB | STM | AT1G62360 | Shoot meristem activity, carpel development | [97,202] |
| | KNAT2 | AT1G70510 | Carpel development | [202] |
| | PFS2/WOX6 | AT2G01500 | Ovule development, petal development | [203] |
| | WUS | AT2G17950 | Basic shoot meristem activity | [145,146,149,150,203] |
| | PHB, PHV, REV, CNA | AT2G34710, AT1G30490, AT5G60690, AT1G52150 | Ovule development, meristem activity and determinacy, organ number | [204] |
| | WOX1, PRS/WOX3 | AT3G18010, AT2G28610 | Sepal and petal development | [203] |
| | HDG1, HDG2, HDG5, HDG12, PDF2 | AT3G61150, AT1G05230, AT2G46680, AT1G17920, AT4G04890 | Stamen development, petal and stamen development | [205] |
| | JAB | AT4G17460 | Determinacy, gynoecium development | [206] |
| | BLR/PNY, PNF | AT5G02030, AT2G27990 | Organ number and specification of petals, stamen and carpels, inflorescence structure | [69,202] |
| | BEL1 | AT5G41410 | Ovule development | [120] |
| LFY | LFY | AT5G61850 | FM identity, floral patterning | [79,86,126] |
| MADS | CAL | AT1G26310 | Floral meristem initiation | [83] |
| | AP1 | AT1G69120 | A-class gene, organ identity, meristem identity | [87,96,182,207] |
| | AP3, PI | AT3G54340, AT5G20240 | B-class gene, petal and stamen identity | [124,182,207] |
| | SHP1, SHP2, STK | AT3G58780, AT2G42830, AT4G09960 | Carpel development, ovule development | [120,207] |
| | AG | AT4G18960 | Floral homeotic C-class gene, determinacy | [125,182,207] |
| | AGL24, SVP | AT4G24540, AT2G22540 | Floral transition, floral meristem identity | [80,102,207] |
| | SEP1, SEP2, SEP3, SEP4 | AT5G15800, AT3G02310, AT1G24260, AT2G03710 | Floral homeotic E-class, organ identity, meristem identity | [20,106,207] |
| | FUL | AT5G60910 | Floral transition, carpel development | [57,207] |
| | MYB65, MYB33 | AT3G11440, AT5G06100 | Flower maturation, anther development | [208] |
| | MYB21, MYB24, MYB57 | AT3G27810, AT5G40350, AT3G01530 | Flower maturation; stamen development | [176,209] |
| NAC | PWR | AT3G52250 | Determinacy, carpel development, petal development | [48] |
| | LMI2 | AT3G61250 | Meristem identity | [85] |
| | CUC1, CUC2, CUC3 | AT3G15170, AT5G53950, AT1G76420 | Boundary formation, gynoecium development, ovule development | [64,179,199] |
| SBP | SPL3, SPL8 | AT2G33810, AT1G02065 | FM identity, gynoecium development, anther development | [210] |
| SPL/NZZ | SPL/NZZ | AT4G27330 | Anther development, ovule development; stamen identity | [186] |
| SRS | STY1, STY2 | AT3G51060, AT4G36260 | Stamen development, gynoecium development | [192] |
| TCP | TCP3, TCP4, TCP5, TCP10, TCP13, TCP17, TCP15 | AT1G53230, AT3G15030, AT5G60970, AT2G31070, AT3G02150, AT5G08070, AT1G69690 | Petal morphology, petal growth, carpel development, flower maturation | [211,212] |
| Trihelix | PTL | AT5G03680 | Petal development, boundary formation | [200] |

fail to accumulate transcripts of the FM identity genes *LFY*, *AP1* and *CAULIFLOWER* (*CAL*) [69]. Recently it was shown that *BOP1/2* are directly repressed by PNY [70]. It has been proposed that *BOP1/2* promote directly the expression of *ATH1*, and by that indirectly *FLC*, that act as repressors of *FD* [70–72]. This may provide a mechanistic explanation for the additional function of PNY in the induction of flowering.

4. Gene-regulatory networks controlling flower initiation

In the flanks of the Arabidopsis IM, FMs are established as lateral meristems in an auxin-dependent manner. The requirement for auxin signaling is shown by the ‘naked pin’-like phenotype of *pin-formed 1* (*pin1*) and *monopteros* (*mp*) mutants, which essentially fail to initiate FMs [73,74]. *PIN1* encodes a major auxin efflux carrier involved in polar auxin transport [75], while *MP/ARF5* encodes an auxin-dependent AUXIN RESPONSE FACTOR (ARF) TF [76]. During the earliest stages of flower development, differentiation is temporarily suppressed in the FM in order to generate a sufficient amount of cells for formation of the inner floral whorls. *MP* directly activates the FM identity gene *LFY* as well as the AP2-type TF genes *AINTEGUMENTA* (*ANT*) and *AINTEGUMENTA-LIKE 6* (*AIL6*), which in turn also up-regulate *LFY* and stimulate FM and organ proliferation [77] (Fig. 1). *MP* activates these genes by switching chromatin state via recruitment of SWITCH/SUCROSE NONFERMENTING (SWI/SNF) -type nucleosome remodelers in an auxin-dependent manner [78].

LFY is a key regulator in the network establishing FM identity and is activated by flowering time genes *AGL24*, *SVP* and *SOC1* [79,31,80,81]. It is expressed prior to FM outgrowth (floral stage 1; for flower developmental stages see Fig. 2) and first activates *LATE MERISTEM IDENTITY1* (*LMI1*) and, in combination with *LMI1*, up-regulates the expression of the MADS-box gene *CAL* [82] (Fig. 3). The MADS-domain TF *AP1* acts in a partly redundant manner with its close paralog *CAL* in controlling FM identity [83]. *CAL* and *LFY*, partly in combination with *LMI2*, activate *AP1* that in turn can increase *LFY* expression in a positive feedback loop in order to maintain FM identity [84–88]. Comparison of the ChIP-seq data generated for the FM identity regulators *LFY* and *AP1* [86,87,89]

shows that approximately one third of *AP1* or *LFY* bound genes are also targeted by the other TF, respectively [90]. Genes involved in gibberellic acid (GA) homeostasis are for instance targeted by *LFY* as well as *AP1*.

GA is a phytohormone that positively regulates floral transition but inhibits FM establishment. Molecular mechanisms underlying these antagonistic roles have only recently started to be elucidated. According to the current model, GA initially up-regulates the expression of *SQUAMOSA PROMOTER BINDING PROTEIN LIKE* (*SPL*) TF genes as well as *LFY* in a concentration-dependent manner by degrading DELLA proteins, transcriptional co-regulators, via the 26S proteasome [91,92]. It is suggested that *LFY* then promotes a reduction of GA levels in part by promoting expression of *EUI-LIKE P450 A1* (*ELA1*) [93], a cytochrome p450 enzyme involved in GA catabolism [94]. Decreasing the GA concentration leads to the accumulation of GA-sensitive DELLA proteins, which in turn are required for up-regulation of *AP1*. This is mediated by binding of DELLA proteins recruited by *SPL9* to the *AP1* promoter [93]. Activation of *AP1* increases *ELA1* expression in a positive feedback loop, thus enforcing a more rapid switch from vegetative to reproductive phase [93].

In order to stabilize the switch to FM identity, *AP1* downregulates repressors of its own expression, such as the AP2 TF genes *TARGET OF EAT* (*TOE*) 1, *TOE3*, *SCHNARCHZAPFEN* and the *AP1* antagonist *TERMINAL FLOWER 1* [87]. *AP1* also establishes FM identity by repressing genes involved in the shoot developmental program and genes that control onset of flowering, like *FD*, *FD PARALOG* and *SPL9* [87]. In addition to that, *AP1* prevents the establishment of meristems in the axils of sepals in order to suppress axillary secondary flower formation [95]. Recently, it has been shown that *AP1* mediates this function by suppressing cytokinin accumulation via direct repression of the cytokinin biosynthetic gene *LONELY GUY1* and activation of the cytokinin degradation gene *CYTOKININ OXIDASE/DEHYDROGENASE3* [96]. Direct regulation is in agreement with binding of *AP1* to these loci [87,96]. By this, *AP1* counteracts the function of *SHOOTMERISTEMLESS* (*STM*), a homeodomain TF essential for shoot apical meristem formation and maintenance that upregulates cytokinin biosynthesis [97]. Changes in cytokinin levels affect the activity and downstream functions of the stem cell regulator *WUS* (see, e.g. [98]).

During early flower development, *AP1* is considered to be an FM identity gene. Later, around stage 3 of flower development, when sepal primordia arise (Fig. 2, stages according to Smyth et al. [99]),




Fig. 1. Auxin mediated floral primordium initiation. At high levels of auxin, repressive Aux/IAA proteins are released from the Auxin Response Factor ARF5/MP. MP then activates expression of *LFY* as well as *ANT* and *AIL6*, which act in parallel with *MP* to promote *LFY* induction. Up-regulation of *LFY* and *ANT/AIL6* activate the gene regulatory networks that initiate the formation of floral primordia.




Fig. 2. Confocal microscopic analysis of early flower bud stages. Overview of an inflorescence meristem and early flower meristem stages 1–6. The early stages are defined as: flower buttress arises (stage 1), flower primordium forms (stage 2), sepal primordia arise (stage 3), sepals overlie floral meristem (stage 4), petal (*) and stamen primordia arise (stage 5), sepals enclose the buds (stage 6). Stages according to [99]. Confocal image shows ubiquitously expressed endomembrane-localized GFP reporter gene.

AP1 expression becomes restricted to the outer whorls, where it specifies sepal and petal identity [95]. A molecular mechanism for the functional change of AP1 could be different protein-protein interactions (Fig. 3): AP1 dimerizes with the MADS-domain TFs SHORT VEGETATIVE PHASE (SVP), AGL24 and SEP3 [100,101]. The expression of AGL24 and SVP is mostly restricted to flower developmental stages 1 and 2, before sepal primordia arise, and decreases afterwards [101,102]. Genetic data, ChIP experiments and protein-protein interaction studies suggest that SVP, AGL24 and AP1 act redundantly in maintaining the FM in a meristematic state by preventing precocious expression of floral homeotic genes and probably do so by forming complexes with each other and recruiting additional factors for SEP3 repression [31,80,102]. In *svp agl24* double mutants, SOC1, which is usually not expressed in FMs, can take over their function [31,102]. Gregis et al. showed that AP1/SVP or AP1/AGL24 complexes interact with SEUSS (SEU) and LEUNIG [101], which form a transcriptional co-repressor complex [103,104]. Furthermore, SEP3 repression by SVP was proposed to be mediated via recruiting the PcG factor TERMINAL FLOWER 2/LIKE HETEROCHROMATIN PROTEIN 1 and changes in H3K27me3 deposition, whereas AGL24 and SOC1 act via recruitment of proteins with histone deacetylase activity in the absence of SVP [31]. Repression of SEP3 in the first stages is crucial for meristem maintenance, since SEP proteins directly activate the B and C class genes in concert with LFY and AP1 [31,105,106]. De-repression of SEP3 in the early stage 3 allows the formation of AP1/SEP3 complexes leading to activation of homeotic gene expression and as a result of that floral organ identity specification, while flowering time genes are kept in a repressed state [31,102, 105–107]. Interestingly, approximately 80% of the putative AP1 target genes are down-regulated in early FM development, but at later stages AP1 together with SEP3 preferentially act as activators of gene expression [87,89]. SEP3 as well has been shown to interact with SEU [108], indicating that recruitment of the co-repressors occurs more frequently by different floral MADS-domain TF complexes.




Fig. 3. LFY-mediated pathway for the establishment of floral meristem identity. During the transition from IM to FM (stages 1–2), LFY activates AP1, possibly by interacting with LMI2. Indirect activation of AP1, by LFY is achieved through LMI1 and CAL. Up-regulation of AP1 results in a positive feedback loop triggering LFY up-regulation in order to stabilize the switch from IM to FM. AP1 and LFY upregulate SEP3 expression, which is required for strong activation of homeotic B- and C-class genes around stage 3 of flower development.

Genome wide identification of SVP target genes in inflorescences identified 265 putative targets that are common to SVP and AP1 [80]. Those genes are enriched for TF genes, including *KANAD11*, *PHABULOSA* and *ARF3* [80]. Furthermore, SVP binds to genes involved in auxin, GA, cytokinin and jasmonic acid (JA) signaling, and to genes with roles in posttranslational regulation in the FM [80]. SVP, SOC1 and AGL24 also promote *LFY* expression while LFY may be an indirect negative regulator of SVP and AGL24 [109].

At later stages of FM development, AGAMOUS, AP1 and SEP3 directly repress SVP, AGL24 and SOC1 in a regulatory feedback loop to allow floral organ patterning and to restrict meristem activity [80,87,101,109, 110]. At the same time, homeotic B and C class genes become activated allowing meristem differentiation and floral organ initiation.

5. Gene-regulatory networks controlling floral organ specification

After initiation of the FM, the distinct floral whorls are established resulting in the formation of the different types of floral organs. FM differentiation is initiated around floral stages 2–3, when the expression of floral homeotic B and C class genes commences in a spatially restricted whorled pattern within the meristem (Fig. 2). At stage 3, sepal primordia appear, followed by the organ primordia of the inner whorls in stage 5 and sepal outgrowth enclosing the buds at stage 6 [99]. With one exception, all classical floral homeotic genes of the ABCE model encode for members of the MADS domain TF family. Their combinatorial interactions in floral organ specification were proposed to be based on formation of organ-specific higher-order protein complexes, as suggested by the ‘floral quartet model’ [10,111]. According to this model, the exact composition of those complexes determines the identity of the organ to be formed. In Arabidopsis, sepal identity specification is mediated by A-class protein AP1 in combination with E-class proteins (SEP1–4), while AP1 together with the B-class proteins APETALA3 (AP3), PISTILLATA (PI) and SEP proteins specify petal identity. Stamens are defined by interactions of the C-class protein AG with AP3 and PI, which is mediated by SEP proteins, whereas carpels are specified by AG/SEP complexes [10,112]. SEP proteins were found to act as mediators of higher-order complex formation and, similar to AP1, possess strong transcriptional activation potential [112]. Only the combined overexpression of the different floral homeotic gene classes leads to the conversion of vegetative leaves to specific types of floral organs [112,113], suggesting that the heteromeric floral homeotic protein complexes are not only necessary, but also sufficient to specify floral organ identity.

In planta immunoprecipitation coupled to mass spectrometry of endogenously expressed MADS-domain TFs further supported the interactions that were proposed in the floral quartet model [114]. The floral homeotic protein complexes can also be assembled *in vitro* [114–116]. Higher-order complex formation can induce short-range looping of the DNA which can be relevant for recruitment to target gene promoters and for target gene regulation [116,117]. Although protein-protein interactions within the MADS-domain TF family are well studied [100, 118], the exact molecular mechanisms of action and determinants of functional specificity are still not well understood. The stabilization of DNA-binding by two TF binding sites at (complex-)specific distances may be one aspect [119]. It is also possible that higher-order complexes show specific characteristics in the recruitment of co-factors or interactions with Non-MADS-TFs. Indeed, floral homeotic MADS-domain TFs have been found to interact with members of other TF families. For example, the Three Amino acid Loop Extension Homeodomain (TALE-HD) TF BELL1 (BEL1) was found to interact with MADS-domain protein complexes consisting of SEP3 and of AG-like proteins [120]. The interaction with BEL1 is important for ovule specification, and the interaction is thought to modify the activity of the SEP-AG complex that otherwise specifies carpel identity. Interestingly, TFs of different families have been identified in AP1-complexes as well: TALE-HD TF BLH1, PNY and KNAT3 as well as ARF2 and SPL8 have been enriched in protein complex

isolation experiments [114]. Incorporating additional factors into ternary complexes may have impact on target gene specificity and adds options for fine-tuning gene expression responses. The protein complex isolation experiments have identified additional complex partners of floral homeotic proteins, which provide clues on how these factors may regulate target gene activity. Most remarkably, the chromatin-remodeling factors BRAHMA (BRM), SPLOYED (SYD) and PICKLE were identified as components of MADS-domain TF complexes indicating an important role during flower development [114,121]. BRM and SYD are regarded as trxG factors and have been shown to reverse Polycomb repression [121]. Loss of BRM activity results in the change of H3K27me3 marks at hundreds of genes [122]. SYD and BRM are important for AP3 and AG activation during flower patterning and are directly recruited to their promoters by LFY and SEP3 [121]. A further link to epigenetic gene regulation of MADS-domain proteins has been revealed by studying the impact of AP1 and SEP3 DNA-binding on chromatin accessibility. Pajoro et al. [89] found that the binding of these TFs to genomic regions precedes changes in chromatin accessibility, in particular ‘opening’ of the chromatin. Therefore, these factors provide the first candidates of “pioneer factors” in plants. Changes in chromatin accessibility induced by AP1 and SEP3 then would allow binding of other TFs to genomic regions thereby mediating spatio-temporal control of gene expression in flower development.

The question how major regulatory TFs of flower initiation and differentiation exert their functions has started to become elucidated by the identification of their target gene networks. Target gene identification has been achieved for TFs involved in FM or floral organ specification such as AP1 [87,89], AP2 [123], AP3 and PI [124], AG [125], LFY [86,126], SEP3 [89,106], as well as for the downstream acting growth regulator JAGGED [127]. For example, target gene analyses of the B-class proteins AP3 and PI, which form obligate heterodimers [128,129], have helped elucidate control of petal and stamen development. AP3 transcripts accumulate in the petal and stamen whorls from stage 3 onwards until expression levels decrease at stage 9, when stamens have formed and petal primordia stalk at the base [130]. PI also shows expression from stage 3 onwards [131]. Due to their expression during different stages of flower development it is assumed that AP3 and PI fulfill stage-specific functions by targeting distinct sets of genes: This is based on perturbation experiments where most dramatic effects on organ identity can be seen when AP3 downregulation occurs in the very early stages, while later disturbances mainly affect stamen maturation [124]. Genome-wide identification of target genes at stage 5, when petal and stamen primordia arise, confirms autoregulatory binding to the AP3 locus [124]. The F-box gene UNUSUAL FLORAL ORGANS (UFO), which encodes a LFY cofactor and thereby up-regulates AP3 expression [132], is also a direct AP3 target. Furthermore, central players of petal and stamen differentiation like RABBIT EARS, PETAL LOSS, SPOROCTELESS/NOZZLE are directly regulated by AP3/PI [124]. Moreover, genes involved in carpal development, like HECATE (HEC) 1 and HEC2, CRABSCLAW (CRC) or SHATTERPROOF 2 are directly repressed in order to prevent ectopic formation of carpels. B class proteins are also involved in hormone signaling: Especially factors involved in auxin (INDOLE-3-ACETIC ACID INDUCIBLE (IAA) 2, IAA18, IAA27, ARF3, ARF8), JA (JASMONATE-ZIM-DOMAIN PROTEIN (JAZ) 9, JAZ10) and GA (GIBBERELLIN 2-OXIDASE (GA2OX) 1, GA2OX3, GA2OX4) signaling are among their downstream targets [124]. Those hormones play a crucial role in determining the size and morphology of floral organs. ARF8 for example limits cell expansion in later stages of petal development in combination with BIG PETAL via combined down-regulation of auxin responsive genes [133–135]. In *arf6 arf8* double mutants, organ elongation in perianth organs and stamen is delayed, probably due to reduced JA levels [136,137]. Feeding experiments suggest that exogenous JA application can promote petal elongation and anther dehiscence, but does not complement other phenotypes, like vascular patterning and changes in petal cell morphology related to *arf6 arf8* mutations [136,137]. Growth defects in stamens often result in stigma protrusion phenotypes, which are related to decreased fertility

due to an imbalanced size ratio of carpel and stamens. This phenotype can be observed in other mutants as well, for example in *arf2* mutants. ARF2 has been shown to be a repressor of GATA TF genes *GATA*, *NITRATE INDUCIBLE*, *CARBON METABOLISM INVOLVED (GNC)* and *GNC-LIKE (GNL)*, and *gnc gnl* loss of function in *arf2* mutant background suppresses the *arf2* stamen growth defect [138]. GNC and GNL have been shown to be regulated by auxin and GA and act downstream of AP3/PI, thus showing that gene-regulatory networks of organogenesis, growth and hormone signaling are tightly interwoven. GNC and GNL are important regulators of different physiological processes, including nitrogen metabolism, sugar sensing and photosynthesis [138–141].

Comparison of genes directly bound by different classes of homeotic TFs show a high overlap [125,142], supporting the idea that they act as part of protein complexes on common sets of target genes. A fraction of the genes appears to be regulated by the different organ-specific complexes in an antagonistic manner, suggesting that the different homeotic protein complexes show partly overlapping DNA-binding activities, but differences in regulatory effects [125,142]. Time-series ChIP-seq experiments of AP1 and SEP3 showed that these proteins are recruited to their target gene promoters in a partly stage-specific manner, supporting the idea that homeotic proteins regulate their genes in a dynamic fashion [89]. The stage-specific differences in promoter occupancy and target gene regulation may depend on availability of cofactors or epigenetic state. However, the detailed molecular mechanisms underlying the stage- and organ-specificity of target gene selection by homeotic proteins remain to be elucidated.

6. Termination of stem cell activity in floral meristems

Stem cell activity in shoot and root meristems is controlled by the complex interplay of TFs, hormonal and cellular signaling pathways (see [143,144] for recent reviews). The homeobox TF WUS is required for the maintenance of the stem cell populations in shoot meristems [145,146], while meristem size is determined by feedback regulation with the CLAVATA signaling pathway, involving movement of WUS within meristems from the organizing center, where it is expressed, to the neighboring stem cells [147]. The stem cell niche in FM is only transiently maintained and a prolonged FM activity in the center of the flower can be observed in several mutants with a compromised determinacy (reviewed by [148]).

Floral determinacy is achieved by exhausting the pool of stem cells in a FM and is tightly linked with shutting down WUS mediated stem cell maintenance in a WUS-AG-feedback loop. During stage 3 of flower development, WUS, together with LFY, promotes AG expression in the center of the flower [149,150], which might be at least partly facilitated by non-cell autonomous activity of WUS [151]. Once AG is activated, it directly and indirectly represses WUS expression. On the one hand, AG directly binds to the promoter of WUS from stage 3 on modifying histone H3 with H3K27me3 marks in a CURLY LEAF dependent manner [30]. On the other hand, AG induces the expression of C2H2-zinc finger TF gene KNUCKLES (KNU) that in turn directly represses WUS from stage 6 onwards, where no WUS mRNA is detectable anymore [30,34,152]. Interestingly KNU itself shows H3K27me3 marks and the activation of its expression was shown to be cell division-dependent. There is evidence showing that AG binding to the KNU promoter interferes with PcG binding, causing a dilution of H3K27me3 marks in a cell division-dependent manner by incorporating unmodified histone H3 [34]. Therefore, AG has a dual and separable function in carpal specification and floral meristem determinacy [152].

The dosage-dependent role of AG in regulating floral determinacy becomes clear from mutants with decreased AG protein levels. For example, the basic leucine zipper TF PERIANTHIA (PAN) is a direct regulator of AG in a whorl specific manner, with a putative repressive function in whorl 1 and a promoting function in whorl 4. Mutations in *pan lfy* and *pan seu* show an increased indeterminacy phenotype [153,154]. In addition to transcriptional regulation of AG, posttranslational processing of

AG pre-mRNA impacts floral determinacy as well. Consistent with this, posttranscriptional regulators of AG show loss of determinacy and compromised *WUS* shutdown (see, e.g. [155–157]). Furthermore, miRNA processing is related to floral stem cell termination: the AG repressors *AP2* and *TOE3* are targeted by miR172, and overexpression of miRNA resistant proteins or mutations in the miR172 sequence result in indeterminate flowers [48,158]. Mutations affecting components of the miRNA processing machinery as *DICER-LIKE 1*, *ARGONAUTE (AGO) 1* or *AGO10* show a determinacy phenotype, partly because of their impact on miR172 activity [159–161]. Another factor in the AGO1 pathway is *SQUINT (SQN)*, which is a cyclophilin 40 homolog and has been identified to enhance weak indeterminacy phenotypes of *crabs claw (crc)*, see below) or *ag* [162–164]. *SQN* promotes the activity of AGO1 in plants via physical interaction with Heat shock protein 90 [165]. In addition to its role in floral meristem termination, Prunet et al. [166] propose an additional role for *SQN* in stem cell homeostasis via the *CLAVATA* signaling pathway.

The *trxG* protein *ULT1* is a negative regulator of stem cell activity, indicated by supernumerary floral organs in *ult1* mutants, and it restricts *WUS* expression to the organizing center [37,38,163]. At the same time *ULT1* interacts with *ARABIDOPSIS HOMOLOG OF TRITHORAX1* in order to reduce repressive H3K27me3 marks at the AG locus [39]. Recently, the Myb-domain TF *ULTRAPETALA1 INTERACTING FACTOR1 (UIF)* has been identified to interact with *ULT1* [167]. *UIF1* is able to bind *WUS* and AG genomic sequences *in vitro*, thereby creating the potential to recruit *ULT1* to these loci.

In an enhancer screen using the weak *ag-10* mutant allele, novel regulators of floral meristem determinacy have been identified recently. One of these identified genes encodes *TOPOISOMERASE1α*, which appears to have a general role decreasing nucleosome density in the genome, thereby mediating PcG recruitment to the *WUS* locus via an unknown mechanism [168]. In the same screen, *ARF3*, which has well-known roles in polarity establishment within leaves and carpels [169,170], has been identified as well [171]. *ARF3* confers its role via repression of *WUS* activity, and this partly depends on AG. *ARF3* is a target of *AP2*, and contributes to mediating its function in floral determinacy [171].

Moreover, target genes of AG may contribute to AG's function in floral determinacy, as has been shown in the case of the *YABBY* TF *CRC*. The *crc* single mutant has no defect in floral determinacy, but when combined with heterozygous *ag* or other mutants that act in floral determinacy, supernumerary whorls of floral organs can be observed [163,172]. *CRC* is a direct target of AG [125], but the exact molecular role of *CRC* in determinacy has not been unraveled yet. Another downstream target of AG is *SUPERMAN (SUP)*, which encodes a C2H2-zinc finger TF that plays a role in boundary maintenance between 3rd and fourth whorl, indicated by supernumerary stamen on the expense of carpels in *sup* [173,174]. However, the exact molecular role of *SUP*, as in the case of *CRC*, in FM determinacy remains to be elucidated.

7. Understanding the networks underlying inflorescence and flower development: current status and outlook

Many important regulatory genes with roles in inflorescence and flower development have been identified over the last two decades. Basic genetic and molecular interactions have been characterized for some of these factors, largely boosted by the technical progress in genome-wide experimental analyses. It has now become clear that gene regulation in plant development is achieved at multiple levels: transcriptional regulation by TFs, epigenetic factors and (long) non-coding RNAs, posttranscriptional regulation by miRNAs, and regulation at the level of protein activity or stability. In particular the regulation at posttranslational level, e.g. the roles of posttranslational modifications of TFs or the assembly of regulatory complexes, is still poorly understood. On the other hand, it has become clear that there is a tight interplay between TFs and epigenetic regulation in the control of

developmental transitions in plants, such as the floral transition and switches in meristem identity. A nice example is the conditional auxin-mediated recruitment of SWI/SNF2-type remodelers by MP, which triggers FM initiation [77]. It is also becoming clear that regulation of most developmentally controlled genes requires a combinatorial input of multiple TFs and other transcriptional regulators (see, e.g. [85,114,120,175,176]). The knowledge on genes and genetic interactions underlying flower development has enabled mathematical modeling approaches, which are able to generate novel, experimentally testable hypotheses on underlying mechanisms [177]. Live imaging using confocal microscopy generates quantitative data that can be used for computational morphodynamics approaches [213]. However, quantification of levels of regulatory molecules and their turnover at cellular resolution *in planta* remains an important challenge. This type of information will help us to understand the complex regulatory interactions that can be observed in gene-regulatory and hormonal signaling networks governing development.

While we do have a good knowledge of important regulatory processes in inflorescence and flower development, we still need to learn more about the impact of environmental fluctuations on specific gene-regulatory interactions. Genetic and molecular analyses are often performed under standardized environmental conditions, while in nature, plants are exposed to significant short and long term fluctuations, e.g. in temperature, light intensity/quality and water supply. It is clear that floral transition is highly adaptive in response to environmental factors, but how environmental robustness of inflorescence and flower development is achieved is largely unclear. Next to *Arabidopsis*, genes controlling inflorescence and flower development are now being systematically identified and analyzed in other model and crop plant species. Among the well-studied crop species are tomato (*Solanum lycopersicum*) and rice (*Oryza sativa*) [181,214,215]. Systematic evolutionary comparison of gene-regulatory networks provides rich information on conserved and non-conserved regulators and regulatory pathways, which has important impact on transferring knowledge from model to crop plant species. For instance, homologs of *UFO* and *LFY* have been implicated in inflorescence patterning in *Solanum*, along with other regulatory factors [181]. Comparative studies are therefore highly valuable in understanding the diversity and evolution of gene-regulatory networks.

In order to discern mechanisms of regulatory evolution, the comparison of closely related species and ecotypes may be particularly interesting (see, e.g. [216]). Here, the combination of genetic analyses (facilitated by novel genome editing tools such as RNA guided Cas9) with genome-wide functional genomics techniques (e.g. ChIP-seq) is likely to provide a new level of understanding of the evolution of inflorescence and flower development in angiosperms in the near future.

Transparency document

The Transparency document associated with this article can be found, in online version.

Acknowledgements

The authors wish to thank Dijun Chen for help with data review in preparation for this manuscript. K. Kaufmann wishes to thank the Alexander-von-Humboldt Foundation and the BMBF for funding. We wish to apologize to all authors of relevant publications whose work could not be cited due to space limitations.

References

- [1] F. Andres, G. Coupland, The genetic basis of flowering responses to seasonal cues, *Nat. Rev. Genet.* 13 (2012) 627–639.
- [2] M. Romera-Branchat, F. Andres, G. Coupland, Flowering responses to seasonal cues: what's new? *Curr. Opin. Plant Biol.* 21 (2014) 120–127.

- [3] S. Duncan, S. Holm, J. Questa, J. Irwin, A. Grant, C. Dean, Seasonal shift in timing of vernalization as an adaptation to extreme winter, *Elife* 4 (2015) e06620.
- [4] J. Hepworth, C. Dean, Flowering locus C's lessons: conserved chromatin switches underpinning developmental timing and adaptation, *Plant Physiol.* 168 (2015) 1237–1245.
- [5] H. Sommer, J.P. Beltran, P. Huijser, H. Pape, W.E. Lonig, H. Saedler, Z. Schwarzommer, Deficiens, a homeotic gene involved in the control of flower morphogenesis in *antirrhinum-majus* - the protein shows homology to transcription factors, *EMBO J.* 9 (1990) 605–613.
- [6] M.F. Yanofsky, H. Ma, J.L. Bowman, G.N. Drews, K.A. Feldmann, E.M. Meyerowitz, The protein encoded by the *Arabidopsis* homeotic gene *agamous* resembles transcription factors, *Nature* 346 (1990) 35–39.
- [7] E.S. Coen, E.M. Meyerowitz, The war of the whorls - genetic interactions controlling flower development, *Nature* 353 (1991) 31–37.
- [8] G.W. Haughn, C.R. Somerville, Genetic-control of morphogenesis in *Arabidopsis*, *Dev. Genet.* 9 (1988) 73–89.
- [9] S. Pelaz, G.S. Ditta, E. Baumann, E. Wisman, M.F. Yanofsky, B and C floral organ identity functions require *SEPALLATA* MADS-box genes, *Nature* 405 (2000) 200–203.
- [10] G. Theissen, Development of floral organ identity: stories from the MADS house, *Curr. Opin. Plant Biol.* 4 (2001) 75–85.
- [11] B. Causier, Z. Schwarz-Sommer, B. Davies, Floral organ identity: 20 years of ABCs, *Semin. Cell Dev. Biol.* 21 (2010) 73–79.
- [12] A.S. Chandrabali, B.A. Berger, D.G. Howarth, P.S. Soltis, D.E. Soltis, Evolving ideas on the origin and evolution of flowers: new perspectives in the genomic era, *Genetics* 202 (2016) 1255–1265.
- [13] G. Capovilla, M. Schmid, D. Posé, Control of flowering by ambient temperature, *J. Exp. Bot.* 66 (2015) 59–69.
- [14] D. Posé, L. Yant, M. Schmid, The end of innocence: flowering networks explode in complexity, *Curr. Opin. Plant Biol.* 15 (2012) 45–50.
- [15] J. Jin, H. Zhang, L. Kong, G. Gao, J. Luo, PlantTFDB 3.0: a portal for the functional and evolutionary study of plant transcription factors, *Nucleic Acids Res.* 42 (2014) D1182–D1187.
- [16] P. Pérez-Rodríguez, D.M. Riaño-Pachón, L.G. Corrêa, S.A. Rensing, B. Kersten, B. Mueller-Roeber, PlantTFDB: updated content and new features of the plant transcription factor database, *Nucleic Acids Res.* 38 (2010) D822–D827.
- [17] D. Wilson, V. Charoensawan, S.K. Kummerfeld, S.A. Teichmann, DBD—taxonomically broad transcription factor predictions: new content and functionality, *Nucleic Acids Res.* 36 (2008) D88–D92.
- [18] J.P. Alvarez, A. Goldschmidt, I. Efroni, J.L. Bowman, Y. Eshed, The *NGATHA* distal organ development genes are essential for style specification in *Arabidopsis*, *Plant Cell* 21 (2009) 1373–1393.
- [19] M. Trigueros, M. Navarrete-Gómez, S. Sato, S.K. Christensen, S. Pelaz, D. Weigel, M.F. Yanofsky, C. Ferrández, The *NGATHA* genes direct style development in the *Arabidopsis* gynoecium, *Plant Cell* 21 (2009) 1394–1409.
- [20] G. Ditta, A. Pinyopich, P. Robles, S. Pelaz, M.F. Yanofsky, The *SEP4* gene of *Arabidopsis thaliana* functions in floral organ and meristem identity, *Curr. Biol.* 14 (2004) 1935–1940.
- [21] K. Kaufmann, A. Pajaro, G.C. Angenent, Regulation of transcription in plants: mechanisms controlling developmental switches, *Nat. Rev. Genet.* 11 (2010) 830–842.
- [22] D.S. O'Maoláidigh, B. Thomson, A. Raganelli, S.E. Wuest, P.T. Ryan, K. Kwaśniewska, C.C. Carles, E. Graciet, F. Wellmer, Gene network analysis of *Arabidopsis thaliana* flower development through dynamic gene perturbations, *Plant J.* 83 (2015) 344–358.
- [23] G. He, A.A. Elling, X.W. Deng, The epigenome and plant development, *Annu. Rev. Plant Biol.* 62 (2011) 411–435.
- [24] M. de la Paz Sanchez, P. Aceves-García, E. Petrone, S. Steckenborn, R. Vega-León, E.R. Álvarez-Buylla, A. Garay-Arroyo, B. García-Ponce, The impact of polycomb group (PcG) and trithorax group (TrxG) epigenetic factors in plant plasticity, *New Phytol.* 208 (2015) 684–694.
- [25] A. Förderer, Y. Zhou, F. Turck, The age of multiplexity: recruitment and interactions of polycomb complexes in plants, *Curr. Opin. Plant Biol.* 29 (2016) 169–178.
- [26] I. Mozgova, L. Hennig, The polycomb group protein regulatory network, *Annu. Rev. Plant Biol.* 66 (2015) 269–296.
- [27] F. Turck, F. Roudier, S. Farrona, M.L. Martin-Magniette, E. Guillaume, N. Buisine, S. Gagnot, R.A. Martienssen, G. Coupland, V. Colot, *Arabidopsis* TFL2/LHP1 specifically associates with genes marked by trimethylation of histone H3 lysine 27, *PLoS Genet.* 3 (2007), e86.
- [28] X. Zhang, S. Germann, B.J. Blus, S. Khorasanizadeh, V. Gaudin, S.E. Jacobsen, The *Arabidopsis* LHP1 protein colocalizes with histone H3 Lys27 trimethylation, *Nat. Struct. Mol. Biol.* 14 (2007) 869–871.
- [29] M. Lafos, P. Kroll, M.L. Hohenstatt, F.L. Thorpe, O. Clarenz, D. Schubert, Dynamic regulation of H3K27 trimethylation during *Arabidopsis* differentiation, *PLoS Genet.* 7 (2011), e1002040.
- [30] X. Liu, Y.J. Kim, R. Müller, R.E. Yumul, C. Liu, Y. Pan, X. Cao, J. Goodrich, X. Chen, AGAMOUS terminates floral stem cell maintenance in *Arabidopsis* by directly repressing *WUSCHEL* through recruitment of polycomb group proteins, *Plant Cell* 23 (2011) 3654–3670.
- [31] C. Liu, W. Xi, L. Shen, C. Tan, H. Yu, Regulation of floral patterning by flowering time genes, *Dev. Cell* 16 (2009) 711–722.
- [32] W. Deng, D.M. Buzas, H. Ying, M. Robertson, J. Taylor, W.J. Peacock, E.S. Dennis, C. Helliwell, *Arabidopsis* polycomb repressive complex 2 binding sites contain putative GAGA factor binding motifs within coding regions of genes, *BMC Genomics* 14 (2013) 593.
- [33] M. Lodha, C.F. Marco, M.C. Timmermans, The ASYMMETRIC LEAVES complex maintains repression of *KNOX* homeobox genes via direct recruitment of polycomb-repressive complex2, *Genes Dev.* 27 (2013) 596–601.
- [34] B. Sun, L.S. Looi, S. Guo, Z. He, E.S. Gan, J. Huang, Y. Xu, W.Y. Wee, T. Ito, Timing mechanism dependent on cell division is invoked by polycomb eviction in plant stem cells, *Science* 343 (2014) 1248559.
- [35] J. Goodrich, P. Puangsomlee, M. Martin, D. Long, E.M. Meyerowitz, G. Coupland, A polycomb-group gene regulates homeotic gene expression in *Arabidopsis*, *Nature* 386 (1997) 44–51.
- [36] M. Calonje, R. Sanchez, L. Chen, Z.R. Sung, EMBRYONIC FLOWER1 participates in polycomb group-mediated AG gene silencing in *Arabidopsis*, *Plant Cell* 20 (2008) 277–291.
- [37] J.C. Fletcher, The ULTRAPETALA gene controls shoot and floral meristem size in *Arabidopsis*, *Development* 128 (2001) 1323–1333.
- [38] C.C. Carles, D. Hoffnes-Inada, K. Reville, K. Lertpiriyapong, J.C. Fletcher, ULTRAPETALA1 encodes a SAND domain putative transcriptional regulator that controls shoot and floral meristem activity in *Arabidopsis*, *Development* 132 (2005) 897–911.
- [39] C.C. Carles, J.C. Fletcher, The SAND domain protein ULTRAPETALA1 acts as a trithorax group factor to regulate cell fate in plants, *Genes Dev.* 23 (2009) 2723–2728.
- [40] M.M. Monfared, C.C. Carles, P. Rossignol, H.R. Pires, J.C. Fletcher, The ULT1 and ULT2 trxG genes play overlapping roles in *Arabidopsis* development and gene regulation, *Mol. Plant* 6 (2013) 1564–1579.
- [41] L. Pu, M.S. Liu, S.Y. Kim, L.F. Chen, J.C. Fletcher, Z.R. Sung, EMBRYONIC FLOWER1 and ULTRAPETALA1 act antagonistically on *Arabidopsis* development and stress response, *Plant Physiol.* 162 (2013) 812–830.
- [42] F. Borges, R.A. Martienssen, The expanding world of small RNAs in plants, *Nat. Rev. Mol. Cell Biol.* 16 (2015) 727–741.
- [43] H.O. Iwakawa, Y. Tomari, The functions of microRNAs: mRNA decay and translational repression, *Trends Cell Biol.* 25 (2015) 651–665.
- [44] X. Chen, A microRNA as a translational repressor of APETALA2 in *Arabidopsis* flower development, *Science* 303 (2004) 2022–2025.
- [45] J. Mathieu, L.J. Yant, F. Mürdter, F. Küttner, M. Schmid, Repression of flowering by the miR172 target SMZ, *PLoS Biol.* 7 (2009), e1000148.
- [46] H. Wollmann, E. Mica, M. Todesco, J.A. Long, D. Weigel, On reconciling the interactions between APETALA2, miR172 and AGAMOUS with the ABC model of flower development, *Development* 137 (2010) 3633–3642.
- [47] M.J. Aukerman, H. Sakai, Regulation of flowering time and floral organ identity by a microRNA and its APETALA2-like target genes, *Plant Cell* 15 (2003) 2730–2741.
- [48] R.E. Yumul, Y.J. Kim, X. Liu, R. Wang, J. Ding, L. Xiao, X. Chen, POWERDRESS and diversified expression of the MIR172 gene family bolster the floral stem cell network, *PLoS Genet.* 9 (2013), e1003218.
- [49] J. Jin, J. Liu, H. Wang, L. Wong, N.H. Chua, PLncDB: plant long non-coding RNA database, *Bioinformatics* 29 (2013) 1068–1071.
- [50] J.A. Chekanova, Long non-coding RNAs and their functions in plants, *Curr. Opin. Plant Biol.* 27 (2015) 207–216.
- [51] S. Swiezewski, F. Liu, A. Magusin, C. Dean, Cold-induced silencing by long antisense transcripts of an *Arabidopsis* polycomb target, *Nature* 462 (2009) 799–802.
- [52] T. Csorba, J.I. Questa, Q. Sun, C. Dean, Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) 16160–16165.
- [53] A. Samach, H. Onouchi, S.E. Gold, G.S. Ditta, Z. Schwarz-Sommer, M.F. Yanofsky, G. Coupland, Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*, *Science* 288 (2000) 1613–1616.
- [54] L. Corbesier, C. Vincent, S. Jang, F. Fornara, Q. Fan, I. Searle, A. Giakountis, S. Farrona, L. Gissot, C. Turnbull, G. Coupland, FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*, *Science* 316 (2007) 1030–1033.
- [55] Z. Tao, L. Shen, C. Liu, L. Liu, Y. Yan, H. Yu, Genome-wide identification of SOC1 and SVP targets during the floral transition in *Arabidopsis*, *Plant J.* 70 (2012) 549–561.
- [56] R.G. Immink, D. Posé, S. Ferrario, F. Ott, K. Kaufmann, F.L. Valentini, S. de Folter, F. van der Wal, A.D. van Dijk, M. Schmid, G.C. Angenent, Characterization of SOC1's central role in flowering by the identification of its upstream and downstream regulators, *Plant Physiol.* 160 (2012) 433–449.
- [57] S. Melzer, F. Lens, J. Gennen, S. Vanneste, A. Rohde, T. Beeckman, Flowering-time genes modulate meristem determinacy and growth form in *Arabidopsis thaliana*, *Nat. Genet.* 40 (2008) 1489–1492.
- [58] M. Abe, Y. Kobayashi, S. Yamamoto, Y. Daimon, A. Yamaguchi, Y. Ikeda, H. Ichinoki, M. Notaguchi, K. Goto, T. Araki, FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex, *Science* 309 (2005) 1052–1056.
- [59] P.A. Wigge, M.C. Kim, K.E. Jaeger, W. Busch, M. Schmid, J.U. Lohmann, D. Weigel, Integration of spatial and temporal information during floral induction in *Arabidopsis*, *Science* 309 (2005) 1056–1059.
- [60] M. Schmid, N.H. Uhlenhaut, F. Godard, M. Demar, R. Bressan, D. Weigel, J.U. Lohmann, Dissection of floral induction pathways using global expression analysis, *Development* 130 (2003) 6001–6012.
- [61] Y. Wang, J. Li, Molecular basis of plant architecture, *Annu. Rev. Plant Biol.* 59 (2008) 253–279.
- [62] Z.W. Teo, S. Song, Y.Q. Wang, J. Liu, H. Yu, New insights into the regulation of inflorescence architecture, *Trends Plant Sci.* 19 (2014) 158–165.
- [63] Y. Han, H. Yang, Y. Jiao, Regulation of inflorescence architecture by cytokinins, *Front. Plant Sci.* 5 (2014) 669.
- [64] A. Peaucelle, H. Morin, J. Traas, P. Laufs, Plants expressing a miR164-resistant CUC2 gene reveal the importance of post-meristematic maintenance of phyllotaxy in *Arabidopsis*, *Development* 134 (2007) 1045–1050.
- [65] S.R. Hepworth, V.A. Pautot, Beyond the divide: boundaries for patterning and stem cell regulation in plants, *Front. Plant Sci.* 6 (2015) 1052.
- [66] H.M. Smith, S. Hake, The interaction of two homeobox genes, *BREVIPEDICELLUS* and *PENNYWISE*, regulates internode patterning in the *Arabidopsis* inflorescence, *Plant Cell* 15 (2003) 1717–1727.
- [67] L. Ragni, E. Belles-Boix, M. Günl, V. Pautot, Interaction of KNAT6 and KNAT2 with *BREVIPEDICELLUS* and *PENNYWISE* in *Arabidopsis* inflorescences, *Plant Cell* 20 (2008) 888–900.
- [68] M. Khan, M. Xu, J. Murmu, P. Tabb, Y. Liu, K. Storey, S.M. McKim, C.J. Douglas, S.R. Hepworth, Antagonistic interaction of BLADE-ON-PETIOLE1 and 2 with *BREVIPEDICELLUS* and *PENNYWISE* regulates *Arabidopsis* inflorescence architecture, *Plant Physiol.* 158 (2012) 946–960.
- [69] M. Khan, L. Ragni, P. Tabb, B.C. Salasini, S. Chatfield, R. Datla, X. Lock, X. Kuai, C. Després, M. Proveniers, Y.G. Cao, D.Q. Xiang, H. Morin, J.P. Rullière, S. Citerne, S.R. Hepworth, V. Pautot, Repression of lateral organ boundary genes by *PENNYWISE*

- and POUND-FOOLISH is essential for meristem maintenance and flowering in *Arabidopsis*, *Plant Physiol.* 169 (2015) 2166–2186.
- [70] F. Andrés, M. Romera-Branchat, R. Martínez-Gallegos, V. Patel, K. Schneeberger, S. Jang, J. Altmüller, P. Nürnberg, G. Coupland, Floral induction in *Arabidopsis* by FLOWERING LOCUS T requires direct repression of BLADE-ON-PETIOLE genes by the homeodomain protein PENNYWISE, *Plant Physiol.* 169 (2015) 2187–2199.
- [71] I. Searle, Y. He, F. Turck, C. Vincent, F. Fornara, S. Kröber, R.A. Amasino, G. Coupland, The transcription factor FLC confers a flowering response to vernalization by repressing meristem competence and systemic signaling in *Arabidopsis*, *Genes Dev.* 20 (2006) 898–912.
- [72] M. Proveniers, B. Rutjens, M. Brand, S. Smeekens, The *Arabidopsis* TALE homeobox gene ATH1 controls floral competency through positive regulation of FLC, *Plant J.* 52 (2007) 899–913.
- [73] K. Okada, J. Ueda, M.K. Komaki, C.J. Bell, Y. Shimura, Requirement of the auxin polar transport system in early stages of *Arabidopsis* floral bud formation, *Plant Cell* 3 (1991) 677–684.
- [74] G.K. Przemeck, J. Mattsson, C.S. Hardtke, Z.R. Sung, T. Berleth, Studies on the role of the *Arabidopsis* gene MONOPTEROS in vascular development and plant cell axialization, *Planta* 200 (1996) 229–237.
- [75] L. Gälweiler, C. Guan, A. Müller, E. Wisman, K. Mendgen, A. Yephremov, K. Palme, Regulation of polar auxin transport by AtPIN1 in *Arabidopsis* vascular tissue, *Science* 282 (1998) 2226–2230.
- [76] C.S. Hardtke, T. Berleth, The *Arabidopsis* gene MONOPTEROS encodes a transcription factor mediating embryo axis formation and vascular development, *EMBO J.* 17 (1998) 1405–1411.
- [77] N. Yamaguchi, M.F. Wu, C.M. Winter, M.C. Berns, S. Nole-Wilson, A. Yamaguchi, G. Coupland, B.A. Krizek, D. Wagner, A molecular framework for auxin-mediated initiation of flower primordia, *Dev. Cell* 24 (2013) 271–282.
- [78] M.F. Wu, N. Yamaguchi, J. Xiao, B. Bargmann, M. Estelle, Y. Sang, D. Wagner, Auxin-regulated chromatin switch directs acquisition of flower primordium founder fate, *Elife* 4 (2015), e09269.
- [79] F. Parcy, O. Nilsson, M.A. Busch, I. Lee, D. Weigel, A genetic framework for floral patterning, *Nature* 395 (1998) 561–566.
- [80] V. Gregis, F. Andrés, A. Sessa, R.F. Guerra, S. Simonini, J.L. Mateos, S. Torti, F. Zambelli, G.M. Prazzoli, K.N. Bjerkan, P.E. Grini, G. Pavesi, L. Colombo, G. Coupland, M.M. Kater, Identification of pathways directly regulated by SHORT VEGETATIVE PHASE during vegetative and reproductive development in *Arabidopsis*, *Genome Biol.* 14 (2013) R56.
- [81] J. Lee, M. Oh, H. Park, I. Lee, SOC1 translocated to the nucleus by interaction with AGL24 directly regulates leafy, *Plant J.* 55 (2008) 832–843.
- [82] L.A. Sadic, B. Huvermann, S. Bezhami, Y. Su, C.M. Winter, C.S. Kwon, R.P. Collum, D. Wagner, The LEAFY target LMI1 is a meristem identity regulator and acts together with LEAFY to regulate expression of CAULIFLOWER, *Development* 133 (2006) 1673–1682.
- [83] S.A. Kempin, B. Savidge, M.F. Yanofsky, Molecular basis of the cauliflower phenotype in *Arabidopsis*, *Science* 267 (1995) 522–525.
- [84] D. Wagner, R.W. Sablowski, E.M. Meyerowitz, Transcriptional activation of APETALA1 by LEAFY, *Science* 285 (1999) 582–584.
- [85] J.J. Pastore, A. Limpuangthip, N. Yamaguchi, M.F. Wu, Y. Sang, S.K. Han, L. Malaspina, N. Chavdaroff, A. Yamaguchi, D. Wagner, LATE MERISTEM IDENTITY2 acts together with LEAFY to activate APETALA1, *Development* 138 (2011) 3189–3198.
- [86] C.M. Winter, R.S. Austin, S. Blanvillain-Baufumé, M.A. Reback, M. Monniaux, M.F. Wu, Y. Sang, A. Yamaguchi, N. Yamaguchi, J.E. Parker, F. Parcy, S.T. Jensen, H. Li, D. Wagner, LEAFY target genes reveal floral regulatory logic, cis motifs, and a link to biotic stimulus response, *Dev. Cell* 20 (2011) 430–443.
- [87] K. Kaufmann, F. Wellmer, J.M. Muñio, T. Ferrier, S.E. Wuest, V. Kumar, A. Serrano-Mislata, F. Madueño, P. Krajewski, E.M. Meyerowitz, G.C. Angenent, J.L. Riechmann, Orchestration of floral initiation by APETALA1, *Science* 328 (2010) 85–89.
- [88] D.A. William, Y. Su, M.R. Smith, M. Lu, D.A. Baldwin, D. Wagner, Genomic identification of direct target genes of LEAFY, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 1775–1780.
- [89] A. Pajoro, P. Madrigal, J.M. Muñio, J.T. Matus, J. Jin, M.A. Mecchia, J.M. Debernardi, J.F. Palatnik, S. Balazadeh, M. Arif, D.S. O'Maoileáidigh, F. Wellmer, P. Krajewski, J.L. Riechmann, G.C. Angenent, K. Kaufmann, Dynamics of chromatin accessibility and gene regulation by MADS-domain transcription factors in flower development, *Genome Biol.* 15 (2014) R41.
- [90] C.M. Winter, N. Yamaguchi, M.F. Wu, D. Wagner, Transcriptional programs regulated by both LEAFY and APETALA1 at the time of flower formation, *Physiol. Plant.* 155 (2015) 55–73.
- [91] M.A. Blazquez, R. Green, O. Nilsson, M.R. Sussman, D. Weigel, Gibberellins promote flowering of *Arabidopsis* by activating the LEAFY promoter, *Plant Cell* 10 (1998) 791–800.
- [92] S. Yu, V.C. Galvão, Y.C. Zhang, D. Horrer, T.Q. Zhang, Y.H. Hao, Y.Q. Feng, S. Wang, M. Schmid, J.W. Wang, Gibberellin regulates the *Arabidopsis* floral transition through miR156-targeted SQUAMOSA promoter binding-like transcription factors, *Plant Cell* 24 (2012) 3320–3332.
- [93] N. Yamaguchi, C.M. Winter, M.F. Wu, Y. Kanno, A. Yamaguchi, M. Seo, D. Wagner, Gibberellin acts positively then negatively to control onset of flower formation in *Arabidopsis*, *Science* 344 (2014) 638–641.
- [94] T. Nomura, H. Magome, A. Hanada, N. Takeda-Kamiya, L.N. Mander, Y. Kamiya, S. Yamaguchi, Functional analysis of *Arabidopsis* CYP714A1 and CYP714A2 reveals that they are distinct gibberellin modification enzymes, *Plant Cell Physiol.* 54 (2013) 1837–1851.
- [95] M.A. Mandel, C. Gustafson-Brown, B. Savidge, M.F. Yanofsky, Molecular characterization of the *Arabidopsis* floral homeotic gene APETALA1, *Nature* 360 (1992) 273–277.
- [96] Y. Han, C. Zhang, H. Yang, Y. Jiao, Cytokinin pathway mediates APETALA1 function in the establishment of determinate floral meristems in *Arabidopsis*, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) 6840–6845.
- [97] S. Scofield, W. Dewitte, J. Nieuwland, J.A. Murray, The *Arabidopsis* homeobox gene SHOOT MERISTEMLESS has cellular and meristem-organisational roles with differential requirements for cytokinin and CYCD3 activity, *Plant J.* 75 (2013) 53–66.
- [98] A. Leibfried, J.P. To, W. Busch, S. Stehling, A. Kehle, M. Demar, J.J. Kieber, J.U. Lohmann, WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response regulators, *Nature* 438 (2005) 1172–1175.
- [99] D.R. Smyth, J.L. Bowman, E.M. Meyerowitz, Early flower development in *Arabidopsis*, *Plant Cell* 2 (1990) 755–767.
- [100] S. de Folter, R.G. Immink, M. Kieffer, L. Parenicová, S.R. Henz, D. Weigel, M. Busscher, M. Kooiker, L. Colombo, M.M. Kater, B. Davies, G.C. Angenent, Comprehensive interaction map of the *Arabidopsis* MADS box transcription factors, *Plant Cell* 17 (2005) 1424–1433.
- [101] V. Gregis, A. Sessa, L. Colombo, M.M. Kater, AGL24, SHORT VEGETATIVE PHASE, and APETALA1 redundantly control AGAMOUS during early stages of flower development in *Arabidopsis*, *Plant Cell* 18 (2006) 1373–1382.
- [102] V. Gregis, A. Sessa, C. Dorca-Fornell, M.M. Kater, The *Arabidopsis* floral meristem identity genes AP1, AGL24 and SVP directly repress class B and C floral homeotic genes, *Plant J.* 60 (2009) 626–637.
- [103] R.G. Franks, C. Wang, J.Z. Levin, Z. Liu, SEUSS, a member of a novel family of plant regulatory proteins, represses floral homeotic gene expression with LEUNIG, *Development* 129 (2002) 253–263.
- [104] V.V. Sridhar, A. Surendrarao, D. Gonzalez, R.S. Conlan, Z. Liu, Transcriptional repression of target genes by LEUNIG and SEUSS, two interacting regulatory proteins for *Arabidopsis* flower development, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 11494–11499.
- [105] C. Castillejo, M. Romera-Branchat, S. Pelaz, A new role of the *Arabidopsis* SEPALLATA3 gene revealed by its constitutive expression, *Plant J.* 43 (2005) 586–596.
- [106] K. Kaufmann, J.M. Muñio, R. Jauregui, C.A. Airoldi, C. Smaczniak, P. Krajewski, G.C. Angenent, Target genes of the MADS transcription factor SEPALLATA3: integration of developmental and hormonal pathways in the *Arabidopsis* flower, *PLoS Biol.* 7 (2009), e1000090.
- [107] C. Liu, J. Zhou, K. Bracha-Drori, S. Yalovsky, T. Ito, H. Yu, Specification of *Arabidopsis* floral meristem identity by repression of flowering time genes, *Development* 134 (2007) 1901–1910.
- [108] V.V. Sridhar, A. Surendrarao, Z. Liu, APETALA1 and SEPALLATA3 interact with SEUSS to mediate transcription repression during flower development, *Development* 133 (2006) 3159–3166.
- [109] V. Grandi, V. Gregis, M.M. Kater, Uncovering genetic and molecular interactions among floral meristem identity genes in *Arabidopsis thaliana*, *Plant J.* 69 (2012) 881–893.
- [110] H. Yu, T. Ito, F. Wellmer, E.M. Meyerowitz, Repression of AGAMOUS-LIKE 24 is a crucial step in promoting flower development, *Nat. Genet.* 36 (2004) 157–161.
- [111] G. Theissen, H. Saedler, Plant biology. Floral quartets, *Nature* 409 (2001) 469–471.
- [112] T. Honma, K. Goto, Complexes of MADS-box proteins are sufficient to convert leaves into floral organs, *Nature* 409 (2001) 525–529.
- [113] S. Pelaz, R. Tapia-López, E.R. Alvarez-Buylla, M.F. Yanofsky, Conversion of leaves into petals in *Arabidopsis*, *Curr. Biol.* 11 (2001) 182–184.
- [114] C. Smaczniak, R.G. Immink, J.M. Muñio, R. Blanvillain, M. Busscher, J. Busscher-Lange, Q.D. Ding, S. Liu, A.H. Westphal, S. Boeren, F. Parcy, L. Xu, C.C. Carles, G.C. Angenent, K. Kaufmann, Characterization of MADS-domain transcription factor complexes in *Arabidopsis* flower development, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 1560–1565.
- [115] M. Egea-Cortines, H. Saedler, H. Sommer, Ternary complex formation between the MADS-box proteins SQUAMOSA, DEFICIENS and GLOBOSA is involved in the control of floral architecture in *Antirrhinum majus*, *EMBO J.* 18 (1999) 5370–5379.
- [116] R. Melzer, G. Theissen, Reconstitution of 'floral quartets' in vitro involving class B and class E floral homeotic proteins, *Nucleic Acids Res.* 37 (2009) 2723–2736.
- [117] M.A. Mendes, R.F. Guerra, M.C. Berns, C. Manzo, S. Masiero, L. Finzi, M.M. Kater, L. Colombo, MADS domain transcription factors mediate short-range DNA looping that is essential for target gene expression in *Arabidopsis*, *Plant Cell* 25 (2013) 2560–2572.
- [118] R.G. Immink, I.A. Tonaco, S. de Folter, A. Shchennikova, A.D. van Dijk, J. Busscher-Lange, J.W. Borst, G.C. Angenent, SEPALLATA3: the 'glue' for MADS box transcription factor complex formation, *Genome Biol.* 10 (2009) R24.
- [119] K. Jetha, G. Theissen, R. Melzer, *Arabidopsis* SEPALLATA proteins differ in cooperative DNA-binding during the formation of floral quartet-like complexes, *Nucleic Acids Res.* 42 (2014) 10927–10942.
- [120] V. Brambilla, R. Battaglia, M. Colombo, S. Masiero, S. Bencivenga, M.M. Kater, L. Colombo, Genetic and molecular interactions between BELL1 and MADS box factors support ovule development in *Arabidopsis*, *Plant Cell* 19 (2007) 2544–2556.
- [121] M.F. Wu, Y. Sang, S. Bezhami, N. Yamaguchi, S.K. Han, Z. Li, Y. Su, T.L. Siewinski, D. Wagner, SWI2/SNF2 chromatin remodeling ATPases overcome polycomb repression and control floral organ identity with the LEAFY and SEPALLATA3 transcription factors, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 3576–3581.
- [122] C. Li, C. Chen, L. Gao, S. Yang, V. Nguyen, X. Shi, K. Siminovich, S.E. Kohalmi, S. Huang, K. Wu, X. Chen, Y. Cui, The *Arabidopsis* SWI2/SNF2 chromatin remodeler BRAHMA regulates polycomb function during vegetative development and directly activates the flowering repressor gene SVP, *PLoS Genet.* 11 (2015), e1004944.
- [123] L. Yant, J. Mathieu, T.T. Dinh, F. Ott, C. Lanz, H. Wollmann, X. Chen, M. Schmid, Orchestration of the floral transition and floral development in *Arabidopsis* by the bi-functional transcription factor APETALA2, *Plant Cell* 22 (2010) 2156–2170.
- [124] S.E. Wuest, D.S. O'Maoileáidigh, L. Rae, K. Kwasniewska, A. Raganelli, K. Hanczaryk, A.J. Lohan, B. Loftus, E. Graciet, F. Wellmer, Molecular basis for the specification of floral organs by APETALA3 and PISTILLATA, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 13452–13457.
- [125] D.S. O'Maoileáidigh, S.E. Wuest, L. Rae, A. Raganelli, P.T. Ryan, K. Kwasniewska, P. Das, A.J. Lohan, B. Loftus, E. Graciet, F. Wellmer, Control of reproductive floral organ identity specification in *Arabidopsis* by the C function regulator AGAMOUS, *Plant Cell* 25 (2013) 2482–2503.

- [126] E. Moyroud, E.G. Minguet, F. Ott, L. Yant, D. Posé, M. Monniaux, S. Blanchet, O. Bastien, E. Thévenon, D. Weigel, M. Schmid, F. Parcy, Prediction of regulatory interactions from genome sequences using a biophysical model for the *Arabidopsis* LEAFY transcription factor, *Plant Cell* 23 (2011) 1293–1306.
- [127] K. Schiessl, J.M. Muiño, R. Sablowski, *Arabidopsis* JAGGED links floral organ patterning to tissue growth by repressing Kip-related cell cycle inhibitors, *Proc. Natl. Acad. Sci. U. S. A.* 111 (2014) 2830–2835.
- [128] B. Davies, M. Egea-Cortines, E. de Andrade Silva, H. Saedler, H. Sommer, Multiple interactions amongst floral homeotic MADS box proteins, *EMBO J.* 15 (1996) 4330–4343.
- [129] J.L. Riechmann, B.A. Krizek, E.M. Meyerowitz, Dimerization specificity of *Arabidopsis* MADS domain homeotic proteins APETALA1, APETALA3, PISTILLATA, and AGAMOUS, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 4793–4798.
- [130] T. Jack, L.L. Brockman, E.M. Meyerowitz, The homeotic gene APETALA3 of *Arabidopsis thaliana* encodes a MADS box and is expressed in petals and stamens, *Cell* 68 (1992) 683–697.
- [131] K. Goto, E.M. Meyerowitz, Function and regulation of the *Arabidopsis* floral homeotic gene PISTILLATA, *Genes Dev.* 8 (1994) 1548–1560.
- [132] E. Chae, Q.K. Tan, T.A. Hill, V.F. Irish, An *Arabidopsis* F-box protein acts as a transcriptional co-factor to regulate floral development, *Development* 135 (2008) 1235–1245.
- [133] F. Brioudes, C. Joly, J. Szécsi, E. Varaud, J. Leroux, F. Bellvert, C. Bertrand, M. Bendahmane, Jasmonate controls late development stages of petal growth in *Arabidopsis thaliana*, *Plant J.* 60 (2009) 1070–1080.
- [134] J. Szécsi, C. Joly, K. Bordji, E. Varaud, J.M. Cock, C. Dumas, M. Bendahmane, BIGPETALp, a bHLH transcription factor is involved in the control of *Arabidopsis* petal size, *EMBO J.* 25 (2006) 3912–3920.
- [135] E. Varaud, F. Brioudes, J. Szécsi, J. Leroux, S. Brown, C. Perrot-Rechenmann, M. Bendahmane, AUXIN RESPONSE FACTOR8 regulates *Arabidopsis* petal growth by interacting with the bHLH transcription factor BIGPETALp, *Plant Cell* 23 (2011) 973–983.
- [136] P. Nagpal, C.M. Ellis, H. Weber, S.E. Ploense, L.S. Barkawi, T.J. Guilfoyle, G. Hagen, J.M. Alonso, J.D. Cohen, E.E. Farmer, J.R. Ecker, J.W. Reed, Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation, *Development* 132 (2005) 4107–4118.
- [137] R. Tabata, M. Ikezaki, T. Fujibe, M. Aida, C.E. Tian, Y. Ueno, K.T. Yamamoto, Y. Machida, K. Nakamura, S. Ishiguro, *Arabidopsis* auxin response factor6 and 8 regulate jasmonic acid biosynthesis and floral organ development via repression of class 1 KNOX genes, *Plant Cell Physiol.* 51 (2010) 164–175.
- [138] R. Richter, C. Behringer, M. Zourelidou, C. Schwechheimer, Convergence of auxin and gibberellin signaling on the regulation of the GATA transcription factors GNC and GNL in *Arabidopsis thaliana*, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 13192–13197.
- [139] Y.M. Bi, Y. Zhang, T. Signorelli, R. Zhao, T. Zhu, S. Rothstein, Genetic analysis of *Arabidopsis* GATA transcription factor gene family reveals a nitrate-inducible member important for chlorophyll synthesis and glucose sensitivity, *Plant J.* 44 (2005) 680–692.
- [140] C.D. Mara, V.F. Irish, Two GATA transcription factors are downstream effectors of floral homeotic gene action in *Arabidopsis*, *Plant Physiol.* 147 (2008) 707–718.
- [141] R. Richter, C. Behringer, I.K. Müller, C. Schwechheimer, The GATA-type transcription factors GNC and GNL/CGA1 repress gibberellin signaling downstream from DELLA proteins and PHYTOCHROME-INTERACTING FACTORS, *Genes Dev.* 24 (2010) 2093–2104.
- [142] W. Yan, D. Chen, K. Kaufmann, Molecular mechanisms of floral organ specification by MADS domain proteins, *Curr. Opin. Plant Biol.* 29 (2016) 154–162.
- [143] C. Gaillochet, J.U. Lohmann, The never-ending story: from pluripotency to plant developmental plasticity, *Development* 142 (2015) 2237–2249.
- [144] A.L. Holt, J.M. van Haperen, E.P. Groot, T. Laux, Signaling in shoot and flower meristems of *Arabidopsis thaliana*, *Curr. Opin. Plant Biol.* 17 (2014) 96–102.
- [145] T. Laux, K.F. Mayer, J. Berger, G. Jürgens, The WUSCHEL gene is required for shoot and floral meristem integrity in *Arabidopsis*, *Development* 122 (1996) 87–96.
- [146] K.F. Mayer, H. Schoof, A. Haecker, M. Lenhard, G. Jürgens, T. Laux, Role of WUSCHEL in regulating stem cell fate in the *Arabidopsis* shoot meristem, *Cell* 95 (1998) 805–815.
- [147] R.K. Yadav, M. Perales, J. Gruel, T. Girke, H. Jönsson, G.V. Reddy, WUSCHEL protein movement mediates stem cell homeostasis in the *Arabidopsis* shoot apex, *Genes Dev.* 25 (2011) 2025–2030.
- [148] B. Sun, T. Ito, Regulation of floral stem cell termination in *Arabidopsis*, *Front. Plant Sci.* 6 (2015) 17.
- [149] M. Lenhard, A. Bohnert, G. Jürgens, T. Laux, Termination of stem cell maintenance in *Arabidopsis* floral meristems by interactions between WUSCHEL and AGAMOUS, *Cell* 105 (2001) 805–814.
- [150] J.U. Lohmann, R.L. Hong, M. Hobe, M.A. Busch, F. Parcy, R. Simon, D. Weigel, A molecular link between stem cell regulation and floral patterning in *Arabidopsis*, *Cell* 105 (2001) 793–803.
- [151] D.S. O'Maoláidigh, E. Graciet, F. Wellmer, Gene networks controlling *Arabidopsis thaliana* flower development, *New Phytol.* 201 (2014) 16–30.
- [152] B. Sun, Y. Xu, K.H. Ng, T. Ito, A timing mechanism for stem cell maintenance and differentiation in the *Arabidopsis* floral meristem, *Genes Dev.* 23 (2009) 1791–1804.
- [153] P. Das, T. Ito, F. Wellmer, T. Vernoux, A. Dedieu, J. Traas, E.M. Meyerowitz, Floral stem cell termination involves the direct regulation of AGAMOUS by PERIANTHIA, *Development* 136 (2009) 1605–1611.
- [154] A.N. Wynn, A.A. Seaman, A.L. Jones, R.G. Franks, Novel functional roles for PERIANTHIA and SEUSS during floral organ identity specification, floral meristem termination, and gynoecial development, *Front. Plant Sci.* 5 (2014) 130.
- [155] X. Chen, E.M. Meyerowitz, HUA1 and HUA2 are two members of the floral homeotic AGAMOUS pathway, *Mol. Cell* 3 (1999) 349–360.
- [156] Y. Cheng, N. Kato, W. Wang, J. Li, X. Chen, Two RNA binding proteins, HEN4 and HUA1, act in the processing of AGAMOUS pre-mRNA in *Arabidopsis thaliana*, *Dev. Cell* 4 (2003) 53–66.
- [157] E. Rodríguez-Cazorla, J.J. Ripoll, A. Andújar, L.J. Bailey, A. Martínez-Laborda, M.F. Yanofsky, A. Vera, K-homology nuclear ribonucleoproteins regulate floral organ identity and determinacy in *Arabidopsis*, *PLoS Genet.* 11 (2015), e1004983.
- [158] J.H. Jung, S. Lee, J. Yun, M. Lee, C.M. Park, The miR172 target TOE3 represses AGAMOUS expression during *Arabidopsis* floral patterning, *Plant Sci.* 215–216 (2014) 29–38.
- [159] S.E. Jacobsen, M.P. Running, E.M. Meyerowitz, Disruption of an RNA helicase/RNase III gene in *Arabidopsis* causes unregulated cell division in floral meristems, *Development* 126 (1999) 5231–5243.
- [160] W. Park, J. Li, R. Song, J. Messing, X. Chen, CARPEL FACTORY, a dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*, *Curr. Biol.* 12 (2002) 1484–1495.
- [161] L. Ji, X. Liu, J. Yan, W. Wang, R.E. Yumul, Y.J. Kim, T.T. Dinh, J. Liu, X. Cui, B. Zheng, M. Agarwal, C. Liu, X. Cao, G. Tang, X. Chen, ARGONAUTE10 and ARGONAUTE1 regulate the termination of floral stem cells through two microRNAs in *Arabidopsis*, *PLoS Genet.* 7 (2011), e1001358.
- [162] T.Z. Berardini, G. Bollman, H. Sun, R.S. Poethig, Regulation of vegetative phase change in *Arabidopsis thaliana* by cyclophilin 40, *Science* 291 (2001) 2405–2407.
- [163] N. Prunet, P. Morel, A.M. Thierry, Y. Eshed, J.L. Bowman, I. Negrutiu, C. Trehin, REBELOTE, SQUINT, and ULTRAPETALA1 function redundantly in the temporal regulation of floral meristem termination in *Arabidopsis thaliana*, *Plant Cell* 20 (2008) 901–919.
- [164] M.R. Smith, M.R. Willmann, G. Wu, T.Z. Berardini, B. Möller, D. Weijers, R.S. Poethig, Cyclophilin 40 is required for microRNA activity in *Arabidopsis*, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 5424–5429.
- [165] K.W. Earley, R.S. Poethig, Binding of the cyclophilin 40 ortholog SQUINT to Hsp90 protein is required for SQUINT function in *Arabidopsis*, *J. Biol. Chem.* 286 (2011) 38184–38189.
- [166] N. Prunet, P. Morel, P. Champelovier, A.M. Thierry, I. Negrutiu, T. Jack, C. Trehin, SQUINT promotes stem cell homeostasis and floral meristem termination in *Arabidopsis* through APETALA2 and CLAVATA signalling, *J. Exp. Bot.* 66 (2015) 6905–6916.
- [167] F. Moreau, E. Thévenon, R. Blanvillain, I. Lopez-Vidriero, J.M. Franco-Zorrilla, R. Dumas, F. Parcy, P. Morel, C. Trehin, C.C. Carles, The Myb-domain protein ULTRAPETALA1 INTERACTING FACTOR 1 controls floral meristem activities in *Arabidopsis*, *Development* 143 (2016) 1108–1119.
- [168] X. Liu, L. Gao, T.T. Dinh, T. Shi, D. Li, R. Wang, L. Guo, L. Xiao, X. Chen, DNA topoisomerase I affects polycomb group protein-mediated epigenetic regulation and plant development by altering nucleosome distribution in *Arabidopsis*, *Plant Cell* 26 (2014) 2803–2817.
- [169] D. Garcia, S.A. Collier, M.E. Byrne, R.A. Martienssen, Specification of leaf polarity in *Arabidopsis* via the trans-acting siRNA pathway, *Curr. Biol.* 16 (2006) 933–938.
- [170] D.R. Kelley, A. Arreola, T.L. Gallagher, C.S. Gasser, ETTIN (ARF3) physically interacts with KANADI proteins to form a functional complex essential for integument development and polarity determination in *Arabidopsis*, *Development* 139 (2012) 1105–1109.
- [171] X. Liu, T.T. Dinh, D. Li, B. Shi, Y. Li, X. Cao, L. Guo, Y. Pan, Y. Jiao, X. Chen, AUXIN RESPONSE FACTOR 3 integrates the functions of AGAMOUS and APETALA2 in floral meristem determinacy, *Plant J.* 80 (2014) 629–641.
- [172] J. Alvarez, D.R. Smyth, CRABS CLAW and SPATULA, two *Arabidopsis* genes that control carpel development in parallel with AGAMOUS, *Development* 126 (1999) 2377–2386.
- [173] J.L. Bowman, H. Sakai, T. Jack, D. Weigel, U. Mayer, E.M. Meyerowitz, SUPERMAN, a regulator of floral homeotic genes in *Arabidopsis*, *Development* 114 (1992) 599–615.
- [174] H. Sakai, L.J. Medrano, E.M. Meyerowitz, Role of SUPERMAN in maintaining *Arabidopsis* floral whorl boundaries, *Nature* 378 (1995) 199–203.
- [175] S. Simonini, I. Roig-Villanova, V. Gregis, B. Colombo, L. Colombo, M.M. Kater, Basic pentacysteine proteins mediate MADS domain complex binding to the DNA for tissue-specific expression of target genes in *Arabidopsis*, *Plant Cell* 24 (2012) 4163–4172.
- [176] T. Qi, H. Huang, S. Song, D. Xie, Regulation of jasmonate-mediated stamen development and seed production by a bHLH-MYB complex in *Arabidopsis*, *Plant Cell* 27 (2015) 1620–1633.
- [177] J. Davila-Velderrain, J.C. Martinez-Garcia, E.R. Alvarez-Buylla, Dynamic network modelling to understanding flowering transition and floral patterning, *J. Exp. Bot.* (2016).
- [178] T. Payne, S.D. Johnson, A.M. Koltunow, KNUCKLES (KNU) encodes a C2H2 zinc-finger protein that regulates development of basal pattern elements of the *Arabidopsis* gynoecium, *Development* 131 (2004) 3737–3749.
- [179] J.W. Chandler, B. Jacobs, M. Cole, P. Comelli, W. Werr, DORNROSCHEN-LIKE expression marks *Arabidopsis* floral organ founder cells and precedes auxin response maxima, *Plant Mol. Biol.* 76 (2011) 171–185.
- [180] M.C. Schruff, M. Spielman, S. Tiwari, S. Adams, N. Fenby, R.J. Scott, The AUXIN RESPONSE FACTOR 2 gene of *Arabidopsis* links auxin signalling, cell division, and the size of seeds and other organs, *Development* 133 (2006) 251–261.
- [181] C.A. MacAlister, S.J. Park, K. Jiang, F. Marcel, A. Bendahmane, Y. Izkovich, Y. Eshed, Z.B. Lippman, Synchronization of the flowering transition by the tomato TERMINATING FLOWER gene, *Nat. Genet.* 44 (2012) 1393–1398.
- [182] J.L. Bowman, D.R. Smyth, E.M. Meyerowitz, Genes directing flower development in *Arabidopsis*, *Plant Cell* 1 (1989) 37–52.
- [183] A. Nag, Y. Yang, T. Jack, DORNROSCHEN-LIKE, an AP2 gene, is necessary for stamen emergence in *Arabidopsis*, *Plant Mol. Biol.* 65 (2007) 219–232.
- [184] N.T. Krogan, K. Hogan, J.A. Long, APETALA2 negatively regulates multiple floral organ identity genes in *Arabidopsis* by recruiting the co-repressor TOPLESS and the histone deacetylase HDA19, *Development* 139 (2012) 4180–4190.
- [185] A. Horstman, V. Willemsen, K. Boutilier, R. Heidstra, AINTEGUMENTA-LIKE proteins: hubs in a plethora of networks, *Trends Plant Sci.* 19 (2014) 146–157.
- [186] R. Sablowski, Control of patterning, growth, and differentiation by floral organ identity genes, *J. Exp. Bot.* 66 (2015) 1065–1073.

- [187] N. Yamaguchi, C.W. Jeong, S. Nole-Wilson, B.A. Krizek, D. Wagner, AINTEGUMENTA and AINTEGUMENTA-LIKE6/PLETHORA3 induce LEAFY expression in response to auxin to promote the onset of flower formation in *Arabidopsis*, *Plant Physiol.* 170 (2016) 283–293.
- [188] J.P. Alvarez, I. Pekker, A. Goldshmidt, E. Blum, Z. Amsellem, Y. Eshed, Endogenous and synthetic microRNAs stimulate simultaneous, efficient, and localized regulation of multiple targets in diverse species, *Plant Cell* 18 (2006) 1134–1151.
- [189] Y. Okushima, I. Mitina, H.L. Quach, A. Theologis, AUXIN RESPONSE FACTOR 2 (ARF2): a pleiotropic developmental regulator, *Plant J.* 43 (2005) 29–46.
- [190] E. Zhu, C. You, S. Wang, J. Cui, B. Niu, Y. Wang, J. Qi, H. Ma, F. Chang, The DYT1-interacting proteins bHLH010, bHLH089 and bHLH091 are redundantly required for *Arabidopsis* anther development and transcriptome, *Plant J.* 83 (2015) 976–990.
- [191] J. Xu, Z. Ding, G. Vizcay-Barrena, J. Shi, W. Liang, Z. Yuan, D. Werck-Reichhart, L. Schreiber, Z.A. Wilson, D. Zhang, ABORTED MICROSPORES acts as a master regulator of pollen wall formation in *Arabidopsis*, *Plant Cell* 26 (2014) 1544–1556.
- [192] G.B. Seymour, L. Østergaard, N.H. Chapman, S. Knapp, C. Martin, Fruit development and ripening, *Annu. Rev. Plant Biol.* 64 (2013) 219–241.
- [193] C. Schuster, C. Gaillochet, J.U. Lohmann, *Arabidopsis* HECATE genes function in phytohormone control during gynoecium development, *Development* 142 (2015) 3343–3350.
- [194] S. Li, A. Lauri, M. Ziemann, A. Busch, M. Bhavé, S. Zachgo, Nuclear activity of ROXY1, a glutaredoxin interacting with TGA factors, is required for petal development in *Arabidopsis thaliana*, *Plant Cell* 21 (2009) 429–441.
- [195] J. Murmu, M.J. Bush, C. DeLong, S. Li, M. Xu, M. Khan, C. Malcolmson, P.R. Fobert, S. Zachgo, S.R. Hepworth, *Arabidopsis* basic leucine-zipper transcription factors TGA9 and TGA10 interact with floral glutaredoxins ROXY1 and ROXY2 and are redundantly required for anther development, *Plant Physiol.* 154 (2010) 1492–1504.
- [196] Q.L. Ranftl, E. Bastakis, C. Klermund, C. Schwechheimer, LLM-domain containing B-GATA factors control different aspects of cytokinin-regulated development in *Arabidopsis thaliana*, *Plant Physiol.* 170 (2016) 2295–2311.
- [197] L. Ding, S. Yan, L. Jiang, W. Zhao, K. Ning, J. Zhao, X. Liu, J. Zhang, Q. Wang, X. Zhang, HANABA TARANU (HAN) bridges meristem and organ primordia boundaries through PINHEAD, JAGGED, BLADE-ON-PETIOLE2 and CYTOKININ OXIDASE 3 during flower development in *Arabidopsis*, *PLoS Genet.* 11 (2015), e1005479.
- [198] X. Zhang, Y. Zhou, L. Ding, Z. Wu, R. Liu, E.M. Meyerowitz, Transcription repressor HANABA TARANU controls flower development by integrating the actions of multiple hormones, floral organ specification genes, and GATA3 family genes in *Arabidopsis*, *Plant Cell* 25 (2013) 83–101.
- [199] M. Cucinotta, L. Colombo, I. Roig-Villanova, Ovule development, a new model for lateral organ formation, *Front. Plant Sci.* 5 (2014) 117.
- [200] E.R. Lampugnani, A. Kilinc, D.R. Smyth, Auxin controls petal initiation in *Arabidopsis*, *Development* 140 (2013) 185–194.
- [201] H.R. Pires, M.M. Monfared, E.A. Shemyakina, J.C. Fletcher, ULTRAPETALA *trxG* genes interact with KANADI transcription factor genes to regulate *Arabidopsis* gynoecium patterning, *Plant Cell* 26 (2014) 4345–4361.
- [202] A. Hay, M. Tsiantis, KNOX genes: versatile regulators of plant development and diversity, *Development* 137 (2010) 3153–3165.
- [203] E. Costanzo, C. Trehin, M. Vandenbussche, The role of WOX genes in flower development, *Ann. Bot.* 114 (2014) 1545–1553.
- [204] M.J. Prigge, D. Otsuga, J.M. Alonso, J.R. Ecker, G.N. Drews, S.E. Clark, Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development, *Plant Cell* 17 (2005) 61–76.
- [205] N. Kamata, A. Sugihara, Y. Komeda, T. Takahashi, Allele-specific effects of PDF2 on floral morphology in *Arabidopsis thaliana*, *Plant Signal. Behav.* 8 (2013), e27417.
- [206] V.M. Zúñiga-Mayo, N. Marsch-Martínez, S. de Folter, JAIBA, a class-II HD-ZIP transcription factor involved in the regulation of meristematic activity, and important for correct gynoecium and fruit development in *Arabidopsis*, *Plant J.* 71 (2012) 314–326.
- [207] C. Smaczniak, R.G. Immink, G.C. Angenent, K. Kaufmann, Developmental and evolutionary diversity of plant MADS-domain factors: insights from recent studies, *Development* 139 (2012) 3081–3098.
- [208] A.A. Millar, F. Gubler, The *Arabidopsis* GAMBY-like genes, MYB33 and MYB65, are microRNA-regulated genes that redundantly facilitate anther development, *Plant Cell* 17 (2005) 705–721.
- [209] X. Chen, H. Huang, T. Qi, B. Liu, S. Song, New perspective of the bHLH-MYB complex in jasmonate-regulated plant fertility in *Arabidopsis*, *Plant Signal. Behav.* 11 (2016), e1135280.
- [210] S. Xing, M. Salinas, A. Garcia-Molina, S. Höhmann, R. Berndtgen, P. Huijser, SPL8 and miR156-targeted SPL genes redundantly regulate *Arabidopsis* gynoecium differential patterning, *Plant J.* 75 (2013) 566–577.
- [211] I. Rubio-Somoza, D. Weigel, Coordination of flower maturation by a regulatory circuit of three microRNAs, *PLoS Genet.* 9 (2013), e1003374.
- [212] T. Huang, V.F. Irish, Temporal control of plant organ growth by TCP transcription factors, *Curr. Biol.* 25 (2015) 1765–1770.
- [213] A.H. Roeder, P.T. Tarr, C. Tobin, X. Zhang, V. Chickarmane, A. Cunha, E.M. Meyerowitz, Computational morphodynamics of plants: integrating development over space and time, *Nat. Rev. Mol. Cell Biol.* 12 (2011) 265–273.
- [214] D. Zhang, Z. Yuan, Molecular control of grass inflorescence development, *Annu. Rev. Plant Biol.* 65 (2014) 553–578.
- [215] C. Périlleux, G. Lobet, P. Tocquin, Inflorescence development in tomato: gene functions within a zigzag model, *Front. Plant Sci.* 5 (2014) 121.
- [216] J.M. Muiño, S. de Bruijn, A. Pajoro, K. Geuten, M. Vingron, G.C. Angenent, K. Kaufmann, Evolution of DNA-binding sites of a floral master regulatory transcription factor, *Mol. Biol. Evol.* 33 (2016) 185–200.
- [217] J.X. Shi, S. Malitsky, S. De Oliveira, C. Branigan, R.B. Franke, L. Schreiber, A. Aharoni, SHINE transcription factors act redundantly to pattern the archetypal surface of *Arabidopsis* flower organs, *PLoS Genet.* 7 (2011) e1001388.