

Bloom to Véraison

This year-round program section covers the major pests of table grapes grown in California. For wine and raisin grapes, see [WINE AND RAISIN GRAPE YEAR-ROUND PROGRAM.\(/agriculture/grape/wine-and-raisin-grapes-bloom-to-vraison/\)](#)

About Bloom to Véraison

- San Joaquin Valley, early May to July; Coachella Valley, April
- [Mitigate pesticide effects\(/agriculture/grape/wine-and-raisin-grapes-pesticide-application-checklist/\)](#) on air and water quality.

What should you be doing during this time?

Monitor for [western flower thrips\(/agriculture/grape/Thrips/\)](#), particularly in vineyards near drying grains. Manage according to the Pest Management Guidelines.

[Monitor leafhoppers, spider mites, and mealybugs\(/agriculture/grape/Monitoring-Insects-and-Spider-Mites/\)](#) weekly.

- Keep records (example [monitoring form\(/PMG/C302/grape-leafhoprmite.pdf\)](#) PDF).
- Manage if needed according to the Pest Management Guidelines.

Examine leaves and shoots for [Botrytis bunch rot\(/agriculture/grape/Botrytis-Bunch-Rot/#MANAGEMENT\)](#) and [powdery mildew\(/agriculture/grape/Powdery-Mildew/#MANAGEMENT\)](#). Manage if needed according to the Pest Management Guidelines.

If [European fruit lecanium scale\(/agriculture/grape/European-Fruit-Lecanium-Scale/#MANAGEMENT\)](#) has been a problem in the past, monitor for egg hatch to time treatment.

Continue to check omnivorous leafroller [pheromone traps\(/agriculture/grape/Pheromone-Traps/\)](#) weekly. Keep records (example [monitoring form\(/PMG/C302/grape-insectdegree-days.pdf\)](#) PDF).

Continue monitoring pheromone traps for [vine mealybug\(/agriculture/grape/Vine-Mealybug/#MANAGEMENT\)](#).

- If males are caught or honeydew, sooty mold, or ants are found, look for nymphs/females on surrounding vines.
- Keep records (example [monitoring form\(/PMG/C302/grape-leafhoprmite.pdf\)](#) PDF).
- If [grape\(/agriculture/grape/Mealybugs-Pseudococcus/\)](#) or [vine mealybug\(/agriculture/grape/Vine-Mealybug/\)](#) nymphs/females are found on the vine, manage according to Pest Management Guidelines

Remove basal leaves and lateral shoots in the fruit zone beginning around berry set to minimize [summer rot\(/agriculture/grape/Summer-Bunch-Rot-Sour-Rot/#MANAGEMENT\)](#), [Botrytis bunch rot\(/agriculture/grape/Botrytis-Bunch-Rot/#MANAGEMENT\)](#), and [leafhopper\(/agriculture/grape/Leafhoppers/#MANAGEMENT\)](#) populations, and to maximize application coverage.

- Time leaf pull before first-generation grape leafhoppers become adults.

- Remove only the leaves on the shaded side of the canopy on non-divided trellis systems to prevent heat damage and sunburn of sensitive varieties (Thompson Seedless, Redglobe).

Treat for Botrytis before rain according to the Pest Management Guidelines.

[Monitor caterpillars\(/agriculture/grape/Monitoring-Caterpillars/\)](/agriculture/grape/Monitoring-Caterpillars/) if they have been a problem in the past:

- Omnivorous leafroller
- Grape leaffolder
- Western grapeleaf skeletonizer

Keep records (example [monitoring form\(/PMG/C302/grape-catmon.pdf\)](/PMG/C302/grape-catmon.pdf) PDF).

Monitor sticky traps for [glassy-winged sharpshooters\(/agriculture/grape/Sharpshooters/#MANAGEMENT\)](/agriculture/grape/Sharpshooters/#MANAGEMENT). Keep records (example [monitoring form\(/PMG/C302/grape-stickytrap.pdf\)](/PMG/C302/grape-stickytrap.pdf) PDF).

Other pests or pest damage you may see:

- [Grape thrips\(/agriculture/grape/Thrips/\)](/agriculture/grape/Thrips/)
- Grasshopper
- Whitefly
- [Black widow spiders\(/agriculture/grape/Black-Widow-Spider/\)](/agriculture/grape/Black-Widow-Spider/)
- [Red-headed and green sharpshooters\(/agriculture/grape/Sharpshooters/\)](/agriculture/grape/Sharpshooters/)
- [False chinch bug\(/agriculture/grape/False-Chinch-Bug/\)](/agriculture/grape/False-Chinch-Bug/)



UC IPM Pest Management Guidelines: Grape
UC ANR Publication 3448

[Authors and Credits \(./authors-and-credits\)](/authors-and-credits)

Text Updated: 07/15