

Agriculture: Grape Pest Management Guidelines

Budbreak

This year-round program section covers the major pests of table grapes grown in California. For wine and raisin grapes, see [WINE AND RAISIN GRAPE YEAR-ROUND PROGRAM.\(/agriculture/grape/wine-and-raisin-grapes-budbreak/\)](#)

About Budbreak

- San Joaquin Valley, March; Coachella Valley, January to February
- [Mitigate pesticide effects\(/agriculture/grape/wine-and-raisin-grapes-pesticide-application-checklist/\)](#) on air and water quality.

What should you be doing during this time?

On a warm day (above 65°F), [monitor trunks, cordons, and spurs\(/agriculture/grape/Delayed-Dormant-And-Budbreak-Monitoring-Table-Grapes/\)](#) for:

- Mealybugs
- Ants associated with mealybugs and European fruit lecanium scale
- Overwintering spider mites (orange)
- Cutworms

Keep records (example [monitoring form\(/PMG/C302/grape-spurmonitor.pdf\)](#) PDF). Manage if needed according to the Pest Management Guidelines.

In San Joaquin Valley continue to check [pheromone traps\(/agriculture/grape/Pheromone-Traps/\)](#) twice weekly for omnivorous leafroller, if biofix has not been reached.

- Record biofix for the first moth (example [monitoring form\(/PMG/C302/grape-insectdegree-days.pdf\)](#) PDF).
 - Check traps weekly after biofix date is established.
-

Monitor for [powdery mildew\(/agriculture/grape/Powdery-Mildew/#MANAGEMENT\)](#) using the risk assessment index followed by visual inspections.

Treat if needed according to the Pest Management Guidelines.

If rainfall is predicted after budbreak, consider treating for [phomopsis cane and leaf spot\(/agriculture/grape/Phomopsis-Cane-and-Leafspot/#MANAGEMENT\)](#) in sensitive varieties (Thompson Seedless, Redglobe).

Note locations of vines showing poor budbreak for future assessment of abiotic disorders or diseases.

Check sticky traps for [glassy-winged sharpshooters\(/agriculture/grape/Sharpshooters/#MANAGEMENT\)](#). Keep records (example [monitoring form\(/PMG/C302/grape-stickytrap.pdf\)](#) PDF).

[Survey weeds\(/PMG/C302/mt302scsprngweed.html\)](/PMG/C302/mt302scsprngweed.html) to plan a weed management strategy if not completed earlier in the season. If herbicides are to be used:

- Record observations (example [survey form\(/PMG/C302/grape-springweed.pdf\)](/PMG/C302/grape-springweed.pdf) PDF).
- Make your selection based on weed survey observations.

Other pests you may see:

- [Grape bud beetle\(/agriculture/grape/Grape-Bud-Beetle/\)](/agriculture/grape/Grape-Bud-Beetle/)
- [Red-headed and green sharpshooters\(/agriculture/grape/Sharpshooters/\)](/agriculture/grape/Sharpshooters/)



UC IPM Pest Management Guidelines: Grape
UC ANR Publication 3448

[Authors and Credits \(../authors-and-credits\)](/authors-and-credits)

Text Updated: 07/15