

Dormancy

This year-round program section covers the major pests of table grapes grown in California. For wine and raisin grapes, see the [WINE AND RAISIN GRAPE YEAR-ROUND PROGRAM\(/agriculture/grape/wine-and-raisin-grapes-dormancy/\)](#).

About Dormancy

- [Mitigate pesticide effects\(/agriculture/grape/wine-and-raisin-grapes-pesticide-application-checklist/\)](#) on air and water quality.

What should you be doing during this time?

In the San Joaquin Valley, sample for [nematodes\(/agriculture/grape/Nematodes/#EVALUATION\)](#) from November to February.

Carry out [preventative practices\(/PMG/C302/mt302rcvnydsntn.html\)](#) for trunk diseases:

- Use delayed pruning or double pruning in February or later.
- Use applications of pruning-wound protectants after pruning and before rain in November through January.

Carry out [dormant-season sanitation\(/PMG/C302/mt302rcvnydsntn.html\)](#) activities.

- Destroy prunings of older infested wood to reduce pest sources.
- Remove dried grape clusters on vines and disc weeds and clusters where orange tortrix or omnivorous leafroller is a problem.
- In vineyards with a history of branch and twig borers, examine old pruning scars and dead parts of vines for brown frass and wood dust.
- If you have vine mealybug, steam sanitize equipment before moving to uninfested area of the vineyard.

[Survey weeds\(/PMG/C302/mt302rcwntrweed.html\)](#) to plan a weed management strategy.

- Pay close attention to weeds that may present a special problem or management challenge.
 - Record your observations (example [survey form\(/PMG/C302/grape-winterweed.pdf\)](#) PDF).
 - Use records to make herbicide selections in vineyards where sprays are planned.
-



UC IPM Pest Management Guidelines: Grape
UC ANR Publication 3448

[Authors and Credits \(./authors-and-credits\)](#)

Text Updated: 07/15