

Most Effective Treatment Timings For Key Diseases

Note: Not all indicated timings may be necessary for disease control.

Disease	Fall	Delayed dormant	Green tip/ White bud	Pink bud/ Full bloom	Petal fall/ Cover sprays
Fire blight	0	1	2	3	3 ⁴
Powdery mildew ³	0	0	1	3	3
Scab ¹	2 ²	2 ²	3	3	3

Rating: 3 = most effective, 2 = moderately effective, 1= least effective, and 0 = ineffective

1	Protection of early tissue is important. Additional applications should be made according to infection periods as determined by the Mills table.
2	Disruption of pseudothecial (ascostroma) development (fall) and inactivation of overwintering twig lesions (delayed dormant); effects of these treatments on disease control is uncertain.
3	Early applications are most effective; additional applications are made if mildew continues.
4	Start management program at the beginning of bloom and continue through bloom including "rat-tail" bloom throughout spring. Several models are available for forecasting infection periods and treatment timing. Models include: Maryblyt, Cougar Blight, etc.

Acknowledgment: Adaskaveg et al., 2025. [Fungicides, Bactericides, Biocontrols, and Natural Products for Deciduous Tree Fruit and Nut, Citrus, Strawberry, and Vine Crops in California\(PDF/PMG/fungicideefficacytiming.pdf\)](#).

UC

PEER

REVIEWED

UC IPM Pest Management Guidelines: Apple

UC ANR Publication 3432

[Authors and Credits \(./Authors-and-Credits\)](#)

Text Updated: 07/24