

Preemergence

This year-round IPM program covers the major pests of both commercial and seed potato production fields in California.

About Preemergence

- [Mitigate pesticide effects\(/agriculture/potato/Pesticide-application-checklist/\)](/agriculture/potato/Pesticide-application-checklist/) on air and water quality.

What should you be doing during this time?

Apply [preemergence herbicide\(/agriculture/potato/Integrated-Weed-Management/#AFTER\)](/agriculture/potato/Integrated-Weed-Management/#AFTER) if needed, according to the Pest Management Guidelines. Consider combining a hilling operation to kill weeds already emerged followed by the herbicide application.

Manage soil moisture to minimize:

- [Fusarium seed piece decay\(/agriculture/potato/Fusarium-Dry-Rot-and-Seed-Piece-Decay/\)](/agriculture/potato/Fusarium-Dry-Rot-and-Seed-Piece-Decay/)
 - [Bacterial soft rot and blackleg\(/agriculture/potato/Bacterial-Soft-Rot-and-Blackleg/\)](/agriculture/potato/Bacterial-Soft-Rot-and-Blackleg/)
 - [Stem and stolon canker \(black scurf\)\(/agriculture/potato/Stem-and-Stolon-Canker/\)](/agriculture/potato/Stem-and-Stolon-Canker/)
-



UC IPM Pest Management Guidelines: Potato
UC ANR Publication 3463

[Authors and Credits \(./authors-and-credits\)](/authors-and-credits/)

Text Updated: 08/19