

Row Closure to Maturity

This year-round IPM program covers the major pests of both commercial and seed potato production fields in California.

About Row Closure to Maturity

- [Mitigate pesticide effects\(/agriculture/potato/Pesticide-application-checklist/\)](/agriculture/potato/Pesticide-application-checklist/) on air and water quality.

What should you be doing during this time?

Monitor [aphids\(/agriculture/potato/Aphids/\)](/agriculture/potato/Aphids/) weekly (Klamath Basin only). Keep records (example [monitoring form\(/PMG/C607/potato-aphidmon.pdf\)](/PMG/C607/potato-aphidmon.pdf) PDF) and manage if needed according to the Pest Management Guidelines.

Watch for [pests of foliage\(/PMG/C607/m607epfolpests.html\)](/PMG/C607/m607epfolpests.html), including:

Insects:

- [Aphids\(/agriculture/potato/Aphids/\)](/agriculture/potato/Aphids/)
- [Cutworms and loopers\(/agriculture/potato/Cutworms/\)](/agriculture/potato/Cutworms/)
- [Flea beetles\(/agriculture/potato/Flea-Beetles/\)](/agriculture/potato/Flea-Beetles/)
- [Psyllids\(/agriculture/potato/Potato-Psyllid/\)](/agriculture/potato/Potato-Psyllid/)
- [Tuberworm\(/agriculture/potato/Potato-Tuberworm/\)](/agriculture/potato/Potato-Tuberworm/)
- [Whiteflies\(/agriculture/potato/Silverleaf-Whitefly/\)](/agriculture/potato/Silverleaf-Whitefly/)

Diseases:

- [Blackleg\(/agriculture/potato/Bacterial-Soft-Rot-and-Blackleg/\)](/agriculture/potato/Bacterial-Soft-Rot-and-Blackleg/)
- [Early blight\(/agriculture/potato/Early-Blight/\)](/agriculture/potato/Early-Blight/)
- [Late blight\(/agriculture/potato/Late-Blight/\)](/agriculture/potato/Late-Blight/)
- [White mold\(/agriculture/potato/White-Mold/\)](/agriculture/potato/White-Mold/)

Note presence of pests or damage (example [monitoring form\(/PMG/C607/potato-foliagepests.pdf\)](/PMG/C607/potato-foliagepests.pdf) PDF) and manage according the Pest Management Guidelines.

Monitor for tuberworm adults. (Central and Southern California)

- Continue monitoring [pheromone traps\(/agriculture/potato/Potato-Tuberworm/#MONITORING\)](/agriculture/potato/Potato-Tuberworm/#MONITORING).
- Keep records (example [monitoring form\(/PMG/C607/potato-tuberwormmon.pdf\)](/PMG/C607/potato-tuberwormmon.pdf) PDF).

Apply insecticides if needed according to the Pest Management Guidelines.

Based on field history, manage diseases according to the Pest Management Guidelines.

- [Pink rot\(/agriculture/potato/Pink-Rot/\)](/agriculture/potato/Pink-Rot/)

- [Pythium leak\(/agriculture/potato/Leak-Water-Rot/\)](/agriculture/potato/Leak-Water-Rot/) (Central and Southern California)
- [Sclerotium stem rot\(/agriculture/potato/Sclerotium-Stem-Rot/\)](/agriculture/potato/Sclerotium-Stem-Rot/) (Southern blight)

If [white mold\(/agriculture/potato/White-Mold/\)](/agriculture/potato/White-Mold/) has been a problem in previous potato crops, apply fungicide at full flowering according to the Pest Management Guidelines.

Manage soil moisture to avoid soil cracking, but avoid over-irrigation to prevent diseases.

Consider application of nematicide via chemigation according to nematode lifecycle and label recommendations.

Apply fertilizer as needed according to soil and petiole tests. Avoid over- or under-fertilization (especially nitrogen).

Survey for weed escapes or late-emerging weeds. Apply postemergence herbicide if necessary.



UC IPM Pest Management Guidelines: Potato
UC ANR Publication 3463

[Authors and Credits \(./authors-and-credits/\)](/authors-and-credits/)

Text Updated: 08/19