

Mathematics for Information Technology

Somkiat Wangsiripitak

somkiat.wa@kmitl.ac.th

Room.518 or Room.506 (MIV Lab)

APPLICATION OF DIFFERENTIATION

Application of Differentiation

- Three basic categories of application.
 - curve sketching,
 - optimization, and
 - approximation techniques.

Extrema on an Interval

I

Objectives

ค่าสูง / ต่ำสุดในช่วงใดช่วงหนึ่ง

- Understand the definition of extrema of a function on an interval.
- Understand the definition of relative extrema of a function on an open interval.
- Find extrema on a closed interval.


Extrema of a Function

DEFINITION OF EXTREMA


Let f be defined on an interval I containing c .

- $f(c)$ is the **minimum** of f on I if $f(c) \leq f(x)$ for all x in I .
- $f(c)$ is the **maximum** of f on I if $f(c) \geq f(x)$ for all x in I .

The **minimum** and **maximum** of a function on an interval are the **extreme values**, or **extrema** (the singular form of extrema is **extremum**), of the function on the interval. The **minimum** and **maximum** of a function on an interval are also called the **absolute minimum** and **absolute maximum**, or the **global minimum** and **global maximum**, on the interval.



Extrema of a Function



(a) f is continuous, $[-1, 2]$ is closed. (b) f is continuous, $(-1, 2)$ is open. (c) g is not continuous, $[-1, 2]$ is closed.

ช่วงปิด, ต่อเนื่อง
หาได้ทั้งคี่



Extrema can occur at interior points or end-points of an interval. Extrema that occur at the endpoints are called **endpoint extrema**.

THEOREM 3.1 THE EXTREME VALUE THEOREM

If f is continuous on a closed interval $[a, b]$, then f has both a minimum and a maximum on the interval.

Relative Extrema and Critical Numbers

local maximum and local minimum



Local Extrema
ตัว


จุดที่ ค. ชัน = 0
หา ค. ชัน ไม่ได้
but
ไม่ต่อเนื่อง
Local เสมอไป

Such a hill and valley can occur in two ways. If the hill (or valley) is smooth and rounded, the graph has a **horizontal tangent line** at the high point (or low point). If the hill (or valley) is sharp and peaked, the graph represents a function that is **not differentiable** at the **high point** (or low point).

Relative Extrema and Critical Numbers

DEFINITION OF A CRITICAL NUMBER

Let f be defined at c . If $f'(c) = 0$ or if f is not differentiable at c , then c is a **critical number** of f .



ไม่ใช่ Local Extrema
เป็น critical num
↑
f' = 0

THEOREM 3.2 RELATIVE EXTREMA OCCUR ONLY AT CRITICAL NUMBERS

If f has a relative minimum or relative maximum at $x = c$, then c is a critical number of f .

an Extrema finden

To find the extrema of a continuous function f on a closed interval $[a, b]$, use the following steps.

- ပုဂ္ဂိုလ်တစ်ဦးချင်းနှင့် ၄ နှစ် အကုန် အများ အနည်း



Confirm with a graph.

On the closed interval $[-1, 2]$, f has a minimum at $(1, -1)$ and a maximum at $(2, 16)$.

note that the critical number $x = 0$ does not yield a relative minimum or a relative maximum. This tells you that the converse of Theorem 3.2 is not true. In other words, the critical numbers of a function need not produce relative extrema.

Critical Number

$$= 0.9666 \times 0$$

" อีฟ ไข่ไข่ "

Solution Begin by differentiating the function.

$$f(x) = 3x^4 - 4x^3$$

Write original function.

$$f'(x) = 12x^3 - 12x^2$$

Differentiate.

To find the critical numbers of f , you must find all x -values for which $f'(x) = 0$ and all x -values for which $f'(x)$ does not exist.

$$f'(x) = 12x^3 - 12x^2 = 0$$

Set $f'(x)$ equal to 0. ทำให้อนุพันธ์เท่ากับ 0

$$12x^2(x - 1) = 0$$

Factor. 664 ៣ ៩ ២ ៥ ៣ ០ ៧

$x = 0, 1$

Critical numbers

Because f' is defined for all x , you can conclude that these are the only critical numbers of f . By evaluating f at these two critical numbers and at the endpoints of $[-1, 2]$, you can determine that the maximum is $f(2) = 16$ and the minimum is $f(1) = -1$, as shown in the table.

Left Endpoint	Critical Number	Critical Number	Right Endpoint
$f(-1) = 7$	$f(0) = 0$	$f(1) = -1$ <u>Minimum</u>	$f(2) = 16$ <u>Maximum</u>

เทียบว่าค่าเงินบาทสูงทุกค่า

ហេតុនេះ យើងបាន $f(x)$

Find the extrema of $f(x) = 2x - 3x^{2/3}$ on the interval $[-1, 3]$.

Solution Begin by differentiating the function.

$$f'(x) = 2 - \frac{2}{x^{2/3}} = 2 \left(\frac{x^{1/3} - 1}{x^{1/3}} \right)$$


$$CN = 1, 0$$

cn $f'(1) = 2 - 3 = -1$

cn $f'(0) = 0 - 0 = 0 \longrightarrow \text{max im km}$

Left $f(-1) = 2(-1) - 3(-1) = -5 \rightarrow \text{minimum}$

Right $f(3) = 6 - 3\sqrt{9} \approx -0.24$



The Mean Value Theorem


THEOREM 3.4 THE MEAN VALUE THEOREM

If f is continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) , then there exists a number c in (a, b) such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

Geometrically, the theorem guarantees the existence of a tangent line that is parallel to the secant line through the points $(a, f(a))$ and $(b, f(b))$, as shown in Figure

In terms of rates of change, the Mean Value Theorem implies that there must be a point in the open interval (a, b) at which the instantaneous rate of change is equal to the average rate of change over the interval $[a, b]$. This is illustrated in Example 4.



Math for IT (Somkiat Wangsiripitak)

29

EXAMPLE 4 Finding an Instantaneous Rate of Change

Two stationary patrol cars equipped with radar are 5 miles apart on a highway, as shown in Figure. As a truck passes the first patrol car, its speed is clocked at 55 miles per hour. Four minutes later, when the truck passes the second patrol car, its speed is clocked at 50 miles per hour. Prove that the truck must have exceeded the speed limit (of 55 miles per hour) at some time during the 4 minutes.


Solution Let $t = 0$ be the time (in hours) when the truck passes the first patrol car. The time when the truck passes the second patrol car is

$$t = \frac{4}{60} = \frac{1}{15} \text{ hour.}$$

By letting $s(t)$ represent the distance (in miles) traveled by the truck, you have $s(0) = 0$ and $s(\frac{1}{15}) = 5$. So, the average velocity of the truck over the five-mile stretch of highway is

$$\text{Average velocity} = \frac{s(1/15) - s(0)}{(1/15) - 0} = \frac{5}{1/15} = 75 \text{ miles per hour.}$$

Assuming that the position function is differentiable, you can apply the Mean Value Theorem to conclude that the truck must have been traveling at a rate of 75 miles per hour sometime during the 4 minutes.



At some time t , the instantaneous velocity is equal to the average velocity over 4 minutes.

Math for IT (Somkiat Wangsiripitak)

First Derivative Test

Objectives

- Apply the First Derivative Test to find relative extrema of a function.

↳ บทนำสู่ Local Extrema

BREAK

Math for IT (Somkiat Wangsiripitak)

36

Math for IT (Somkiat Wangsiripitak)


37

The First Derivative Test

THEOREM 3.6 THE FIRST DERIVATIVE TEST

Let c be a critical number of a function f that is continuous on an open interval I containing c . If f is differentiable on the interval, except possibly at c , then $f(c)$ can be classified as follows.

- If $f'(x)$ changes from negative to positive at c , then f has a relative minimum at $(c, f(c))$.
- If $f'(x)$ changes from positive to negative at c , then f has a relative maximum at $(c, f(c))$.
- If $f'(x)$ is positive on both sides of c or negative on both sides of c , then $f(c)$ is neither a relative minimum nor a relative maximum.



44

EXAMPLE 3 Applying the First Derivative Test

Find the relative extrema of

$$f(x) = (x^2 - 4)^{2/3}$$

Solution Begin by noting that f is continuous on the entire real number line. The derivative of f

$$f'(x) = \left[\frac{2}{3} (x^2 - 4)^{-1/3} \right] (2x) = \frac{4x}{3(x^2 - 4)^{1/3}}$$

General Power Rule


Simplify.

$$x^2 - 4 = x^2 - 2^2 = (x - 2)(x + 2)$$

is 0 when $x = 0$ and does not exist when $x = \pm 2$. So, the critical numbers are $x = -2$, $x = 0$, and $x = 2$. The table summarizes the testing of the four intervals determined by these three critical numbers.

Interval	$-\infty < x < -2$	$-2 < x < 0$	$0 < x < 2$	$2 < x < \infty$
Test Value	$x = -3$	$x = -1$	$x = 1$	$x = 3$
Sign of $f'(x)$	$f'(-3) < 0$	$f'(-1) > 0$	$f'(1) < 0$	$f'(3) > 0$
Conclusion	Decreasing	Increasing	Decreasing	Increasing

By applying the First Derivative Test, you can conclude that f has a relative minimum at the point $(-2, 0)$, a relative maximum at the point $(0, \frac{3}{16})$, and another relative minimum at the point $(2, 0)$.



48

EXAMPLE 4 Applying the First Derivative Test

Find the relative extrema of $f(x) = \frac{x^4 + 1}{x^2}$

Solution

$$\begin{aligned} f(x) &= x^2 + x^{-2} \\ &= 2x - 2x^{-3} \\ &= 2x - \frac{2}{x^3} \\ &= \frac{2x^4 - 2}{x^3} \\ &= \frac{2(x^4 - 1)}{x^3} \\ &= \frac{2(x^2 + 1)(x - 1)(x + 1)}{x^3} \end{aligned}$$

$x = \pm 1$
 $x = 0$
 $x = 1$
 $x = -1$
 $x = 0$

EXAMPLE 4 Applying the First Derivative Test

Find the relative extrema of $f(x) = \frac{x^4 + 1}{x^2}$.

Solution

$f(x) = x^2 + x^{-2}$ Rewrite original function.
 $f'(x) = 2x - 2x^{-3}$ Differentiate.
 $= 2x - \frac{2}{x^3}$ Rewrite with positive exponent.
 $= \frac{2(x^4 - 1)}{x^3}$ Simplify.
 $= \frac{2(x^2 + 1)(x - 1)(x + 1)}{x^3}$ Factor.


So, $f'(x)$ is zero at $x = \pm 1$. Moreover, because $x = 0$ is not in the domain of f , you should use this x -value along with the critical numbers to determine the test intervals.

$x = \pm 1$ Critical numbers, $f'(\pm 1) = 0$
 $x = 0$ 0 is not in the domain of f .

The table summarizes the testing of the four intervals determined by these three x -values.

Interval	$-\infty < x < -1$	$-1 < x < 0$	$0 < x < 1$	$1 < x < \infty$
Test Value	$x = -2$	$x = -\frac{1}{2}$	$x = \frac{1}{2}$	$x = 2$
Sign of $f'(x)$	$f'(-2) < 0$	$f'(-\frac{1}{2}) > 0$	$f'(\frac{1}{2}) < 0$	$f'(2) > 0$
Conclusion	Decreasing	Increasing	Decreasing	Increasing

Local minimum
 Local maximum
 Local minimum
 not in domain



50

Math for IT (Somkiat Wangsiripitak)

52

Math for IT (Somkiat Wangsiripitak)

Optimization Problems

II

Objectives

หาค่าต่ำสุดที่สุด ex. ห่อของขวัญ / ภาชนะ

Solve applied minimum and maximum problems.

$$\begin{aligned} & [(x^2 + 144)^{\frac{1}{2}} + (x^2 - 60x + 1684)^{\frac{1}{2}}]' \\ & \frac{1}{2}(x^2 + 144)^{-\frac{1}{2}}(2x) + \frac{1}{2}(x^2 - 60x + 1684)^{-\frac{1}{2}}(2x - 60) \\ & \frac{x}{\sqrt{x^2 + 144}} + \frac{x - 30}{\sqrt{x^2 - 60x + 1684}} \end{aligned}$$

Math for IT (Somkiat Wangsiripitak)

53

GUIDELINES FOR SOLVING APPLIED MINIMUM AND MAXIMUM PROBLEMS

ให้มา / ให้มา

ต้องการให้มา / ต้องการให้มา

1. Identify all given quantities and all quantities to be determined. If possible, make a sketch.
2. Write a primary equation for the quantity that is to be maximized or minimized.
3. Reduce the primary equation to one having a single independent variable. This may involve the use of secondary equations relating the independent variables of the primary equation.
4. Determine the feasible domain of the primary equation. That is, determine the values for which the stated problem makes sense.
5. Determine the desired maximum or minimum value by the calculus techniques.

or maximum / minimum

Math for IT (Somkiat Wangsiripitak)

56

EXAMPLE 4 Finding Minimum Length

Two posts, one 12 feet high and the other 28 feet high, stand 30 feet apart. They are to be stayed by two wires, attached to a single stake, running from ground level to the top of each post. Where should the stake be placed to use the least amount of wire?

Solution Let W be the wire length to be minimized. you can write

$$W = y + z \quad \text{Primary equation}$$

In this problem, rather than solving for y in terms of z (or vice versa), you can solve for both y and z in terms of a third variable x . From the Pythagorean Theorem, you obtain

$$\begin{aligned} x^2 + 12^2 &= y^2 \\ (30 - x)^2 + 28^2 &= z^2 \end{aligned} \quad a^2 + b^2 = c^2$$

which implies that


$$\begin{aligned} y &= \sqrt{x^2 + 144} \\ z &= \sqrt{x^2 - 60x + 1684} \end{aligned}$$

So, W is given by

$$W = y + z = \sqrt{x^2 + 144} + \sqrt{x^2 - 60x + 1684}, \quad 0 \leq x \leq 30$$

Differentiating W with respect to x yields

$$\frac{dW}{dx} = \frac{x}{\sqrt{x^2 + 144}} + \frac{x - 30}{\sqrt{x^2 - 60x + 1684}}$$



The quantity to be minimized is length. From the diagram, you can see that x varies between 0 and 30.


By letting $dW/dx = 0$, you obtain

$$\begin{aligned} \frac{x}{\sqrt{x^2 + 144}} + \frac{x - 30}{\sqrt{x^2 - 60x + 1684}} &= 0 \\ x\sqrt{x^2 - 60x + 1684} &= (30 - x)\sqrt{x^2 + 144} \\ x^2(x^2 - 60x + 1684) &= (30 - x)^2(x^2 + 144) \\ x^4 - 60x^3 + 1684x^2 &= x^4 - 60x^3 + 1044x^2 - 8640x + 129,600 \\ 640x^2 + 8640x - 129,600 &= 0 \\ 320(x - 9)(2x + 45) &= 0 \\ x &= 9, -22.5 \end{aligned}$$

Because $x = -22.5$ is not in the domain and

$$W(0) \approx 53.04, \quad W(9) = 50, \quad \text{and} \quad W(30) \approx 60.31$$

you can conclude that the wire should be staked at 9 feet from the 12-foot pole.



Math for IT (Somkiat Wangsiripitak)

59

EXAMPLE 5 An Endpoint Maximum

Four feet of wire is to be used to form a square and a circle. How much of the wire should be used for the square and how much should be used for the circle to enclose the maximum total area?

$$\begin{aligned}
 A &= (\text{area of square}) + (\text{area of circle}) \\
 A &= x^2 + \pi r^2 \\
 \text{total length of wire} &= 4 \text{ feet;} \\
 4 &= (\text{perimeter of square}) + (\text{circumference of circle}) \\
 4 &= 4x + 2\pi r \\
 r &= 2(1-x)/\pi; \\
 A &= x^2 + \pi \left[\frac{2(1-x)}{\pi} \right]^2 \\
 &= x^2 + \frac{4(1-x)^2}{\pi} \\
 &= \frac{1}{\pi} [(\pi+4)x^2 - 8x + 4].
 \end{aligned}$$

domain $0 \leq x \leq 1$;


$$\frac{dA}{dx} = \frac{2(\pi+4)x - 8}{\pi} \stackrel{\text{set to } 0}{=}$$

$$\begin{aligned}
 2(\pi+4)x - 8 &= 0 \\
 (\pi+4)x &= \frac{8}{2} \\
 x &= \frac{4}{\pi+4}
 \end{aligned}$$

the only critical number in $(0, 1)$ is $x = 4/(\pi+4) \approx 0.56$. So, using
 $A(0) \approx 1.273$, $A(0.56) \approx 0.56$, and $A(1) = 1$
 you can conclude that the maximum area occurs when $x = 0$. That is, *all the wire is used for the circle.*

Math for IT (Somkiat Wangsiripitak)

60



Newton's Method

Approximating the zeros of a function

Newton's Method uses tangent lines to approximate the graph of the function near its x-intercepts.

To see how Newton's Method works, consider a function f that is continuous on the interval $[a, b]$ and differentiable on the interval (a, b) . If $f(a)$ and $f(b)$ differ in sign, then, by the Intermediate Value Theorem, f must have at least one zero in the interval (a, b) . Suppose you estimate this zero to occur at


$$x = x_1 \quad \text{First estimate}$$

Newton's Method is based on the assumption that the graph of f and the tangent line at $(x_1, f(x_1))$ both cross the x -axis at *about* the same point. Because you can easily calculate the x -intercept for this tangent line, you can use it as a second (and, usually, better) estimate of the zero of f . The tangent line passes through the point $(x_1, f(x_1))$ with a slope of $f'(x_1)$. In point-slope form, the equation of the tangent line is therefore

$$\begin{aligned}
 y - f(x_1) &= f'(x_1)(x - x_1) \\
 y &= f'(x_1)(x - x_1) + f(x_1).
 \end{aligned}$$

Letting $y = 0$ and solving for x produces

$$x = x_1 - \frac{f(x_1)}{f'(x_1)}.$$



Newton's Method

III

Objectives


- Approximate a zero of a function using Newton's Method.

So, from the initial estimate x_1 you obtain a new estimate

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} \quad \text{Second estimate}$$

You can improve on x_2 and calculate yet a third estimate

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)} \quad \text{Third estimate}$$



Repeated application of this process is called Newton's Method.

NEWTON'S METHOD FOR APPROXIMATING THE ZEROS OF A FUNCTION

Let $f(c) = 0$, where f is differentiable on an open interval containing c . Then, to approximate c , use the following steps.

- Make an initial estimate x_1 that is close to c . (A graph is helpful.)
- Determine a new approximation

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}.$$

- If $|x_n - x_{n+1}|$ is within the desired accuracy, let x_{n+1} serve as the final approximation. Otherwise, return to Step 2 and calculate a new approximation.

Each successive application of this procedure is called an **iteration**.

4

Math for IT (Somkiat Wangsiripitak)

62

EXAMPLE 1 Using Newton's Method


Calculate three iterations of Newton's Method to approximate a zero of $f(x) = x^2 - 2$. Use $x_1 = 1$ as the initial guess.

Solution Because $f(x) = x^2 - 2$, you have $f'(x) = 2x$, and the iterative process is given by the formula

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^2 - 2}{2x_n}.$$

The calculations for three iterations are shown in the table.

n	x_n	$f(x_n)$	$f'(x_n)$	$\frac{f(x_n)}{f'(x_n)}$	$x_n - \frac{f(x_n)}{f'(x_n)}$
1	1.000000	-1.000000	2.000000	-0.500000	1.500000
2	1.500000	0.250000	3.000000	0.083333	1.416667
3	1.416667	0.006945	2.833334	0.002451	1.414216
4	1.414216				



Of course, in this case you know that the two zeros of the function are $\pm\sqrt{2}$. To six decimal places, $\sqrt{2} = 1.414214$. So, after only three iterations of Newton's Method, you have obtained an approximation that is within 0.000002 of an actual root. The first iteration of this process is shown in Figure

65

L'Hôpital's Rule

THEOREM 8.4 L'HÔPITAL'S RULE

Let f and g be functions that are differentiable on an open interval (a, b) containing c , except possibly at c itself. Assume that $g'(x) \neq 0$ for all x in (a, b) , except possibly at c itself. If the limit of $f(x)/g(x)$ as x approaches c produces the indeterminate form $0/0$, then

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)}$$

provided the limit on the right exists (or is infinite). This result also applies if the limit of $f(x)/g(x)$ as x approaches c produces any one of the indeterminate forms ∞/∞ , $(-\infty)/\infty$, $\infty/(-\infty)$, or $(-\infty)/(-\infty)$.

L'Hôpital's Rule can also be applied to one-sided limits. For instance, if the limit of $f(x)/g(x)$ as x approaches c from the right produces the indeterminate form $0/0$, then

$$\lim_{x \rightarrow c^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c^+} \frac{f'(x)}{g'(x)}$$

provided the limit exists (or is infinite).

Another form of L'Hôpital's Rule states that if the limit of $f(x)/g(x)$ as x approaches ∞ (or $-\infty$) produces the indeterminate form $0/0$ or ∞/∞ , then

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)}$$

provided the limit on the right exists.

L'Hôpital's Rule

EXAMPLE Evaluate $\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x}$.

Solution Because direct substitution results in the indeterminate form $0/0$

$$\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x} \rightarrow \begin{cases} \lim_{x \rightarrow 0} (e^{2x} - 1) = 0 \\ \lim_{x \rightarrow 0} x = 0 \end{cases}$$

$$\begin{aligned} \frac{d}{dx}[e^x] &= e^x \\ \frac{d}{dx}[e^u] &= e^u \frac{du}{dx} \end{aligned}$$

you can apply L'Hôpital's Rule, as shown below.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x} &= \lim_{x \rightarrow 0} \frac{\frac{d}{dx}[e^{2x} - 1]}{\frac{d}{dx}[x]} \quad \text{Apply L'Hôpital's Rule.} \\ &= \lim_{x \rightarrow 0} \frac{2e^{2x}}{1} \quad \text{Differentiate numerator and denominator.} \\ &= 2 \quad \text{Evaluate the limit.} \end{aligned}$$

71

L'Hôpital's Rule

EXAMPLE Evaluate $\lim_{x \rightarrow -\infty} \frac{x^2}{e^{-x}} = \frac{\infty}{\infty} \rightarrow \text{indeterminate form}$

Solution Because direct substitution results in the indeterminate form ∞/∞ , you can apply L'Hôpital's Rule.

$$\begin{aligned} \lim_{x \rightarrow -\infty} \frac{x^2}{e^{-x}} &= \lim_{x \rightarrow -\infty} \frac{2x}{-e^{-x}} \\ &= \lim_{x \rightarrow -\infty} \frac{2}{(-1)(-1)e^{-x}} \\ &= \lim_{x \rightarrow -\infty} \frac{2}{e^{-x}} \\ &= \frac{2}{\infty} \rightarrow 0 \quad \text{2 หาร เลข 6666666666666666} \\ &= 0 \quad \text{= 6666666666666666} \end{aligned}$$

$$\begin{aligned} \frac{d}{dx}[e^x] &= e^x \\ \frac{d}{dx}[e^u] &= e^u \frac{du}{dx} \end{aligned}$$


Math for IT (Somkiat Wangsiripitak)

72

Homework

Find the area of the largest isosceles triangle that can be inscribed in a circle of radius 6 (see the figure).

Solve by writing the area as a function of h .



- Write down a step-by-step solution in a piece of A4 paper (or on an electronic paper).
- **DO NOT TYPE.**
- Submit your solution in MS Team.
- Do not forget to write your name and student ID.