

Vector Spaces #1

Instructor: Asst. Prof. Dr. Praphan Pavarangkoon

Office: Room No. 418-5, 4th Floor

Email: praphan@it.kmitl.ac.th

Office hours: Thursday at 13:00 – 16:00 (Before Midterm Examination)
or Tuesday at 9:00 – 12:00 (After Midterm Examination)
or as an advance appointment

Outline



- Vectors in the Plane
- Operations with Vectors in the Plane
- Vectors in R^n
- Length and Dot Product in R^n
- Angle Between Two Vectors
- The Cross Product of Two Vectors

Scalars and Vectors

ซึ่งขึ้นอยู่กับทิศทาง (ขนาด & ทิศทาง)

ขนาด Only

อัตราเร่ง

- A **scalar quantity** has only **magnitude**. of acceleration
- A **vector quantity** has both **magnitude** and **direction**.

Scalar Quantities

length, area, volume,
speed, mass, density,
pressure, temperature,
energy, entropy, work,
power



Scalar

24

Vector

$\begin{bmatrix} 2 & -8 & 7 \end{bmatrix}$

row

or

column

$\begin{bmatrix} 2 \\ -8 \\ 7 \end{bmatrix}$

Vector Quantities

displacement, velocity,
acceleration, momentum,
force, lift, drag, thrust,
weight

ความเร็ว
= A. เวกเตอร์ที่เพิ่มขึ้นใน
1 หน่วยเวลา



Matrix



$\begin{bmatrix} 6 & 4 & 24 \\ 1 & -9 & 8 \end{bmatrix}$

row(s) × column(s)

Vectors in the Plane

- A **vector in the plane** is represented geometrically by a **directed line segment** whose **initial point** is the origin and whose **terminal point** is the point (x_1, x_2) .
- This vector is represented by the same ordered pair used to represent its terminal point. That is,

$$\mathbf{x} = (x_1, x_2)$$



Vectors in the Plane (cont.)

- The coordinates x_1 and x_2 are called the **components** of the vector $\mathbf{x}$.
- Two vectors in the plane $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$ are **equal** if and only if $u_1 = v_1$ and $u_2 = v_2$.

vector ~~may~~ vectors
-> to carry

Vectors in the Plane (cont.)

Use a directed line segment to represent each vector in the plane.

(a) $\mathbf{u} = (2, 3)$ (b) $\mathbf{v} = (-1, 2)$

Operations with Vectors in the Plane


$\mathbb{R}^2$ 2 มิติ

Vector Addition

- To add two vectors in the plane, add their corresponding components. That is, the **sum** of **u** and **v** is the vector

$$\mathbf{u} + \mathbf{v} = (u_1, u_2) + (v_1, v_2) = (u_1 + v_1, u_2 + v_2).$$

- Geometrically, the sum of two vectors in the plane is represented as the diagonal of a parallelogram having **u** and **v** as its adjacent sides.



Adding Two Vectors in the Plane

Find the sum of the vectors.

(a) $\mathbf{u} = (1, 4)$, $\mathbf{v} = (2, -2)$ (b) $\mathbf{u} = (3, -2)$, $\mathbf{v} = (-3, 2)$ (c) $\mathbf{u} = (2, 1)$, $\mathbf{v} = (0, 0)$

(a) $\mathbf{u} + \mathbf{v} = (1, 4) + (2, -2) = (3, 2)$

(b) $\mathbf{u} + \mathbf{v} = (3, -2) + (-3, 2) = (0, 0)$

ผลรวมของ Vector เป็น $\mathbf{0}$ ทั้งหมด
เรียกว่า Vector $\mathbf{0}$


Operations with Vectors in the Plane (cont.)

Scalar Multiplication


- To multiply a vector $\mathbf{v}$ by a scalar c , multiply each of the components of $\mathbf{v}$ by c . That is,

$$c\mathbf{v} = c(v_1, v_2) = (cv_1, cv_2).$$

- The word *scalar* is used to mean a real number.
- In general, for a scalar c , the vector $c\mathbf{v}$ will be $|c|$ times as long as $\mathbf{v}$. If c is **positive**, then $c\mathbf{v}$ and $\mathbf{v}$ have the same direction, and if c is **negative**, then $c\mathbf{v}$ and $\mathbf{v}$ have opposite directions.



$c\mathbf{v}$ จะไปในทิศทาง
เดียวกันกับ $\mathbf{v}$



จะไปในทิศทาง
ตรงข้ามกับ $\mathbf{v}$

Operations with Vectors in the Plane (cont.)

Provided with $\mathbf{v} = (-2, 5)$ and $\mathbf{u} = (3, 4)$, find each vector.

(a) $\frac{1}{2}\mathbf{v}$ (b) $\mathbf{u} - \mathbf{v}$ (c) $\frac{1}{2}\mathbf{v} + \mathbf{u}$

$$(a) \left(\frac{1}{2}(-2), \frac{1}{2}(5) \right) = \left(-1, \frac{5}{2} \right)$$

Properties of Vector Addition and Scalar Multiplication in the Plane

เวกเตอร์ในระนาบ $\mathbb{R}^2$

Let $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ be vectors in the plane, and let c and d be scalars.

1. $\mathbf{u} + \mathbf{v}$ is a vector in the plane.
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. $c\mathbf{u}$ is a vector in the plane.
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1(\mathbf{u}) = \mathbf{u}$

Closure under addition คุณสมบัติปิดภายใต้การบวก

Commutative property of addition การสลับที่การบวก

Associative property of addition การเปลี่ยนหมู่ของการบวก

Additive identity property เวกเตอร์เอกลักษณ์การบวก

Additive inverse property Inverse การบวก

Closure under scalar multiplication คุณสมบัติปิดภายใต้การคูณด้วย scalar

Distributive property การแจกแจงด้วย scalar

Distributive property " $\forall$ scalar

Associative property of multiplication การเปลี่ยนหมู่ของการคูณ

Multiplicative identity property เวกเตอร์เอกลักษณ์การคูณ

Vectors in R^n

វ៉ិចទ័រ ក្នុង លំហ n វ៉ិចទ័រ

triple $R^3 \Rightarrow (x_1, x_2, x_3)$

quadruple $R^4 \Rightarrow (x_1, x_2, x_3, x_4)$

n -tuple $R^n \Rightarrow (x_1, x_2, \dots, x_n)$

- A vector in n -space is represented by an **ordered n -tuple**.
- The set of all n -tuples is called **n -space** and is denoted by R^n .

$R^1 = 1\text{-space} = \text{set of all real numbers}$

$R^2 = 2\text{-space} = \text{set of all ordered pairs of real numbers}$

$R^3 = 3\text{-space} = \text{set of all ordered triples of real numbers}$

$R^4 = 4\text{-space} = \text{set of all ordered quadruples of real numbers}$

$\vdots$

$R^n = n\text{-space} = \text{set of all ordered } n\text{-tuples of real numbers}$

- The practice of using an ordered pair to represent either a point or a vector in R^2 continues in R^n .
- That is, an n -tuple $(x_1, x_2, x_3, \dots, x_n)$ can be viewed as a **point** in R^n with the x_i 's as its coordinates or as a **vector**

$$\mathbf{x} = (x_1, x_2, x_3, \dots, x_n)$$

Vector in R^n

with the x_i 's as its components.

- As with vectors in the plane, two vectors in R^n are **equal** if and only if corresponding components are equal.

Vector Addition and Scalar Multiplication in $\mathbb{R}^n$


Let $\mathbf{u} = (u_1, u_2, u_3, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, v_3, \dots, v_n)$ be vectors in $\mathbb{R}^n$ and let c be a real number. Then the sum of $\mathbf{u}$ and $\mathbf{v}$ is defined as the vector

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, u_3 + v_3, \dots, u_n + v_n),$$

and the **scalar multiple** of $\mathbf{u}$ by c is defined as the vector

$$c\mathbf{u} = (cu_1, cu_2, cu_3, \dots, cu_n).$$

Vectors in R^n (cont.)



Vector Operations in R^3

Provided that $\mathbf{u} = (-1, 0, 1)$ and $\mathbf{v} = (2, -1, 5)$ in R^3 , find each vector.

(a) $\mathbf{u} + \mathbf{v}$ (b) $2\mathbf{u}$ (c) $\mathbf{v} - 2\mathbf{u}$

$(1, -1, 6)$ $(-2, 0, 2)$ $(4, -1, 3)$

Properties of Vector Addition and Scalar Multiplication in R^n

Let $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ be vectors in R^n , and let c and d be scalars.

- | | |
|--|--|
| 1. $\mathbf{u} + \mathbf{v}$ is a vector in R^n . | Closure under addition |
| 2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ | Commutative property of addition |
| 3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ | Associative property addition |
| 4. $\mathbf{u} + \mathbf{0} = \mathbf{u}$ | Additive identity property |
| 5. $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ | Additive inverse property |
| 6. $c\mathbf{u}$ is a vector in R^n . | Closure under scalar multiplication |
| 7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ | Distributive property |
| 8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ | Distributive property |
| 9. $c(d\mathbf{u}) = (cd)\mathbf{u}$ | Associative property of multiplication |
| 10. $1(\mathbf{u}) = \mathbf{u}$ | Multiplicative identity property |

Vector Operations in $\mathbb{R}^4$

Let $\mathbf{u} = (2, -1, 5, 0)$, $\mathbf{v} = (4, 3, 1, -1)$, and $\mathbf{w} = (-6, 2, 0, 3)$ be vectors in $\mathbb{R}^4$. Solve for $\mathbf{x}$.

(a) $\mathbf{x} = 2\mathbf{u} - (\mathbf{v} + 3\mathbf{w})$ (b) $3(\mathbf{x} + \mathbf{w}) = 2\mathbf{u} - \mathbf{v} + \mathbf{x}$

$$\begin{aligned} \text{(a)} \quad \mathbf{x} &= 2\mathbf{u} - (\mathbf{v} + 3\mathbf{w}) \\ &= 2\mathbf{u} - \mathbf{v} - 3\mathbf{w} \\ &= (4, -2, 10, 0) - (4, 3, 1, -1) - (-18, 6, 0, 9) \\ &= (4 - 4 + 18, -2 - 3 - 6, 10 - 1 - 0, 0 + 1 - 9) \\ &= (18, -11, 9, -8) \end{aligned}$$

Properties of Additive Identity and Additive Inverse

The zero vector $\mathbf{0}$ in R^n is called the **additive identity** in R^n . Similarly, the vector $-\mathbf{v}$ is called the **additive inverse** of $\mathbf{v}$. The theorem below summarizes several important properties of the additive identity and additive inverse in R^n .

Let $\mathbf{v}$ be a vector in R^n , and let c be a scalar. Then the following properties are true.

1. The additive identity is unique. That is, if $\mathbf{v} + \mathbf{u} = \mathbf{v}$, then $\mathbf{u} = \mathbf{0}$.
2. The additive inverse of $\mathbf{v}$ is unique. That is, if $\mathbf{v} + \mathbf{u} = \mathbf{0}$, then $\mathbf{u} = -\mathbf{v}$.
3. $0\mathbf{v} = \mathbf{0}$
4. $c\mathbf{0} = \mathbf{0}$
5. If $c\mathbf{v} = \mathbf{0}$, then $c = 0$ or $\mathbf{v} = \mathbf{0}$.
6. $-(-\mathbf{v}) = \mathbf{v}$

Length and Dot Product in R^n


- You will begin by reviewing the definition of the length of a vector in R^2 .
- If $\mathbf{v} = (v_1, v_2)$ is a vector in the plane, then the **length**, or **magnitude**, of denoted by $\|\mathbf{v}\|$, is defined as

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2}.$$

norm (ค่าบรรทัดฐาน)

- This definition corresponds to the usual notion of length in Euclidean geometry.
- That is, the vector is thought of as the hypotenuse of a right triangle whose sides have lengths of $|v_1|$ and $|v_2|$.
- Applying the Pythagorean Theorem produces

$$\|\mathbf{v}\|^2 = |v_1|^2 + |v_2|^2 = v_1^2 + v_2^2.$$



Length of a Vector in R^n

The **length**, or **magnitude**, of a vector $\mathbf{v} = (v_1, v_2, \dots, v_n)$ in R^n is given by

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}.$$

အားသာသော Vector $\rightarrow$
REMARK: The length of a vector is also called its **norm**. If $\|\mathbf{v}\| = 1$, then the vector $\mathbf{v}$ is called a **unit vector**. *vector ၁ ခုသာ*

This definition shows that the length of a **vector cannot be negative**. That is, $\|\mathbf{v}\| \geq 0$. Moreover, $\|\mathbf{v}\| = 0$ if and only if $\mathbf{v}$ is the zero vector $\mathbf{0}$.

The Length of a Vector in R^n

$$60 + 16 + 0 + 0 + 4 = 80$$

(a) In R^5 , the length of $\mathbf{v} = (0, -2, 1, 4, -2)$ is

(b) In R^3 , the length of $\mathbf{v} = (2/\sqrt{17}, -2/\sqrt{17}, 3/\sqrt{17})$ is

Length of a Scalar Multiple

Let $\mathbf{v}$ be a vector in R^n and let c be a scalar. Then

$$\|c\mathbf{v}\| = |c| \|\mathbf{v}\|,$$

where $|c|$ is the absolute value of c .

Unit Vector in the Direction of $\mathbf{v}$

If $\mathbf{v}$ is a nonzero vector in R^n , then the vector

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$$

has length 1 and has the same direction as $\mathbf{v}$. This vector $\mathbf{u}$ is called the **unit vector in the direction of $\mathbf{v}$** .

Finding a Unit Vector

Find the unit vector in the direction of $\mathbf{v} = (3, -1, 2)$, and verify that this vector has length 1.

$$\frac{\mathbf{v}}{\|\mathbf{v}\|}$$

$$\sqrt{3^2 + (-1)^2 + 2^2}$$

Distance Between Two Vectors in R^n

distance between two vectors $\vec{u}$ and $\vec{v}$ in R^n

The **distance between two vectors $\mathbf{u}$ and $\mathbf{v}$ in R^n** is

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|.$$

You can easily verify the three properties of distance listed below.

1. $d(\mathbf{u}, \mathbf{v}) \geq 0$
2. $d(\mathbf{u}, \mathbf{v}) = 0$ if and only if $\mathbf{u} = \mathbf{v}$.
3. $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$

Finding the Distance Between Two Vectors

The distance between $\mathbf{u} = (0, 2, 2)$ and $\mathbf{v} = (2, 0, 1)$ is

$$\|\mathbf{u} - \mathbf{v}\|$$

Scalar Product

Dot Product in R^n

The **dot product** of $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is the *scalar* quantity

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n.$$

REMARK: Notice that the dot product of two vectors is a scalar, not another vector.

Dot $\Rightarrow$ Scalar

Cross $\Rightarrow$ Vector

Finding the Dot Product of Two Vectors

The dot product of $\mathbf{u} = (1, 2, 0, -3)$ and $\mathbf{v} = (3, -2, 4, 2)$ is

$$(1)(3) + (2)(-2) + (0)(4) + (-3)(2)$$

Properties of the Dot Product

If $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are vectors in R^n and c is a scalar, then the following properties are true.

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$

2. $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$

3. $c(\mathbf{u} \cdot \mathbf{v}) = (c\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (c\mathbf{v})$

4. $\mathbf{v} \cdot \mathbf{v} = \|\mathbf{v}\|^2$

5. $\mathbf{v} \cdot \mathbf{v} \geq 0$, and $\mathbf{v} \cdot \mathbf{v} = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

Finding Dot Products

Given $\mathbf{u} = (2, -2)$, $\mathbf{v} = (5, 8)$, and $\mathbf{w} = (-4, 3)$, find

- (a) $\mathbf{u} \cdot \mathbf{v}$. (b) $(\mathbf{u} \cdot \mathbf{v})\mathbf{w}$. (c) $\mathbf{u} \cdot (2\mathbf{v})$. (d) $\|\mathbf{w}\|^2$. (e) $\mathbf{u} \cdot (\mathbf{v} - 2\mathbf{w})$.

$$(a) \quad (2)(5) + (-2)(8) = -6$$

$$(b) \quad -6(\mathbf{w}) = -6(-4, 3) = (24, -18)$$

$$(c) \quad 2(\mathbf{u} \cdot \mathbf{v}) = 2(-6) = -12$$

$$(d) \quad \mathbf{w} \cdot \mathbf{w} = (-4)(-4) + (3)(3) = 25$$

Using Properties of the Dot Product

Provided with two vectors $\mathbf{u}$ and $\mathbf{v}$ in $\mathbb{R}^n$ such that $\mathbf{u} \cdot \mathbf{u} = 39$, $\mathbf{u} \cdot \mathbf{v} = -3$, and $\mathbf{v} \cdot \mathbf{v} = 79$, evaluate $(\mathbf{u} + 2\mathbf{v}) \cdot (3\mathbf{u} + \mathbf{v})$.

$$(\mathbf{u} + 2\mathbf{v}) \cdot (3\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot 3\mathbf{u} + \mathbf{u} \cdot \mathbf{v} + (2\mathbf{v}) \cdot 3\mathbf{u} + 2\mathbf{v} \cdot \mathbf{v}$$

The Angle Between Two Vectors in R^n

Cauchy - Schwarz
Inequality

The **angle** θ between two nonzero vectors in R^n is given by

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}, \quad 0 \leq \theta \leq \pi.$$

REMARK: The angle between the zero vector and another vector is not defined.

Finding the Angle Between Two Vectors

$\mathbb{R}^4$


The angle between $\mathbf{u} = (-4, 0, 2, -2)$ and $\mathbf{v} = (2, 0, -1, 1)$ is

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{-12}{\sqrt{24} \sqrt{6}} = \frac{-12}{\sqrt{144}} = -1$$

$$\theta = \cos^{-1} \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} \right)$$

↳ arc cosine

↳ inverse cosine function



moving degrees

radius

The Angle Between Two Vectors in R^n (cont.)

Note that because $\|\mathbf{u}\|$ and $\|\mathbf{v}\|$ are always positive, $\mathbf{u} \cdot \mathbf{v}$ and $\cos \theta$ will always have the same sign. Moreover, because the cosine is positive in the first quadrant and negative in the second quadrant, the sign of the dot product of two vectors can be used to determine whether the angle between them is acute or obtuse, as shown in Figure 5.6.




Figure 5.6

Orthogonal Vectors

Two vectors $\mathbf{u}$ and $\mathbf{v}$ in R^n are **orthogonal** if

$$\mathbf{u} \cdot \mathbf{v} = 0.$$

REMARK: Even though the angle between the zero vector and another vector is not defined, it is convenient to extend the definition of orthogonality to include the zero vector. In other words, the vector $\mathbf{0}$ is said to be orthogonal to every vector.

Orthogonal Vectors in R^n

- (a) The vectors $\mathbf{u} = (1, 0, 0)$ and $\mathbf{v} = (0, 1, 0)$ are orthogonal because
- (b) The vectors $\mathbf{u} = (3, 2, -1, 4)$ and $\mathbf{v} = (1, -1, 1, 0)$ are orthogonal because

Finding Orthogonal Vectors

Determine all vectors in R^2 that are orthogonal to $\mathbf{u} = (4, 2)$.

The Cross Product of Two Vectors in Space

Vector Product

Many problems in linear algebra involve finding a vector orthogonal to each vector in a set. Here you will look at a vector product that yields a vector in R^3 orthogonal to two vectors. This vector product is called the **cross product**, and it is most conveniently defined and calculated with vectors written in standard unit vector form.

$$\mathbf{v} = (v_1, v_2, v_3) = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$$

ឥឡូវនេះ យើងនឹងសិក្សាពីផលគុណខ្លោង

Let $\mathbf{u} = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$ and $\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$ be vectors in R^3 . The **cross product** of $\mathbf{u}$ and $\mathbf{v}$ is the vector

$$\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2)\mathbf{i} - (u_1v_3 - u_3v_1)\mathbf{j} + (u_1v_2 - u_2v_1)\mathbf{k}.$$

REMARK: The cross product is defined only for vectors in R^3 . The cross product of two vectors in R^2 or of vectors in R^n , $n > 3$, is not defined here.

A convenient way to remember the formula for the cross product $\mathbf{u} \times \mathbf{v}$ is to use the determinant form shown below.

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

← Components of $\mathbf{u}$

← Components of $\mathbf{v}$

Finding the Cross Product of Two Vectors

$$\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$$

$$\mathbf{v} \times \mathbf{v} = \mathbf{0} \quad \text{matrix 2 row, missing det (0)}$$

Provided that $\mathbf{u} = \mathbf{i} - 2\mathbf{j} + \mathbf{k}$ and $\mathbf{v} = 3\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, find

(a) $\mathbf{u} \times \mathbf{v}$.

(b) $\mathbf{v} \times \mathbf{u}$.

(c) $\mathbf{v} \times \mathbf{v}$.

$$(a) \quad \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

Algebraic Properties of the Cross Product

If $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are vectors in R^3 and c is a scalar, then the following properties are true.

1. $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$
2. $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w})$
3. $c(\mathbf{u} \times \mathbf{v}) = c\mathbf{u} \times \mathbf{v} = \mathbf{u} \times c\mathbf{v}$
4. $\mathbf{u} \times \mathbf{0} = \mathbf{0} \times \mathbf{u} = \mathbf{0}$
5. $\mathbf{u} \times \mathbf{u} = \mathbf{0}$
6. $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$

Geometric Properties of the Cross Product


If $\mathbf{u}$ and $\mathbf{v}$ are nonzero vectors in R^3 , then the following properties are true.

1. $\mathbf{u} \times \mathbf{v}$ is orthogonal to both $\mathbf{u}$ and $\mathbf{v}$.
2. The angle θ between $\mathbf{u}$ and $\mathbf{v}$ is given by

$$\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\| \|\mathbf{v}\| \sin \theta.$$

3. $\mathbf{u}$ and $\mathbf{v}$ are parallel if and only if $\mathbf{u} \times \mathbf{v} = \mathbf{0}$.
4. The parallelogram having $\mathbf{u}$ and $\mathbf{v}$ as adjacent sides has an area of $\|\mathbf{u} \times \mathbf{v}\|$.

$$\text{Area} = \overbrace{\|\mathbf{u}\|}^{\text{Base}} \overbrace{\|\mathbf{v}\| \sin \theta}^{\text{Height}} = \|\mathbf{u} \times \mathbf{v}\|.$$



Finding a Vector Orthogonal to Two Given Vectors

Find a unit vector orthogonal to both

$$\mathbf{u} = \mathbf{i} - 4\mathbf{j} + \mathbf{k}$$

and

$$\mathbf{v} = 2\mathbf{i} + 3\mathbf{j}.$$

Finding the Area of a Parallelogram

រកផ្ទៃក្រឡា $\square$ ជុំវិញវ៉ិចទ័រ


Find the area of the parallelogram that has

$$\mathbf{u} = -3\mathbf{i} + 4\mathbf{j} + \mathbf{k}$$

and

$$\mathbf{v} = -2\mathbf{j} + 6\mathbf{k}$$

as adjacent sides, as shown in Figure 5.31.



The area of the parallelogram is given by
 $\|\mathbf{u} \times \mathbf{v}\| = \sqrt{1036}.$

Figure 5.31



Q & A