

Mathematics for Information Technology

Somkiat Wangsiripitak

somkiat.wa@kmitl.ac.th

Room.518 or Room.506 (MIV Lab)


INTEGRATION (II)

The Fundamental Theorem of Calculus ^{III}

Objectives

- Evaluate a definite integral using the Fundamental Theorem of Calculus.
- Understand and use the Mean Value Theorem for Integrals.
- Find the average value of a function over a closed interval.
- Understand and use the Second Fundamental Theorem of Calculus.
- Understand and use the Net Change Theorem.

The Fundamental Theorem of Calculus



(a) Differentiation

(b) Definite integration

Differentiation and definite integration have an "inverse" relationship.

THEOREM 4.9 THE FUNDAMENTAL THEOREM OF CALCULUS

If a function f is continuous on the closed interval $[a, b]$ and F is an antiderivative of f on the interval $[a, b]$, then

$$\int_a^b f(x) dx = F(b) - F(a).$$

GUIDELINES FOR USING THE FUNDAMENTAL THEOREM OF CALCULUS

1. Provided you can find an antiderivative of f , you now have a way to evaluate a definite integral without having to use the limit of a sum.
2. When applying the Fundamental Theorem of Calculus, the following notation is convenient.

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a)$$

For instance, to evaluate $\int_1^3 x^3 dx$, you can write

$$\int_1^3 x^3 dx = \frac{x^4}{4} \Big|_1^3 = \frac{3^4}{4} - \frac{1^4}{4} = \frac{81}{4} - \frac{1}{4} = 20.$$

3. It is not necessary to include a constant of integration C in the antiderivative because

$$\begin{aligned} \int_a^b f(x) dx &= [F(x) + C]_a^b \\ &= [F(b) + C] - [F(a) + C] \\ &= F(b) - F(a). \end{aligned}$$

EXAMPLE 1 Evaluating a Definite Integral

Evaluate each definite integral.

a. $\int_1^2 (x^2 - 3) dx$ b. $\int_1^4 3\sqrt{x} dx$ c. $\int_0^{\pi/4} \sec^2 x dx$

Solution

a. $\int_1^2 (x^2 - 3) dx = \left[\frac{x^3}{3} - 3x \right]_1^2 = \frac{2^3}{3} - 3(2) - \left(\frac{1^3}{3} - 3(1) \right) = \frac{8}{3} - 6 - \frac{1}{3} + 3 = \frac{8-18-1+9}{3} = -\frac{2}{3}$

b. $\int_1^4 3\sqrt{x} dx = 3 \int_1^4 x^{1/2} dx = 3 \left[\frac{2}{3} x^{3/2} \right]_1^4 = 2 \left[\sqrt{x} \right]_1^4 = 2(\sqrt{4} - \sqrt{1}) = 2(2 - 1) = 2$

c. $\int_0^{\pi/4} \sec^2 x dx = \left[\tan x \right]_0^{\pi/4} = \tan\left(\frac{\pi}{4}\right) - \tan(0) = 1 - 0 = 1$


EXAMPLE 2 A Definite Integral Involving Absolute Value

Evaluate $\int_0^2 |2x - 1| dx$.
 $2x - 1 > 0 \Rightarrow x > \frac{1}{2}$
 $2x - 1 < 0 \Rightarrow x < \frac{1}{2}$

Solution

$\int_0^2 |2x - 1| dx = \int_{\frac{1}{2}}^2 (2x - 1) dx + \int_0^{\frac{1}{2}} -(2x - 1) dx$
 $= \left[x^2 - x \right]_{\frac{1}{2}}^2 + \left[-x^2 + x \right]_0^{\frac{1}{2}}$


$= (4 - 2) - \left(\frac{1}{4} - \frac{1}{2} \right) + \left(-\frac{1}{4} + \frac{1}{2} - 0 + 0 \right) = 2 - \frac{1}{4} + \frac{1}{2} - \frac{1}{4} + \frac{1}{2} = 2$



EXAMPLE 3 Using the Fundamental Theorem to Find Area

Find the area of the region bounded by the graph of $y = 2x^2 - 3x + 2$, the x -axis, and the vertical lines $x = 0$ and $x = 2$, as shown in Figure


$$\begin{aligned}
 & \int_0^2 (2x^2 - 3x + 2) dx \\
 &= \left[\frac{2x^3}{3} - \frac{3x^2}{2} + 2x \right]_0^2 \\
 &= \frac{16}{3} - \frac{12}{2} + 4 - 0 \\
 &= \frac{32 - 36 + 24}{6} = \frac{20}{6} = \frac{10}{3}
 \end{aligned}$$



Math for IT (Somkiat Wangsiripitak)

The Mean Value Theorem for Integrals

Mean Value Theorem for Integrals states that somewhere “between” the inscribed and circumscribed rectangles there is a rectangle whose area is precisely equal to the area of the region under the curve, as shown in Figure



Mean value rectangle:

$$f(c)(b - a) = \int_a^b f(x) dx$$

THEOREM 4.10 MEAN VALUE THEOREM FOR INTEGRALS

If f is continuous on the closed interval $[a, b]$, then there exists a number c in the closed interval $[a, b]$ such that

$$\int_a^b f(x) dx = f(c)(b - a).$$

NOTE Notice that Theorem 4.10 does not specify how to determine c . It merely guarantees the existence of at least one number c in the interval.


Average Value of a Function

The value of $f(c)$ given in the Mean Value Theorem for Integrals is called the **average value** of f on the interval $[a, b]$.

DEFINITION OF THE AVERAGE VALUE OF A FUNCTION ON AN INTERVAL

If f is integrable on the closed interval $[a, b]$, then the **average value** of f on the interval is

$$\frac{1}{b - a} \int_a^b f(x) dx.$$



NOTE Notice in Figure that the area of the region under the graph of f is equal to the area of the rectangle whose height is the average value.


Math for IT (Somkiat Wangsiripitak)

EXAMPLE 4 Finding the Average Value of a Function

Find the average value of $f(x) = 3x^2 - 2x$ on the interval $[1, 4]$.

Solution The average value is given by


$$\begin{aligned}
 & \frac{1}{b - a} \int_a^b f(x) dx \\
 &= \frac{1}{4 - 1} \int_1^4 (3x^2 - 2x) dx \\
 &= \frac{1}{3} \left[x^3 - x^2 \right]_1^4 \\
 &= \frac{1}{3} (64 - 16 - 1 + 1) \\
 &= \frac{1}{3} (48) = 16
 \end{aligned}$$




Math for IT (Somkiat Wangsiripitak)

The Second Fundamental Theorem of Calculus

The Definite Integral as a Number



The Definite Integral as a Function of x



EXAMPLE 6 The Definite Integral as a Function

Evaluate the function


$$F(x) = \int_0^x \cos t \, dt$$

at $x = 0, \pi/6, \pi/4, \pi/3$, and $\pi/2$.

Solution You could evaluate five different definite integrals, one for each of the given upper limits. However, it is much simpler to fix x (as a constant) temporarily to obtain

$$\int_0^x \cos t \, dt = \sin t \Big|_0^x = \sin x - \sin 0 = \sin x.$$

Now, using $F(x) = \sin x$, you can obtain the results shown in Figure



$F(x) = \int_0^x \cos t \, dt$ is the area under the curve $f(t) = \cos t$ from 0 to x .

an accumulation function
accumulating the area

THEOREM 4.11 THE SECOND FUNDAMENTAL THEOREM OF CALCULUS

If f is continuous on an open interval I containing a , then, for every x in the interval,

$$\frac{d}{dx} \left[\int_a^x f(t) \, dt \right] = f(x).$$

It can be used to avoid the long process of integration followed by differentiation

EXAMPLE 8 Using the Second Fundamental Theorem of Calculus

Find the derivative of $F(x) = \int_{\pi/2}^{x^3} \cos t \, dt$.

Solution Using $u = x^3$, you can apply the Second Fundamental Theorem of Calculus with the Chain Rule as shown.

$$\begin{aligned} F'(x) &= \frac{dF}{du} \frac{du}{dx} \\ &= \frac{d}{du} [F(x)] \frac{du}{dx} \\ &= \frac{d}{du} \left[\int_{\pi/2}^{x^3} \cos t \, dt \right] \frac{du}{dx} \\ &= \frac{d}{du} \left[\int_{\pi/2}^u \cos t \, dt \right] \frac{du}{dx} \\ &= (\cos u)(3x^2) \\ &= (\cos x^3)(3x^2) \end{aligned}$$

Chain Rule

Definition of $\frac{dF}{du}$

Substitute $\int_{\pi/2}^{x^3} \cos t \, dt$ for $F(x)$.

Substitute u for x^3 .

Apply Second Fundamental Theorem of Calculus.

Rewrite as function of x .

$$\frac{d}{dx} \left[\int_a^x f(t) \, dt \right] = f(x)$$

Alternative

Because the integrand in Example 8 is easily integrated, you can verify the derivative as follows.

$$F(x) = \int_{\pi/2}^{x^3} \cos t \, dt = \sin t \Big|_{\pi/2}^{x^3} = \sin x^3 - \sin \frac{\pi}{2} = (\sin x^3) - 1$$

In this form, you can apply the Power Rule to verify that the derivative is the same as that obtained in Example 8.

$$F'(x) = (\cos x^3)(3x^2)$$

Net Change Theorem

THEOREM 4.12 THE NET CHANGE THEOREM

The definite integral of the rate of change of a quantity $F'(x)$ gives the total change, or net change, in that quantity on the interval $[a, b]$.

$$\int_a^b F'(x) dx = F(b) - F(a) \quad \text{Net change of } F$$

EXAMPLE 9 Using the Net Change Theorem

A chemical flows into a storage tank at a rate of $180 + 3t$ liters per minute, where $0 \leq t \leq 60$. Find the amount of the chemical that flows into the tank during the first 20 minutes.

Solution Let $c(t)$ be the amount of the chemical in the tank at time t . Then $c'(t)$ represents the rate at which the chemical flows into the tank at time t . During the first 20 minutes, the amount that flows into the tank is

$$\begin{aligned} \int_0^{20} c'(t) dt &= \int_0^{20} (180 + 3t) dt \\ &= \left[180t + \frac{3}{2}t^2 \right]_0^{20} \\ &= 3600 + 600 = 4200. \end{aligned}$$

So, the amount that flows into the tank during the first 20 minutes is 4200 liters.


EXAMPLE 10 Solving a Particle Motion Problem

A particle is moving along a line so that its velocity is $v(t) = t^3 - 10t^2 + 29t - 20$ feet per second at time t .

- What is the displacement of the particle on the time interval $1 \leq t \leq 5$?
- What is the total distance traveled by the particle on the time interval $1 \leq t \leq 5$?

Solution

$$\begin{aligned} \text{a. } \int_1^5 v(t) dt &= \int_1^5 (t^3 - 10t^2 + 29t - 20) dt \\ &= \int_1^5 \left(\frac{t^4}{4} - \frac{10t^3}{3} + \frac{29t^2}{2} - 20t \right) dt \\ &= \left[\frac{t^4}{4} - \frac{10t^3}{3} + \frac{29t^2}{2} - 20t \right]_1^5 \\ &= \left(\frac{625}{4} - \frac{1250}{3} + \frac{725}{2} - 100 \right) - \left(\frac{1}{4} - \frac{10}{3} + \frac{29}{2} - 20 \right) \\ &= \frac{36}{3} \end{aligned}$$




EXAMPLE 10 Solving a Particle Motion Problem

A particle is moving along a line so that its velocity is $v(t) = t^3 - 10t^2 + 29t - 20$ feet per second at time t .

- What is the displacement of the particle on the time interval $1 \leq t \leq 5$?
- What is the total distance traveled by the particle on the time interval $1 \leq t \leq 5$?

Solution

$$\text{b. } \int_1^5 |v(t)| dt \rightarrow \begin{aligned} &t^3 - 10t^2 + 29t - 20 \\ &= (t-1)(t-4)(t-5) \\ &v(t) > 0 \text{ on } [1, 4] \\ &v(t) \leq 0 \text{ on } [4, 5] \end{aligned}$$



BREAK

Integration by Substitution

IV

Objectives

- Use pattern recognition to find an indefinite integral.
- Use a change of variables to find an indefinite integral.
- Use the General Power Rule for Integration to find an indefinite integral.
- Use a change of variables to evaluate a definite integral.

Pattern Recognition

In this section you will study techniques for integrating composite functions. The discussion is split into two parts—pattern recognition and change of variables. Both techniques involve a u -substitution. With pattern recognition you perform the substitution mentally, and with change of variables you write the substitution steps.

THEOREM 4.13 ANTIDIFFERENTIATION OF A COMPOSITE FUNCTION


Let g be a function whose range is an interval I , and let f be a function that is continuous on I . If g is differentiable on its domain and F is an antiderivative of f on I , then

$$\int f(g(x))g'(x) dx = F(g(x)) + C.$$

Letting $u = g(x)$ gives $du = g'(x) dx$ and

$$\int f(u) du = F(u) + C.$$

NOTE The statement of Theorem 4.13 doesn't tell how to distinguish between $f(g(x))$ and $g'(x)$ in the integrand. As you become more experienced at integration, your skill in doing this will increase. Of course, part of the key is familiarity with derivatives.



EXAMPLE 1 Recognizing the $f(g(x))g'(x)$ Pattern

Find $\int (x^2 + 1)^2(2x) dx$.

Solution Letting $g(x) = x^2 + 1$, you obtain

$$g'(x) = 2x$$

and

$$f(g(x)) = f(x^2 + 1) = (x^2 + 1)^2.$$

From this, you can recognize that the integrand follows the $f(g(x))g'(x)$ pattern. Using the Power Rule for Integration and Theorem 4.13, you can write

$$\int \overbrace{(x^2 + 1)^2(2x)}^{f(g(x)) \quad g'(x)} dx = \frac{1}{3}(x^2 + 1)^3 + C.$$

Try using the Chain Rule to check that the derivative of $\frac{1}{3}(x^2 + 1)^3 + C$ is the integrand of the original integral.

Constant Multiple Rule

$$\int kf(x) dx = k \int f(x) dx.$$

Many integrands contain the essential part (the variable part) of $g'(x)$ but are missing a constant multiple. In such cases, you can multiply and divide by the necessary constant multiple, as shown in Example 3.

EXAMPLE 3 Multiplying and Dividing by a Constant

Find $\int x(x^2 + 1)^2 dx$.

Solution This is similar to the integral given in Example 1, except that the integrand is missing a factor of 2. Recognizing that $2x$ is the derivative of $x^2 + 1$, you can let $g(x) = x^2 + 1$ and supply the $2x$ as follows.

$$\begin{aligned} \int x(x^2 + 1)^2 dx &= \int (x^2 + 1)^2 \left(\frac{1}{2}\right)(2x) dx && \text{Multiply and divide by 2.} \\ &= \frac{1}{2} \int \overbrace{(x^2 + 1)^2}^{f(g(x))} \overbrace{(2x)}^{g'(x)} dx && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left[\frac{(x^2 + 1)^3}{3} \right] + C && \text{Integrate.} \\ &= \frac{1}{6} (x^2 + 1)^3 + C && \text{Simplify.} \end{aligned}$$

EXAMPLE 1

Find $\int (x^2 + 1)^2(2x) dx$.

EXAMPLE 3 Multiplying and Dividing by a Constant

Find $\int x(x^2 + 1)^2 dx$.

Solution

In practice, most people would not write as many steps as are shown in Example 3. For instance, you could evaluate the integral by simply writing

$$\begin{aligned} \int x(x^2 + 1)^2 dx &= \frac{1}{2} \int (x^2 + 1)^2 2x dx \\ &= \frac{1}{2} \left[\frac{(x^2 + 1)^3}{3} \right] + C \\ &= \frac{1}{6} (x^2 + 1)^3 + C. \end{aligned}$$

Change of Variables

With a formal **change of variables**, you completely rewrite the integral in terms of u and du (or any other convenient variable). Although this procedure can involve more written steps than the pattern recognition illustrated in Examples 1 to 3, it is useful for complicated integrands. The change of variables technique uses the Leibniz notation for the differential. That is, if $u = g(x)$, then $du = g'(x) dx$, and the integral in Theorem 4.13 takes the form

$$\int f(g(x))g'(x) dx = \int f(u) du = F(u) + C.$$

EXAMPLE 4 Change of Variables

Find $\int \sqrt{2x-1} dx$.

Solution First, let u be the inner function, $u = 2x - 1$. Then calculate the differential du to be $du = 2 dx$. Now, using $\sqrt{2x-1} = \sqrt{u}$ and $dx = du/2$, substitute to obtain

$$\begin{aligned} \int \sqrt{2x-1} dx &= \int \sqrt{u} \left(\frac{du}{2} \right) && \text{Integral in terms of } u \\ &= \frac{1}{2} \int u^{1/2} du && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left(\frac{u^{3/2}}{3/2} \right) + C && \text{Antiderivative in terms of } u \\ &= \frac{1}{3} u^{3/2} + C && \text{Simplify.} \\ &= \frac{1}{3} (2x-1)^{3/2} + C. && \text{Antiderivative in terms of } x \end{aligned}$$

EXAMPLE 5 Change of Variables

Find $\int x\sqrt{2x-1} dx$. $u = 2x-1$
 $du = 2 dx$ $dx = \frac{du}{2}$

$$\begin{aligned} \int x\sqrt{2x-1} dx &= \int \frac{u+1}{2} \sqrt{u} \frac{du}{2} \\ &= \int \frac{(u+1)\sqrt{u}}{2} du \\ &= \int \frac{u\sqrt{u} + \sqrt{u}}{2} du \\ &= \frac{1}{2} \int u\sqrt{u} + \sqrt{u} du \\ &= \frac{1}{2} \left(\frac{u^{5/2}}{5/2} + \frac{u^{3/2}}{3/2} \right) \\ &= \frac{1}{2} \left(\frac{2u^{5/2}}{5} + \frac{2u^{3/2}}{3} \right) \\ &= \frac{1}{2} \left(\frac{2(2x-1)^{5/2}}{5} + \frac{2(2x-1)^{3/2}}{3} \right) \\ &= \frac{(2x-1)^{5/2}}{5} + \frac{(2x-1)^{3/2}}{3} \end{aligned}$$

GUIDELINES FOR MAKING A CHANGE OF VARIABLES

1. Choose a substitution $u = g(x)$. Usually, it is best to choose the *inner* part of a composite function, such as a quantity raised to a power.
2. Compute $du = g'(x) dx$.
3. Rewrite the integral in terms of the variable u .
4. Find the resulting integral in terms of u .
5. Replace u by $g(x)$ to obtain an antiderivative in terms of x .
6. Check your answer by differentiating.

The General Power Rule for Integration

THEOREM 4.14 THE GENERAL POWER RULE FOR INTEGRATION

If g is a differentiable function of x , then

$$\int [g(x)]^n g'(x) dx = \frac{[g(x)]^{n+1}}{n+1} + C, \quad n \neq -1.$$

Equivalently, if $u = g(x)$, then

$$\int u^n du = \frac{u^{n+1}}{n+1} + C, \quad n \neq -1.$$

EXAMPLE 7 Substitution and the General Power Rule

a. $\int 3(3x - 1)^4 dx = \int \overbrace{(3x - 1)^4}^{u^4} \overbrace{3}^{du} dx = \frac{(3x - 1)^5}{5} + C$

b. $\int (2x + 1)(x^2 + x) dx = \int \overbrace{(x^2 + x)}^{u^1} \overbrace{(2x + 1)}^{du} dx = \frac{(x^2 + x)^2}{2} + C$

c. $\int 3x^2 \sqrt{x^3 - 2} dx = \int \overbrace{\sqrt{x^3 - 2}}^{u^{\frac{1}{2}}} \overbrace{(3x^2)}^{du} dx = \frac{2(x^3 - 2)^{\frac{3}{2}}}{\frac{3}{2}} = \frac{(2x^3 - 4)\sqrt{x^3 - 2}}{3} + C$

d. $\int \frac{-4x}{(1 - 2x^2)^2} dx =$

e. $\int \cos^2 x \sin x dx =$

Some integrals whose integrands involve quantities raised to powers cannot be found by the General Power Rule. Consider the two integrals

$$\int x(x^2 + 1)^2 dx \quad \text{and} \quad \int (x^2 + 1)^2 dx.$$

The substitution $u = x^2 + 1$ works in the first integral but not in the second. In the second, the substitution fails because the integrand lacks the factor x needed for du . Fortunately, for this particular integral, you can expand the integrand as $(x^2 + 1)^2 = x^4 + 2x^2 + 1$ and use the (simple) Power Rule to integrate each term.

Integration by Substitution

EXAMPLE

$$\begin{aligned} \int (2x + 3)^7 dx &= \frac{1}{2} \int 2(2x + 3)^7 dx \\ &= \frac{(2x + 3)^8}{8} + C \end{aligned}$$

Integration by Substitution

EXAMPLE

$$\int x(x^2 + 1)^4 dx \quad ?$$

$$\begin{aligned} & \frac{1}{2} \int (x^2 + 1)^4 (2x) dx \\ &= \frac{(x^2 + 1)^5}{10} \end{aligned}$$

Integration by Substitution

EXAMPLE

$$\begin{aligned} & \int 3x^2 \sqrt{x^3 + 5} dx \quad ? \\ &= \frac{(x^3 + 5)^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2x^3 + 10 \sqrt{x^3 + 5}}{3} + C \end{aligned}$$

Integration by Substitution

EXAMPLE

$$\frac{d}{dx}[\ln u] = \frac{1}{u} \frac{du}{dx} = \frac{u'}{u}, \quad u > 0 \quad \left| \quad \int \frac{1}{u} du = \ln|u| + C \right|$$

$$\begin{aligned} \int \frac{x}{3x^2 + 4} dx \quad ? &= \int x(3x^2 + 4)^{-1} dx \\ &= \frac{1}{6} \int \frac{6x}{3x^2 + 4} dx \\ &= \frac{1}{6} \ln|3x^2 + 4| + C \end{aligned}$$

Integration by Substitution

EXAMPLE

$$\begin{aligned} & \int \sin(3x + 2) dx \quad ? \quad \begin{array}{l} u = 3x + 2 \\ du = 3 dx \end{array} \\ & \frac{1}{3} \int \sin(u) du \\ &= \frac{-\cos(3x + 2)}{3} + C \end{aligned}$$

Pattern
 $\frac{1}{3} \int 3 \sin(3x + 2) dx$

Integration by Substitution

EXAMPLE $\int \sin x \cos x \, dx$?

$u = \sin x$
 $du = \cos x \, dx$ (2003 sin)

$\int u \, du = \frac{u^2}{2} = \frac{\sin^2(x)}{2} + C$

$= -\frac{1}{9} \cos(2x) + C$

Math for IT (Somkiat Wangsiripitak)

67

Change of Variables for Definite Integrals

When using u -substitution with a definite integral, it is often convenient to determine the limits of integration for the variable u rather than to convert the antiderivative back to the variable x and evaluate at the original limits. This change of variables is stated explicitly in the next theorem.

THEOREM 4.15 CHANGE OF VARIABLES FOR DEFINITE INTEGRALS

If the function $u = g(x)$ has a continuous derivative on the closed interval $[a, b]$ and f is continuous on the range of g , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du.$$

Math for IT (Somkiat Wangsiripitak)

69

EXAMPLE 8 Change of Variables

Evaluate $\int_0^1 x(x^2 + 1)^3 \, dx$.

Solution To evaluate this integral, let $u = x^2 + 1$. Then, you obtain

$$u = x^2 + 1 \Rightarrow du = 2x \, dx.$$

Before substituting, determine the new upper and lower limits of integration.

Lower Limit

When $x = 0$, $u = 0^2 + 1 = 1$.

Upper Limit

When $x = 1$, $u = 1^2 + 1 = 2$.

Now, you can substitute to obtain

$$\int_0^1 x(x^2 + 1)^3 \, dx = \frac{1}{2} \int_1^2 (x^2 + 1)^3 (2x) \, dx$$

Integration limits for x

$$= \frac{1}{2} \int_1^2 u^3 \, du$$

Integration limits for u

$$= \frac{1}{2} \left[\frac{u^4}{4} \right]_1^2$$

$$= \frac{1}{2} \left(4 - \frac{1}{4} \right)$$

$$= \frac{15}{8}$$

Try rewriting the antiderivative $\frac{1}{2}(u^4/4)$ in terms of the variable x and evaluate the definite integral at the original limits of integration, as shown.

$$\frac{1}{2} \left[\frac{u^4}{4} \right]_1^2 = \frac{1}{2} \left[\frac{(x^2 + 1)^4}{4} \right]_0^1$$

$$= \frac{1}{2} \left(4 - \frac{1}{4} \right) = \frac{15}{8}$$

Notice that you obtain the same result.

EXAMPLE 9 Change of Variables

Evaluate $A = \int_1^5 \frac{x}{\sqrt{2x-1}} \, dx$.

u = \sqrt{2x-1}

u^2 + 1 = 2x - 1

u^2 + 1 = x

u du = dx

when $x=1$
 $u = \sqrt{2-1} = 1$
 $x=5$
 $u = \sqrt{10-1} = 3$

$$\begin{aligned} \int_1^5 \frac{x}{\sqrt{2x-1}} \, dx &= \int_1^3 \frac{1}{\cancel{u}} \left(\frac{u^2+1}{2} \right) \cancel{u} \, du \\ &= \frac{1}{2} \int_1^3 (u^2+1) \, du \\ &= \frac{1}{2} \left[\frac{u^3}{3} + u \right]_1^3 \\ &= \frac{1}{2} \left(9 + 3 - \frac{1}{3} - 1 \right) \\ &= \frac{16}{3} \end{aligned}$$

Math for IT (Somkiat Wangsiripitak)

72

Homework

Find

1) $\int_{-1}^1 \left(6x^5 - \frac{2}{\sqrt[5]{x^3}} + 3 \right) dx$

2) $\int_1^4 |x - 3| dx$

3) $\int \frac{(2x^2+1)}{(2x^3+3x)^2} dx$

- Write down a step-by-step solution in a piece of A4 paper (or on an electronic paper).
- **DO NOT TYPE.**
- Submit your solution in MS Team.
- Do not forget to write your name and student ID.