

Vector Spaces #2

Instructor: Dr. Praphan Pavarangkoon

Office: 4th Floor

Email: praphan@it.kmitl.ac.th

**Office hours: Thursday at 9:00 – 11:00
or as an advance appointment**

Outline

- Vector Spaces
- Subspaces of Vector Spaces
- Linear Combination of Vectors
- Spanning Sets
- Linear Independence

สัจพจน์ (Axiom)

- หมายถึง ข้อความที่ยอมรับว่าเป็นจริงโดยไม่ต้องพิสูจน์ ซึ่งตรงข้ามกับคำว่า “ทฤษฎีบท” ซึ่งจะถูกรับว่าเป็นจริงได้ก็ต่อเมื่อมีการพิสูจน์
- ดังนั้นสัจพจน์จึงถูกใช้เป็นจุดเริ่มต้นในการพิสูจน์ทางคณิตศาสตร์ และทฤษฎีบททุกอัน จะต้องอนุมาน (inference) มายังสัจพจน์ได้ เช่น
 - เราสามารถลากเส้นตรงผ่านจุดสองจุดได้
 - เราสามารถขยายส่วนของเส้นตรงไปเป็นเส้นตรงได้เส้นเดียว
 - มุมจากทุกมุมย่อมเท่ากัน
 - สมการที่ถูกรวบรวมด้วยค่าเท่ากันทั้งสองข้างก็ยังเป็นสมการอยู่

เซต (Set)

- หมายถึง กลุ่มของสิ่งใดสิ่งหนึ่ง
- เซต = {สมาชิกของเซตตัวที่ 1, สมาชิกของเซตตัวที่ 2, ... }
- สมาชิกของเซต จะอยู่ภายใต้เครื่องหมาย **ปีกกา { }**

เช่น เซต A คือของจำนวนคู่ที่เป็นบวก จะได้ว่า

$$A = \{2, 4, 6, \dots\}$$

การเป็นสมาชิกของเซต ใช้เครื่องหมาย $\in$ เช่น

- เซตที่เป็นที่รู้จัก เช่น
 - เซตของจำนวนจริง แทนด้วย R
 - เซตของคู่ลำดับในระนาบ แทนด้วย R^2
 - เซตของจำนวนเต็มบวก แทนด้วย I^+
 - เซตของเมทริกซ์ ขนาด $m \times n$ แทนด้วย $M_{m \times n}$

Vector Spaces

- Any set that satisfies these properties (or **axioms**) is called a **vector space**, and the objects in the set are called **vectors**.

Let V be a set on which two operations (**vector addition** and **scalar multiplication**) are defined. If the listed axioms are satisfied for every $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ in V and every scalar (real number) c and d , then V is called a **vector space**.

Addition:

1. $\mathbf{u} + \mathbf{v}$ is in V .
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
4. V has a **zero vector** $\mathbf{0}$ such that for every $\mathbf{u}$ in V , $\mathbf{u} + \mathbf{0} = \mathbf{u}$.
5. For every $\mathbf{u}$ in V , there is a vector in V denoted by $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$.

Closure under addition

Commutative property

Associative property

Additive identity

Additive inverse

Scalar Multiplication:

6. $c\mathbf{u}$ is in V .
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1(\mathbf{u}) = \mathbf{u}$

Closure under scalar multiplication

Distributive property

Distributive property

Associative property

Scalar identity

R^2 with the Standard Operations Is a Vector Space

The set of all ordered pairs of real numbers R^2 with the standard operations is a vector space. To verify this, look back at Theorem 4.1. Vectors in this space have the form

$$\mathbf{v} = (v_1, v_2).$$

Let $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ be vectors in the plane, and let c and d be scalars.

- | | |
|--|--|
| 1. $\mathbf{u} + \mathbf{v}$ is a vector in the plane. | Closure under addition |
| 2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ | Commutative property of addition |
| 3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ | Associative property of addition |
| 4. $\mathbf{u} + \mathbf{0} = \mathbf{u}$ | Additive identity property |
| 5. $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ | Additive inverse property |
| 6. $c\mathbf{u}$ is a vector in the plane. | Closure under scalar multiplication |
| 7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ | Distributive property |
| 8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ | Distributive property |
| 9. $c(d\mathbf{u}) = (cd)\mathbf{u}$ | Associative property of multiplication |
| 10. $1(\mathbf{u}) = \mathbf{u}$ | Multiplicative identity property |

R^n with the Standard Operations Is a Vector Space

The set of all ordered n -tuples of real numbers R^n with the standard operations is a vector space. This is verified by Theorem 4.2. Vectors in this space are of the form

$$\mathbf{v} = (v_1, v_2, v_3, \dots, v_n).$$

Let $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ be vectors in R^n , and let c and d be scalars.

- | | |
|--|--|
| 1. $\mathbf{u} + \mathbf{v}$ is a vector in R^n . | Closure under addition |
| 2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ | Commutative property of addition |
| 3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ | Associative property addition |
| 4. $\mathbf{u} + \mathbf{0} = \mathbf{u}$ | Additive identity property |
| 5. $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ | Additive inverse property |
| 6. $c\mathbf{u}$ is a vector in R^n . | Closure under scalar multiplication |
| 7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ | Distributive property |
| 8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$ | Distributive property |
| 9. $c(d\mathbf{u}) = (cd)\mathbf{u}$ | Associative property of multiplication |
| 10. $1(\mathbf{u}) = \mathbf{u}$ | Multiplicative identity property |

The Vector Space of All 2×3 Matrices

Show that the set of all 2×3 matrices with the operations of matrix addition and scalar multiplication is a vector space.

$$\text{ex. } u = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix} \quad v = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix}$$

$$u+v = \begin{bmatrix} a_{11}+b_{11} & a_{12}+b_{12} & a_{13}+b_{13} \\ a_{21}+b_{21} & a_{22}+b_{22} & a_{23}+b_{23} \end{bmatrix}$$

The Vector Space of All Polynomials of Degree 2 or Less

Let P_2 be the set of all polynomials of the form

$$p(x) = a_2x^2 + a_1x + a_0,$$

where a_0 , a_1 , and a_2 are real numbers. The *sum* of two polynomials $p(x) = a_2x^2 + a_1x + a_0$ and $q(x) = b_2x^2 + b_1x + b_0$ is defined in the usual way by

$$p(x) + q(x) = (a_2 + b_2)x^2 + (a_1 + b_1)x + (a_0 + b_0),$$

and the *scalar multiple* of $p(x)$ by the scalar c is defined by

$$cp(x) = ca_2x^2 + ca_1x + ca_0.$$

Show that P_2 is a vector space.

Properties of Scalar Multiplication

Let $\mathbf{v}$ be any element of a vector space V , and let c be any scalar. Then the following properties are true.

1. $0\mathbf{v} = \mathbf{0}$

2. $c\mathbf{0} = \mathbf{0}$

3. If $c\mathbf{v} = \mathbf{0}$, then $c = 0$ or $\mathbf{v} = \mathbf{0}$.

4. $(-1)\mathbf{v} = -\mathbf{v}$

The Set of Integers Is Not a Vector Space

The set of all integers (with the standard operations) does not form a vector space because it is not closed under scalar multiplication. For example,

$$\frac{1}{2}(1) = \frac{1}{2}.$$

Scalar Integer Noninteger

The Set of Second-Degree Polynomials Is Not a Vector Space

The set of all second-degree polynomials is not a vector space because it is not closed under addition. To see this, consider the second-degree polynomials

$$p(x) = x^2 \quad \text{and} \quad q(x) = -x^2 + x + 1,$$

whose sum is the first-degree polynomial

$$p(x) + q(x) = x + 1.$$

A Set That Is Not a Vector Space

Let $V = \mathbb{R}^2$, the set of all ordered pairs of real numbers, with the standard operation of addition and the *nonstandard* definition of scalar multiplication listed below.

$$c(x_1, x_2) = (cx_1, 0)$$

Show that V is not a vector space.

Subspaces of Vector Spaces

- In most important applications in linear algebra, vector spaces occur as **subspaces** of larger spaces.
- For instance, you will see that the solution set of a homogeneous system of linear equations in n variables is a subspace of R^n .

A nonempty subset W of a vector space V is called a **subspace** of V if W is a vector space under the operations of addition and scalar multiplication defined in V .

REMARK: Note that if W is a subspace of V , it must be closed under the operations inherited from V .

A Subspace of R^3

Show that the set $W = \{(x_1, 0, x_3) : x_1 \text{ and } x_3 \text{ are real numbers}\}$ is a subspace of R^3 with the standard operations.

Test for a Subspace

If W is a nonempty subset of a vector space V , then W is a subspace of V if and only if the following closure conditions hold.

1. If $\mathbf{u}$ and $\mathbf{v}$ are in W , then $\mathbf{u} + \mathbf{v}$ is in W .
2. If $\mathbf{u}$ is in W and c is any scalar, then $c\mathbf{u}$ is in W .

The Subspace of $M_{2,2}$

Let W be the set of all 2×2 symmetric matrices. Show that W is a subspace of the vector space $M_{2,2}$, with the standard operations of matrix addition and scalar multiplication.

The Set of Singular Matrices Is Not a Subspace of $M_{n,n}$

Let W be the set of singular matrices of order 2. Show that W is not a subspace of $M_{2,2}$ with the standard operations.

The Set of First Quadrant Vectors Is Not a Subspace of $\mathbb{R}^2$

Show that $W = \{(x_1, x_2): x_1 \geq 0 \text{ and } x_2 \geq 0\}$, with the standard operations, is not a subspace of $\mathbb{R}^2$.

Linear Combination of Vectors

- This section begins to develop procedures for representing each vector in a vector space as a **linear combination** of a select number of vectors in the space.

A vector $\mathbf{v}$ in a vector space V is called a **linear combination** of the vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ in V if $\mathbf{v}$ can be written in the form

$$\mathbf{v} = c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + \dots + c_k\mathbf{u}_k,$$

where $c_1, c_2, \dots, c_k$ are scalars.

Examples of Linear Combinations

(a) For the set of vectors in R^3 ,

$$S = \{ \overset{\mathbf{v}_1}{(1, 3, 1)}, \overset{\mathbf{v}_2}{(0, 1, 2)}, \overset{\mathbf{v}_3}{(1, 0, -5)} \},$$

$\mathbf{v}_1$ is a linear combination of $\mathbf{v}_2$ and $\mathbf{v}_3$ because

$$\begin{aligned} \mathbf{v}_1 &= 3\mathbf{v}_2 + \mathbf{v}_3 = 3(0, 1, 2) + (1, 0, -5) \\ &= (1, 3, 1). \end{aligned}$$

(b) For the set of vectors in $M_{2,2}$,

$$S = \left\{ \overset{\mathbf{v}_1}{\begin{bmatrix} 0 & 8 \\ 2 & 1 \end{bmatrix}}, \overset{\mathbf{v}_2}{\begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix}}, \overset{\mathbf{v}_3}{\begin{bmatrix} -1 & 3 \\ 1 & 2 \end{bmatrix}}, \overset{\mathbf{v}_4}{\begin{bmatrix} -2 & 0 \\ 1 & 3 \end{bmatrix}} \right\},$$

$\mathbf{v}_1$ is a linear combination of $\mathbf{v}_2$, $\mathbf{v}_3$, and $\mathbf{v}_4$ because

$$\begin{aligned} \mathbf{v}_1 &= \mathbf{v}_2 + 2\mathbf{v}_3 - \mathbf{v}_4 \\ &= \begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix} + 2 \begin{bmatrix} -1 & 3 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} -2 & 0 \\ 1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 8 \\ 2 & 1 \end{bmatrix}. \end{aligned}$$

Finding a Linear Combination

Write the vector $\mathbf{w} = (1, 1, 1)$ as a linear combination of vectors in the set S .

$$S = \{(\overset{\mathbf{v}_1}{1}, \overset{\mathbf{v}_2}{2}, \overset{\mathbf{v}_3}{3}), (0, 1, 2), (-1, 0, 1)\}$$

Finding a Linear Combination

If possible, write the vector $\mathbf{w} = (1, -2, 2)$ as a linear combination of vectors in the set S from Example 2.



$$c_9 = t$$

Spanning Sets

- If every vector in a vector space can be written as a linear combination of vectors in a set S , then S is called a **spanning set** of the vector space.

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ be a subset of a vector space V . The set S is called a **spanning set** of V if *every* vector in V can be written as a linear combination of vectors in S . In such cases it is said that S **spans** V .

Examples of Spanning Sets

- (a) The set $S = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ spans R^3 because any vector $\mathbf{u} = (u_1, u_2, u_3)$ in R^3 can be written as

$$\mathbf{u} = u_1(1, 0, 0) + u_2(0, 1, 0) + u_3(0, 0, 1) = (u_1, u_2, u_3).$$

- (b) The set $S = \{1, x, x^2\}$ spans P_2 because any polynomial function $p(x) = a + bx + cx^2$ in P_2 can be written as

$$\begin{aligned} p(x) &= a(1) + b(x) + c(x^2) \\ &= a + bx + cx^2. \end{aligned}$$

A Spanning Set of $\mathbb{R}^3$

Show that the set $S = \{(1, 2, 3), (0, 1, 2), (-2, 0, 1)\}$ spans $\mathbb{R}^3$.

A Set That Does Not Span R^3

From Example 3 you know that the set

$$S = \{(1, 2, 3), (0, 1, 2), (-1, 0, 1)\}$$

does not span R^3 because $\mathbf{w} = (1, -2, 2)$ is in R^3 and cannot be expressed as a linear combination of the vectors in S .

The Span of a Set

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a set of vectors in a vector space V , then the **span of S** is the set of all linear combinations of the vectors in S ,

$$\text{span}(S) = \{c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k : c_1, c_2, \dots, c_k \text{ are real numbers}\}.$$

The span of S is denoted by $\text{span}(S)$ or $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$. If $\text{span}(S) = V$, it is said that V is **spanned** by $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$, or that S **spans** V .

Span(S) Is a Subspace of V

If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ is a set of vectors in a vector space V , then $\text{span}(S)$ is a subspace of V . Moreover, $\text{span}(S)$ is the smallest subspace of V that contains S , in the sense that every other subspace of V that contains S must contain $\text{span}(S)$.

PROOF To show that $\text{span}(S)$, the set of all linear combinations of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$, is a subspace of V , show that it is closed under addition and scalar multiplication. Consider any two vectors $\mathbf{u}$ and $\mathbf{v}$ in $\text{span}(S)$,

$$\mathbf{u} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k$$

$$\mathbf{v} = d_1\mathbf{v}_1 + d_2\mathbf{v}_2 + \dots + d_k\mathbf{v}_k,$$

where $c_1, c_2, \dots, c_k$ and $d_1, d_2, \dots, d_k$ are scalars. Then

$$\mathbf{u} + \mathbf{v} = (c_1 + d_1)\mathbf{v}_1 + (c_2 + d_2)\mathbf{v}_2 + \dots + (c_k + d_k)\mathbf{v}_k$$

and

$$c\mathbf{u} = (cc_1)\mathbf{v}_1 + (cc_2)\mathbf{v}_2 + \dots + (cc_k)\mathbf{v}_k,$$

which means that $\mathbf{u} + \mathbf{v}$ and $c\mathbf{u}$ are also in $\text{span}(S)$ because they can be written as linear combinations of vectors in S . So, $\text{span}(S)$ is a subspace of V . It is left to you to prove that $\text{span}(S)$ is the smallest subspace of V that contains S .

Linear Dependence and Linear Independence

A set of vectors $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ in a vector space V is called **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}$$

has only the trivial solution, $c_1 = 0, c_2 = 0, \dots, c_k = 0$. If there are also nontrivial solutions, then S is called **linearly dependent**.

Examples of Linearly Dependent Sets

(a) The set $S = \{(1, 2), (2, 4)\}$ in R^2 is linearly dependent because

$$-2(1, 2) + (2, 4) = (0, 0).$$

(b) The set $S = \{(1, 0), (0, 1), (-2, 5)\}$ in R^2 is linearly dependent because

$$2(1, 0) - 5(0, 1) + (-2, 5) = (0, 0).$$

(c) The set $S = \{(0, 0), (1, 2)\}$ in R^2 is linearly dependent because

$$1(0, 0) + 0(1, 2) = (0, 0).$$

Testing for Linear Independence

Determine whether the set of vectors in R^3 is linearly independent or linearly dependent.

$$S = \{ \overset{\mathbf{v}_1}{(1, 2, 3)}, \overset{\mathbf{v}_2}{(0, 1, 2)}, \overset{\mathbf{v}_3}{(-2, 0, 1)} \}$$

Testing for Linear Independence and Linear Dependence

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ be a set of vectors in a vector space V . To determine whether S is linearly independent or linearly dependent, perform the following steps.

1. From the vector equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}$, write a homogeneous system of linear equations in the variables $c_1, c_2, \dots$, and c_k .
2. Use Gaussian elimination to determine whether the system has a unique solution.
3. If the system has only the trivial solution, $c_1 = 0, c_2 = 0, \dots, c_k = 0$, then the set S is linearly independent. If the system also has nontrivial solutions, then S is linearly dependent.

Testing for Linear Independence

Determine whether the set of vectors in P_2 is linearly independent or linearly dependent.

$$S = \{\overset{\mathbf{v}_1}{1 + x - 2x^2}, \overset{\mathbf{v}_2}{2 + 5x - x^2}, \overset{\mathbf{v}_3}{x + x^2}\}$$

Testing for Linear Independence

Determine whether the set of vectors in $M_{2,2}$ is linearly independent or linearly dependent.

$$S = \left\{ \overset{\mathbf{v}_1}{\begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}}, \overset{\mathbf{v}_2}{\begin{bmatrix} 3 & 0 \\ 2 & 1 \end{bmatrix}}, \overset{\mathbf{v}_3}{\begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}} \right\}$$

A Property of Linearly Dependent Sets

A set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$, $k \geq 2$, is linearly dependent if and only if at least one of the vectors $\mathbf{v}_j$ can be written as a linear combination of the other vectors in S .

PROOF To prove the theorem in one direction, assume S is a linearly dependent set. Then there exist scalars $c_1, c_2, c_3, \dots, c_k$ (not all zero) such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + \dots + c_k\mathbf{v}_k = \mathbf{0}.$$

Because one of the coefficients must be nonzero, no generality is lost by assuming $c_1 \neq 0$. Then solving for $\mathbf{v}_1$ as a linear combination of the other vectors produces

$$c_1\mathbf{v}_1 = -c_2\mathbf{v}_2 - c_3\mathbf{v}_3 - \dots - c_k\mathbf{v}_k$$

$$\mathbf{v}_1 = -\frac{c_2}{c_1}\mathbf{v}_2 - \frac{c_3}{c_1}\mathbf{v}_3 - \dots - \frac{c_k}{c_1}\mathbf{v}_k.$$

Conversely, suppose the vector $\mathbf{v}_1$ in S is a linear combination of the other vectors. That is,

$$\mathbf{v}_1 = c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + \dots + c_k\mathbf{v}_k.$$

Then the equation $-\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + \dots + c_k\mathbf{v}_k = \mathbf{0}$ has at least one coefficient, -1 , that is nonzero, and you can conclude that S is linearly dependent.

Writing a Vector as a Linear Combination of Other Vectors

In Example 9, you determined that the set

$$S = \{1 + x - 2x^2, 2 + 5x - x^2, x + x^2\}$$

is linearly dependent. Show that one of the vectors in this set can be written as a linear combination of the other two.

Corollary

Two vectors $\mathbf{u}$ and $\mathbf{v}$ in a vector space V are linearly dependent if and only if one is a scalar multiple of the other.

REMARK: The zero vector is always a scalar multiple of another vector in a vector space.

Testing for Linear Dependence of Two Vectors

(a) The set

$$S = \{(\overset{\mathbf{v}_1}{1}, \overset{\mathbf{v}_1}{2}, 0), (\overset{\mathbf{v}_2}{-2}, \overset{\mathbf{v}_2}{2}, 1)\}$$


is linearly independent because $\mathbf{v}_1$ and $\mathbf{v}_2$ are not scalar multiples of each other, as shown in Figure 4.17(a).

(b) The set

$$S = \{(\overset{\mathbf{v}_1}{4}, \overset{\mathbf{v}_1}{-4}, -2), (\overset{\mathbf{v}_2}{-2}, \overset{\mathbf{v}_2}{2}, 1)\}$$

is linearly dependent because $\mathbf{v}_1 = -2\mathbf{v}_2$, as shown in Figure 4.17(b).


(a)



$$S = \{(1, 2, 0), (-2, 2, 1)\}$$

The set S is linearly independent.

(b)



$$S = \{(4, -4, -2), (-2, 2, 1)\}$$

The set S is linearly dependent because $\mathbf{v}_1 = -2\mathbf{v}_2$.

Figure 4.17

Q & A