

[[Xavier: Measuring precise radia velocities and cross-correlation function line-profile variations using a Skew Normal distribution]]

U. Simola

X. Dumusque

Jessi Cisewski-Kehe

May 15, 2018

Abstract

[[Xavier: Stellar activity is the main limitation to the detection of small-mass exoplanets using the radial-velocity (RV) technique. Stellar activity can be probed by measuring the cross-correlation function (CCF) shape variations as a function of time. Those variations are measured using the different moments of the CCF, therefore measuring with the best precision those moments is crucial]] to de-correlate exoplanet signals from spurious RV signals originating from stellar activity.

We propose here to measure those moments using a Skew Normal (SN) distribution, which compared to the Normal distribution generally used, naturally provides an extra parameter to model the CCF natural asymmetry induced by convective blueshift.

We analyze 5 stars with different activity levels and different signal-to-noise ratio levels. In each case, we compare the results obtained by fitting to the CCF respectively a Normal and a SN. We also estimate rigorous errors for the different moments of the CCF using a bootstrap analysis.

The correlations between the RVs and the CCF asymmetry or RVs and the CCF width is always stronger when using the parameters derived from the SN [[Xavier: in the case of real observations]]. Therefore the CCF asymmetry and width derived using a SN are more sensitive to stellar activity, which [[Xavier: allows to probe with a better precision stellar rotational periods, in addition to characterize more precisely stellar activity signals.]] [[Xavier: The precision on the RVs, derived using the median of the fitted SN distributions]] are on average 10% smaller than the RV errors calculated on the mean of the Normal. In addition, the asymmetry parameter derived from the SN is on average 15% better than the one calculated on the common Bisector Inverse Slope Span (BIS SPAN).

[[Xavier: We strongly encourage the use of the SN distribution rather than the Normal distribution, because this allows to retrieve in one single fit the different moments of the CCF, because the derived moments probe better stellar activity signals and because the standard errors on the RV and the the asymmetry are smaller.]] [[Xavier: I just left this comment here so that we can check at the end that we answered to all referee's major comments, however, I believe it is not relevant anymore with our new analysis.]] [[Referee: You simply cannot make this claim. What you have

shown is that the SN method produces errors that are 60% larger than the Normal distribution and you hint that this is due to it being more sensitive to activity. You have not demonstrated this by looking at the "activity" signal that the SN isolates to see if it is actually more sensitive at finding rotation periods, activity cycles, etc. Where is the sensitivity in actually getting out useful information? Furthermore, after correcting the SN RVs for the activity, you arrive at the same answer as the Normal distribution method. I would conclude that the SN offers no real advantages over Normal so the community can continue to use the standard method. Grammatically this is an ambiguous statement. Are you looking for "better signals", i.e. ones that you like, or probing stellar signals in a better way? (This is what you meant, but not what was stated.))]

1 Introduction

[[**Jessi:** I added subsections here for our reference to make sure the bases are covered, but we should remove them before submission.]]

1.1 Goal of RV analysis

[[**Xavier:** When working with radial velocities, the main limitation to the detection of small-mass exoplanets is not anymore the precision of the instruments used, but the different noises induce by the stars we are observing (Dumusque et al. 2017). Indeed, stellar oscillations, granulation phenomena and stellar activity all induce RV signals (e.g. Saar and Donahue 1997; Queloz et al. 2001; Desort et al. 2007; Dumusque et al. 2011; Dumusque 2016) that are beyond the meter-per-second precision reached by the best high-resolution spectrographs. It is therefore mandatory to understand better stellar signals and to find ways to correct for them, if in the near future we want to detect or confirm an Earth-twin planet using the RV technique. This is even more true now that instrument like ESPRESSO (Pepe et al. 2014) and EXPRESS (Fischer et al. 2017) should have the stability to detect such signals. However, if solutions are not found to mitigate the impact of stellar activity, the detection or confirmation of potential Earth-twins will be extremely challenging.]]

1.2 Stellar activity effect on CCF, Normal fit plus FWHM and BIS SPAN indicators

[[**Jessi:** Eventually three methods are introduced for detecting CCF asymmetry: FWHM, bisector span, and the Figueira approach - it seems that these methods need to be better explained. Somewhere in the intro it needs to become clear that we are (i) providing a new way of measuring CCF asymmetry, (ii) this is a unified approach that also provides an RV estimate, and (iii) it (hopefully) provides a better indicator.]]

[[**Xavier:** Among the different stellar signals we are aware of, the one that is the most difficult to characterize and to correct for is the signal induced by stellar activity. Stellar activity is responsible for creating magnetic regions on the surface of stars, and those regions change locally the temperature and the convection, inducing spurious RV variations (e.g. Meunier et al. 2010; Dumusque et al. 2014). In theory, it should be easy to differentiate between the pure Doppler-shift

induced by a planet, that will shift the entire stellar spectrum, and stellar activity that modifies the shape of spectral lines and by doing so create a spurious shift of the stellar spectrum (Saar and Donahue 1997; Hatzes 2002; Kurster et al. 2003; Lindegren and Dravins 2003; Desort et al. 2007; Lagrange et al. 2010; Meunier et al. 2010; Dumusque et al. 2014). However, on quiet GKM dwarfs, the main target for precise RV measurements, stellar activity induce signals of a few m s^{-1} . This corresponds physically to variations smaller than 1/100th of a pixel on the detector. To be able to measure such tiny variations, we average the information of all the line in the spectrum by cross correlating the stellar spectrum with a synthetic (Baranne et al. 1996; Pepe et al. 2002) or an observed stellar template (Anglada-Escudé and Butler 2012). The result of this operation gives us the cross-correlation function (CCF). To measure the Doppler-shift between different spectra, and therefore get the RV of a star as a function of time, we calculate the variations of the CCF barycenter. The barycenter is generally estimated by fitting a Normal distribution to the CCF, and taking its mean. Variation in line shape between different spectra, which indicate the presence of signals induced by stellar activity, is measured by analysing the different moments of the CCF. Generally, the width of the CCF is estimated using the FWHM of the fitted Normal distribution, and its asymmetry using the the bisector inverse slope span (BIS SPAN, Queloz et al. 2001).]]

1.3 Why FWHM and BIS SPAN important?

[[**Xavier:** If a RV signal is induced by activity, generally a strong correlation will be observed between the RV and chromospheric activity indicators like $\log(R'_{HK})$ or $H-\alpha$ (e.g. Boisse et al. 2009; Dumusque et al. 2012; Robertson et al. 2014), but also between the RV and the FWHM of the CCF or its BIS SPAN (Queloz et al. 2001; Boisse et al. 2009; Queloz et al. 2009; Dumusque 2016). It is therefore common now, that when fitting for a planetray signal, in addition to a Keplerian, the model includes in addition linear dependancies with the $\log(R'_{HK})$, the FWHM and the BIS SPAN (e.g. Dumusque et al. 2017; Feng et al. 2017). It is also common to add a Gaussian process to the model to account for the correlated noise induced by stellar activity. The hyperparameters of the Gaussian process are generally trained on the different activity indicators (e.g. Haywood et al. 2014; Rajpaul et al. 2015). It is therefore essential for mitigating stellar activity to obtain activity indicators that are the most correlated to the RVs but also for which we can obtain the best precision.]]

1.4 Figueira indicators of stellar activity + other

[[**Xavier:** Several works have derived indicators that are more sensitive to line asymmetry than the BIS SPAN. In Boisse et al. (2011), the authors develop V_{span} a new indicator to derive the CCF asymmetry, that is more sensitive than the BIS SPAN at low signal-to-noise ratio (SNR). Figueira et al. (2013) studied the use of two new indicators, bi-Gauss and V_{asy} . The authors were able to show that when using bi-Gauss, the amplitude in asymmetry is 30% larger, therefore allowing the detection of smaller-amplitude correlations with radial-velocity variations. They also demonstrated that V_{asy} seems to be the best indicator of line asymmetry at high SNR, as its correlation with RV

is more significant than any other asymmetry indicators.]]

1.5 Why using a SN distribution?

[[**Xavier:** In all the method described above, the RV and the FWHM is derived using a Normal distribution fitted to the CCF, and the asymmetry is estimated using another approach. In this paper we propose to use a Skew Normal (SN) distribution to derive at the same time the RV, the FWHM and the asymmetry of the CCF, as this function includes a skewness parameter (Azzalini 1985). In addition, we know that for solar-type stars and cooler dwarfs, the bisector of the CCF as a "C"-shape due to convective blueshift (e.g. Dravins et al. 1981; Gray 2009). Therefore, fitting the CCF using a model that naturally includes an asymmetry, like the SN distribution, should give more precise results.]]

1.6 Outline of paper

[[**TODO:** This paragraph will need to be updated once the rest is finished.]] The paper is organized as follow. In Sec. 2 we introduce the SN distribution and describe its applicability for modeling the CCF, show that the SN distribution is a better representation of observed CCF than a Normal distribution, and study how the SN parameters relate to the RV, FWHM and BIS SPAN of the CCF. In Sec. 1 we present a simple model to correct for stellar activity. In Sec. 4, we compare on real observations the sensitivity of the SN parameters to stellar activity with respect to other existing indicators. In Sec. 5 we derive error bars for the different CCF parameters, and finally we discuss our results and conclude in Sec. 6 and Sec. 7.

2 The Skew Normal distribution

The Skew Normal (SN) distribution is a class of probability distributions which includes the Normal distribution as a special case (Azzalini 1985). The SN distribution has, in addition to a location and a scale parameter analogous to the Normal distribution's mean and standard deviation, a third parameter which describes the asymmetry, or the skewness, of the distribution. Considering a random variable $Y \in \mathbb{R}$ (where \mathbb{R} is the real line) which follows a SN distribution with location parameter $\xi \in \mathbb{R}$, scale parameter $\omega \in \mathbb{R}^+$ (i.e., the positive real line), and skewness parameter $\alpha \in \mathbb{R}$, its density at some value $Y = y$ can be written as

$$SN(y; \xi, \omega, \alpha) = \frac{2}{\omega} \phi\left(\frac{y - \xi}{\omega}\right) \Phi\left(\frac{\alpha(y - \xi)}{\omega}\right), \quad (1)$$

where ϕ and Φ are respectively the density function and the distribution function of a *standard* Normal distribution¹ and $\alpha \in \mathbb{R}$ is the skewness parameter which quantifies the asymmetry of the SN. We then write $Y \sim SN(\xi, \omega^2, \alpha)$ to mean that the random variable Y follows the noted

¹A standard Normal distribution is a Normal distribution with a mean of 0 and a standard deviation of 1.

SN distribution. Examples of SN densities under different skewness parameter values and the same location and scale parameters ($\xi = 0$ and $\omega = 1$) are displayed in Fig. 1. A usual Normal distribution is the special case of the SN distribution when the skewness parameter, α , is equal to 0.² For reasons related to the interpretation of the parameters in Equation (1) and computational issues




Figure 1: Density function of a random variable $Y \sim SN(\xi, \omega^2, \alpha)$ with location parameter $\xi = 0$, scale parameter $\omega = 1$ and different values of the skewness parameter α indicated by different colors and line types. Note that the solid black line has an $\alpha = 0$, making it a Normal distribution.

with estimating α near 0, a different parametrization is used, which is referred to as the *centered parametrization* (CP). We will be using the CP in this work, which includes a mean parameter μ , a variance parameter σ^2 and a skewness parameter γ . In order to define the CP, we need to express the CP parameters (μ, σ^2, γ) as a function of the one used in the Equation (1) with (ξ, ω^2, α) by

$$\mu = \xi + \omega\beta, \quad \sigma^2 = \omega^2(1 - \beta^2), \quad \gamma = \frac{1}{2}(4 - \pi)\beta^3(1 - \beta^2)^{-3/2}, \quad (2)$$

where $\beta = \sqrt{\frac{2}{\pi}} \left(\frac{\alpha}{\sqrt{1 + \alpha^2}} \right)$.

By using Eq. (2), the new set of parameters (μ, σ^2, γ) provides a more clear interpretation of the behavior of the SN distribution. For the α values used in Fig. 1, the corresponding values of μ , σ^2 , γ are displayed in Table 1. In particular, μ and σ^2 are the actual mean and variance of

²This can be seen from Eq. (1). If $\alpha = 0$ then $\Phi\left(\frac{\alpha(y-\xi)}{\omega}\right) = \Phi(0)$; this is the probability a standard Normal random variable is less than or equal to 0, which is 0.5. The 0.5 cancels with the 2 in the denominator and what remains is the usual Normal density, $\frac{1}{\omega} \phi\left(\frac{y-\xi}{\omega}\right)$

the distribution (rather than simply a location and scale parameter) and γ becomes an index for evaluating the skewness of the SN.

Beyond the mean of the SN, we introduce a second location parameter that will be largely used in the analyses: the median. The median of the SN, and in general the median of an absolute continuous random variable, is defined as that value m such that

$$\int_{-\infty}^m SN(y; \xi, \omega, \alpha) = \frac{1}{2}, \quad (3)$$

where $SN(y; \xi, \omega, \alpha)$ follows Eq. (1).³

α	μ	σ^2	γ
-3	-0.757	0.427	-0.667
0	0.000	1.000	0.000
2	0.714	0.491	0.454
6	0.787	0.381	0.891
10	0.794	0.370	0.956

Table 1: CP values, (μ, σ^2, γ) , corresponding to the α values from Fig. 1 (with location parameter $\xi = 0$ and scale parameter $\omega = 1$) using Eq. (2). Values are rounded to three decimal places.

Further details about the parametrization from Eq. (1) (called *Direct Parametrization* or DP), the CP, and general statistical properties of the SN are treated in rigorous mathematical and statistical viewpoints in the book by Azzalini and Capitanio (2014).

2.1 Fitting the Skew Normal distribution to the CCF

The SN density shape is used to model the CCF. In particular, using least-squares algorithm, we fit the following function:

$$f_{CCF}(y_i) = C - A \times SN(y_i; \mu, \sigma^2, \gamma), \quad i = 1, \dots, n \quad (4)$$

where C is an unknown offset fitting the continuum of the CCF, A is an unknown amplitude parameter and y_1, \dots, y_n are the set of RV's considered for the CCF. Note that the CCF is expressed in flux as a function of the lag of the cross-correlation template, expressed in RV.

Since the CCF presents a natural asymmetry due the convective blueshift, the SN distribution should in principle better catch this aspect, together with other changes in asymmetry, respect the Normal fit. To initially check this assumption, we compared the CCF residuals after fitting a Normal and a SN distribution for 2 stars. The first star is Alpha Centauri b, whose CCFs have high signal-to-noise ratio (SNR). The second star is Corot-7, whose CCFs have low SNR. Fig. 2 shows that the SN seems to be a slightly better model to explain the shape of the CCF, in particular as the SNR decreases.

³We recall that when using a symmetric distribution such as the Normal distribution, the mean and the median are equivalent.




Figure 2: Comparison between the Normal (black circles) and the SN (red crosses) residuals using CCFs from the star Alpha Centauri b (left) and Corot-7 (right). When looking at the residuals corresponding to the tails of the CCF, the results derived by the two fits are comparable. However, when focusing on the center of the CCF, the SN fit leads to slightly better results for both the stars. Moreover, as the SNR decreases, the SN distribution shows smaller residuals respect the Normal ones.

In the following of the paper, we define RV as the mean of the Normal distribution. Concerning the fit of the CCF using the SN, we present at first 2 indicators that define the RV of the star: the mean of the SN, defined as RV mean SN and the median of the SN, defined as RV median SN (i.e. realling Eq.(3), $m = \text{RV median SN}$). We will discuss advantages and limits for both these choices in Section 4 and Section 5. For the width of the CCF, we use the FWHM of the Normal, which is $2\sqrt{2\ln 2}\sigma$. The width of the SN, SN FWHM, is defined in the same way⁴. Being a Normal distribution symmetric, there is not such a parameter that evaluates the asymmetry of the distribution, so the BIS SPAN is used. The BIS SPAN will be compared to the asymmetric parameter γ of the SN, named SN GAMMA. To test the strength of the correlation between the RVs and the different indicators for stellar activity, we calculated the Pearson correlation coefficient, R . A p -value for the statistical test having null hypothesis $H_0 : R = 0$ (i.e., no correlation) is provided, along with a 95 % confidence interval for R .

3 Simulation Study

In order to evaluate the performance of the proposed SN approach for modeling the CCF, we begin by considering a simulation study using spectra generated from the Spot Oscillation And Planet

⁴Note that SN FWHM does not correspond to the width of the SN distribution at half maximum like in the Normal case.

(SOAP) 2.0 code (Dumusque et al. 2014).

[[**Umberto:** Add more information on SOAP, the way we retrieve the CCFs with SOAP, the characteristics of the CCF (i.e. how many points, ...) and the reasons why we need CCFs simulated with SOAP.]] [[**Umberto:** Specify characteristic of the star.]]

[[**Umberto:** Add info on the standard errors about the asymmetry parameter and the width and make a comparison with the results from the bootstrap. Then right discussions and conclusions,]]

3.1 Faculae

[[**Umberto:** Specify characteristic of the faculae.]]

Fig. 3 shows the results obtained when a faculae is present on the photosphere of the star. It is possible to note that, although there is not a planet, the presence of the faculae leads to spurious variations in RVs for all the proposed indicators. SN mean RV seems to have the smallest spurious variations caused by the faculae.

Since in this case we know that the variations in RV are only caused by the faculae, the evaluation of the correlations between the set of RVs and the asymmetry parameters can provide relevant information to understand the cause of the variations observed in RVs. Fig. 4 shows the correlations obtained using as RV respectively RV mean SN, RV median SN and RV. The γ parameter shows a correlation with RV median SN equal to -0.944 , which is stronger than the correlations between the other asymmetry indicators and their estimated RVs. The correlation between the width of the CCF and RV mean SN is the strongest one ($R^2 = 0.655$).

3.2 Spot

[[**Umberto:** Specify characteristic of the spot.]]

Fig. 5 shows the results obtained when a spot is present on the photosphere of the star. Similarly to the previous case, the presence of the spot leads to spurious variations in RVs for all the proposed indicators, but N mean RV seems to have the smallest ones. We note however that in this case all the three indicators are similarly effected by the spurious changes in RVs caused by the spot.

Fig. 6 shows the correlations between the asymmetry parameters and respectively RV mean SN, RV median SN and RV. The correlation between γ and RV median SN is equal to -0.89 while the correlation between γ and RV median SN is equal to -0.82 . Both these correlations are stronger than the correlation between the BIS SPAN and RV ($R^2 = -0.77$). The correlations between the width of the CCF and the corresponding RVs are slightly stronger when fitting a SN rather than a Normal distribution, as shown in the left series of plots in Fig. 6. The results shows anyway a correlation close to 0, suggesting that the presence of a spot on the photosphere of star changes the shape of the CCF rather than its width.

3.3 Spot and planet

[[**Umberto:** Specify characteristic of the spot.]]




Figure 3: (RVs changes as function of the orbital phase in the case in which a faculae is present on the photosphere of the star. SN mean RV seems to have the smallest spurious variations caused by the faculae.




Figure 4: Evaluation of the correlation between the RVs and the asymmetry parameters when a faculae is present on the photosphere of the star. In this case both the shape and the width of the CCF changes as the faculae moves, producing statistically significant correlations between the RVs and respectively the asymmetry parameter and the width parameter.




Figure 5: RVs changes as function of the orbital phase in the case in which a spot is present on the photosphere of the star. SN mean RV seems to have the smallest spurious variations caused by the faculae.




Figure 6: Evaluation of the correlation between the RVs and the asymmetry parameters when a spot is present on the photosphere of the star. In this case only the shape of the CCF changes as the spot moves, producing statistically significant correlations only between the RVs and the asymmetry parameter.

Fig. 7 shows the results obtained when a spot is present on the photosphere of the star and a planet is injected. The planet, having an amplitude of 10 m s^{-1} , produces a pure doppler shift on the CCF, without further changing its shape. In this case N mean RV seems to have the largest variations caused by the combined action of spot and planet.

Fig. 8 shows the correlations between γ and respectively RV mean SN and RV median SN and the correlation between BIS SPAN and RV. In this case the correlations are weaker than the ones derived when only a spot is present on the photosphere of the star. Anyway the γ parameter shows a correlation with the median SN of -0.433 , which is stronger than what the correlation between the other asymmetry indicators and their corresponding RVs. The correlations between the width of the CCF and the corresponding RVs are comparable and, similarly to the previous case, close to 0.




Figure 7: RVs changes as function of the orbital phase in the case in which a spot is present on the photosphere of the star and a planet is injected. N mean RV seems to have the largest variations caused by the combined action of spot and planet.

As final step we removed to the RVs obtained in this case those spurious signals coming from the spot. The results, presented in Fig. 9, show that the periodic pure doppler shift caused by the planet is detected.




Figure 8: Evaluation of the correlation between the RVs and the asymmetry parameters when a spot is present on the photosphere of the star and a planet is injected. In this case only the shape of the CCF changes as the spot moves, producing statistically significant correlations only between the RVs and the asymmetry parameter. The correlations between the RVs and the width parameter of the CCF is weaker than the previous case that considers only the presence of a spot on the photosphere of the star.




Figure 9: RVs changes as function of the orbital phase after removed the spurious effect of the spot. In green the real signal of the planet is displayed.

In this Section we presented a first implementation of the SN fit to CCF, using the simulation environment SOAP. Before moving to the next Section, where the analyses on five real stars are presented, we need to provide further considerations. First of all, looking at the analyses conducted with SOAP, it seems that the largest correlation between an asymmetry parameter and a set of RVs happens to be when respectively γ and RV median SN are used. This is a bit surprising, since as the shape of the CCF changes, we expect RV median SN to be more robust than RV mean SN. [\[\[Umberto: A possible justification of this ...\]\]](#). As second, when searching for stellar activity by deriving the correlation between the set of RVs and either an asymmetry parameter or the width of the CCF, the latter leads to weaker and hence less conclusive results if the active region is a spot. When stellar activity is dominated by faculae, both the shape and the width of the CCF changes as the faculae evolves on the photosphere of the star. Finally, the correlations involving the common indicators (i.e. RV, FWHM and BIS SPAN) are systematically weaker than the correlations obtained by fitting the SN to the CCF, suggesting that this distribution could be helpful when searching for active regions. We recall moreover that all the quantities needed for conducting the analyses of the CCF are directly available by just fitting the SN.

We remand the reader to Section 5 for the evaluation of the standard errors associated with the parameters estimated for all the three presented cases. This step will point out how RV median SN minimizes the uncertainties respect using both RV mean SN and RV, therefore suggesting its use in order to properly define the set of RVs of the star.

4 Real data applications

In this Section we present the analyses conducted on Alpha Centauri b, comparing the result of fitting a CCF using the SN distribution defined in Section 2.1 with the approach based on the Normal distribution. Other four stars have been analyzed with the proposed method and details can be found in the Appendix A. For all the stars that have been considered in the present work, we selected those CCFs having SNR larger than 10.

A comparison with the results obtained by the classic approach is done, where the RVs of the star are estimated by retrieving the mean of the Normal distribution used to fit the CCF, along with the BIS SPAN or the other asymmetric parameters defined in Figueira et al. (2013). The latter parameters are calculated separately from the Normal fit that leads to the set of RVs of the star.

4.1 Alpha Centauri B

A total of 1808 CCFs measured in 2010 have been analysed from the star Alpha Centauri B. Several measurement in 2010 are contaminated by light from Alpha Centauri A. To remove contaminated spectra and thus CCFs, we performed the same selection as presented in Dumusque et al. (2012). Moreover, as noted in Dumusque et al. (2012) and Thompson et al. (2017), this dataset presents a strong stellar activity signal.

We begin the analyses by evaluating the correlation between γ and the BIS SPAN. In Fig. 10, we see that the relationship between γ and the BIS SPAN is linear, with a slope equal to 0.72

and a strong Pearson correlation coefficient of 0.954. This comparison is useful because γ is an adimensional parameter taking information about the asymmetry of the SN while the BIS SPAN, beyond this, has got unit of measure of km s^{-1} . In other words, by using Fig. 10, it is possible to provide a physical meaning to γ .




Figure 10: Correlation between γ and the BIS SPAN for Alpha Centauri B. Because γ is adimensional, retrieving the slope between γ and the BIS SPAN, which is expressed in km s^{-1} , allows us to provide physical meaning to γ .

Fig. 11 shows the comparison between the RVs retrieved using the SN shape and the ones obtained with the Normal shape. It is possible to appreciate the presence of a strong stellar activity signal, as expected (Dumusque et al. 2012; Thompson et al. 2017). When using RV mean SN, it is possible to observe more variations than the ones measured by the Normal fitting. This happens because the mean of the SN is more sensitive to stellar activity. In fact, because the SN includes an asymmetry parameter, RV mean SN gets more shifted in the direction of the asymmetry induced by stellar activity. On the other hand, when using RV median SN, smaller variations in RV are caused by changes in the asymmetry of the CCF, because this second location parameter is a more robust indicator than the mean. The bottom plot of Fig. 11 captures this aspect. Both indicators can be used to capture and summarise the different information available in the CCF, as will be shown in

the remainder of this work.




Figure 11: (top) RVs as function of Julian Day for Alpha Centauri b. The RVs are retrieved using the mean of the Normal (red triangles), RV mean SN (black circles), RV median SN (cyan crosses). (bottom) RV differences between Normal RV and RV mean SN (black circles) and between Normal RV and a RV median SN (cyan crosses).

Similar to Figueira et al. (2013), we compare the correlation between the different activity indicators and the RVs of the star in Fig. 12. The correlation between γ and RV mean SN and the correlation between γ and RV median SN are much stronger than the correlations calculated between the other asymmetry parameters and their corresponding RVs. In particular the correlation between γ and RV mean SN is almost twice the correlation between the other asymmetry parameters and their corresponding RVs.

Because the median is a more robust index than the mean, the correlation between γ and RV median SN is not as large as the correlation between γ and RV mean SN, but it is nonetheless 1.5 times larger than the correlation between the other common asymmetry parameters and their corresponding RVs. In other words, changes in the asymmetry of the CCF are better captured when using the RV mean SN. The correlation between FWHM and the RVs, either by using RV mean SN or RV median SN, is as well stronger when fitting a SN distribution rather than a Normal. All

the correlations are statistically different from 0. Recalling the analyses presented in Section 3, we could infer that Alpha Centauri b is dominated by faculae, because the correlations between the RVs and the width of the CCF are strong (in particular the correlation between RV mean SN and SN FWHM is 0.817).




Figure 12: Correlation between the asymmetry parameters and the RVs for Alpha Centauri B. The last three plots show the correlation between the FWHM and the RVs for Alpha Centauri B using respectively the SN (RV mean SN and RV median SN) and the Normal fits. The correlations are always stronger when using parameters derived from the SN fit than the Normal one.

Both the proposed indicators coming from the SN distribution have desirable and undesirable behaviors: RV mean SN better catches changes in the asymmetry of the CCF but the resulting set of RVs ends up being contaminated by those spurious shifts caused by stellar activity that have been shortly presented in Section 1. When using RV median SN, the final set of radial velocity is less affected by those spurious shifts caused by stellar activity, but at the same time this indicator is not able to catch as well as RV mean SN changes in the shape and in the width of the CCF. Anyway, both RV mean SN and RV median SN are useful to catch different aspects of the CCF and our suggestion is to use RV mean SN when interested in retrieving information about changes in the shape and/or the width of the CCF. In order to provide a set of RVs containing the smallest

amount of spurious contamination imputable to stellar activity, our suggestion is to use instead RV median SN.

4.2 Doppler shift added to Alpha Centauri B

We also consider a real-data example using HARPS spectra from the star Alpha Centauri B with an imputed Doppler shift added...

[[Umberto: to be done]]

In the next Section we further motivate the reasons to define the RVs derived by the CCF by calculating RV median SN. In order to do that, we retrieve the standard errors associated with RV mean SN, RV median SN and RV.

5 Estimation of standard errors for the CCF parameters

In this section, we perform a bootstrap analysis to measure the standard errors associated to RV mean SN, RV median SN, RV, FWHM, SN FWHM, BIS SPAN and γ . Because a CCF is obtained from a cross-correlation, each point in a CCF is correlated with each other. Therefore, we cannot do a bootstrap analysis on perturbing independently each CCF point with a Gaussian distribution scaled to the error of each given point. To bypass this problem, we bootstrap a hundred times the stellar spectrum given the photon-noise error of each wavelength and calculate for each realization a new CCF. We then fit a Normal or a SN to each of these CCFs and then calculate the standard deviations of the distribution for the location parameters (RV, RV mean SN or RV median SN), the width parameters (FWHM or SN FWHM) and the parameters of asymmetry (BIS SPAN or γ).

[[Referee: It is not clear what you are doing here. I can understand bootstrapping a time series: you shuffle all the values keeping the time stamps fixed. How do you bootstrap a stellar spectrum? Do you shuffle the intensity values keeping the wavelengths fixed? That would produce a mess. Do you just add different levels of random noise? But the spectra already have noise in them, they are real observations. You need to state more clearly here what you did.]]

5.1 Estimation of standard errors for the CCF parameters for the simulation study

We start by calculating the standard errors of the parameters retrieved in Section 3, where with SOAP we produced CCFs contaminated by a faculae or a spot. In the third and final case we considered, beyond the spot, a planetary signal that produces pure doppler shifts i the CCFs.

Fig. 13 shows the results of the bootstrap analysis performed when a faculae is present on the photosphere of the star. The series of three plots in the top of Fig. 13 show the different errors for the RVs, defined as RV (red triangles), RV mean SN (black circles) or RV median SN (cyan crosses), the width and the asymmetry of the CCFs. In the three plots in the bottom of Fig. 13 we show the ratio between the parameters derived from the bootstrap analysis fitting the SN and the parameters derived from the bootstrap analysis fitting the Normal distribution. Concerning the standard errors related to the RVs, the ratio between the RV error measured by the bootstrap using

the SN and Normal fitting is 1.5 when using RV mean SN and 0.9 when using RV median SN. By using RV median SN we get standard errors 10% smaller than using the Normal fit and its corresponding mean. Regarding the errors in width of the CCF, we see that the bootstrap analysis for the Normal and the SN are comparable. Therefore, the precision in the width of the CCF is the comparable if we fit a Normal or a SN to the CCF. Finally, for the errors in evaluating the asymmetry of the CCF, we see that, when fitting the SN to the CCF, the asymmetry errors are 20% smaller. Therefore, the SN fit gives a better precision in CCF asymmetry than what can be reached using BIS SPAN.




Figure 13: Faculae Case. Comparison between the standard errors using the bootstrap analysis for the RVs, the FWHM and the asymmetry parameter. When using RV mean SN (black circles), the standard errors are in average 50% larger than the standard errors retrieved fitting a Normal (red triangles). However, if using RV median SN (cyan crosses), the standard errors are in average 10% smaller than the standard errors coming from the Normal fit. To use as asymmetry parameter γ of the SN leads to standard errors in average 20% smaller than the standard errors related to the BIS SPAN. **[[Umberto: explain what happens for those CCF 15 to 19 where s.e. decrease.]]** Note that for the asymmetry, the error in BIS SPAN is in km s^{-1} . To be able to compare the errors in γ and BIS SPAN, we multiplied the error in γ by the slope of the correlation between γ and BIS SPAN.

Fig. 14 shows the results of the bootstrap analysis performed when a spot is present on the photosphere of the star. The series of plots follows the specifications outlined for the previous case. Concerning the standard errors related to the RVs, the ratio between the RV error measured by the bootstrap using the SN and Normal fitting is 1.4 when using RV mean SN and 0.9 when using RV median SN. Regarding the errors in width of the CCF, we see that the bootstrap analysis for the Normal and the SN are comparable. Therefore, the precision in the width of the CCF is the comparable if we fit a Normal or a SN to the CCF. Finally, for the errors in evaluating the asymmetry of the CCF, we see that, when fitting the SN to the CCF, the asymmetry errors are 20% smaller.




Figure 14: Spot case. Comparison between the standard errors using the bootstrap analysis for the RVs, the FWHM and the asymmetry parameter. When using RV mean SN (black circles), the standard errors are in average 40% larger than the standard errors retrieved fitting a Normal (red triangles). However, if using RV median SN (cyan crosses), the standard errors are in average 10% smaller than the standard errors coming from the Normal fit. To use as asymmetry parameter γ of the SN leads to standard errors in average 20% smaller than the standard errors related to the BIS SPAN. Note that for the asymmetry, the error in BIS SPAN is in km s^{-1} . To be able to compare the errors in γ and BIS SPAN, we multiplied the error in γ by the slope of the correlation between γ and BIS SPAN.

Fig. 15 shows the results of the bootstrap analysis performed when a spot is present on the photosphere of the star. The series of plots follows the specifications outlined for the previous two cases. The conclusions are comparable to the case in which only a spot is present on the photosphere of the star. The ratio between the RV error measured by the bootstrap using the SN and Normal fitting is 1.4 when using RV mean SN and 0.9 when using RV median SN. The errors in width of the CCF are comparable and the errors in evaluating the asymmetry of the CCF are 20% smaller when using the asymmetry parameter γ of the SN.




Figure 15: Spot and Planet case. Comparison between the standard errors using the bootstrap analysis for the RVs, the FWHM and the asymmetry parameter. When using RV mean SN (black circles), the standard errors are in average 50% larger than the standard errors retrieved fitting a Normal (red triangles). However, if using RV median SN (cyan crosses), the standard errors are in average 10% smaller than the standard errors coming from the Normal fit. To use as asymmetry parameter γ of the SN leads to standard errors in average 20% smaller than the standard errors related to the BIS SPAN. Note that for the asymmetry, the error in BIS SPAN is in km s^{-1} . To be able to compare the errors in γ and BIS SPAN, we multiplied the error in γ by the slope of the correlation between γ and BIS SPAN.

5.2 Estimation of standard errors for the CCF parameters for real stars

In the top plots of Fig. 16 we show the different errors for the RVs, either defined as RV (red triangles), RV mean SN (black circles) or RV median SN (cyan crosses), the width and the asymmetry of the CCFs for three stars, HD215152, HD192310 and Corot-7, that are all at different SNR levels. The parameter SN50 corresponds to the SNR in order 50, which defines a wavelength of 550 nm. In the bottom plots, we show the ratio between the parameters derived from the bootstrap analysis fitting the SN and the parameters derived from the bootstrap analysis fitting the Normal distribution. We first see that the errors on the CCF parameters only depend on the SNR and do not depend on the spectral type. This is true if the spectral types are not too different though, like here where we show the results for G and K dwarfs.

Concerning the standard errors related to the RVs, the ratio between the RV error measured by the bootstrap using the SN and Normal fitting is 1.6 when using RV mean SN and 0.9 when using RV median SN. In other words, by using RV median SN as parameter that defines the radial velocity of the star given a CCF, we get standard errors 10% smaller than using the Normal fit and its corresponding mean. This result is consistent with what we observed with the simulation from SOAP presented in Section 3.

Regarding the errors in width of the CCF, we see that the bootstrap analysis for the Normal and the SN are comparable. Therefore, the precision in the width of the CCF is comparable if we fit a Normal or a SN to the CCF.

Finally, for the errors in evaluating the asymmetry of the CCF, we see that, when fitting the SN to the CCF, the asymmetry errors are 15% smaller. Therefore, the SN fit gives a better precision in CCF asymmetry than what can be reached using BIS SPAN. We recall moreover that, using the SN, all parameters are automatically retrieved in 1 single step, while in the common approach the RV and the BIS SPAN are calculated separately.

6 Discussion

An analysis of the CCF residuals after fitting a Normal or SN distribution shows that the SN is a slightly better model to explain the shape of the CCF. This comes from the fact that CCFs present a natural asymmetry due to the convective blueshift.

We tested at first our assumptions by using simulated CCFs retrieved using the software SOAP 2.0. We then compared for five real stars the difference between the RVs (defined as mean of a Normal, mean of the SN or median of the SN), FWHM and asymmetry (BIS SPAN in the Normal case and γ in the SN case). The γ parameter is linearly dependent on the BIS SPAN, with always a strong correlation coefficient. The slope of this linear correlation changes depending on the studied star. This is probably because the spectral type is different, therefore the effects from stellar activity are different.

When using as parameter for the RV the mean of the SN, the standard errors are in average 60% larger than the standard errors retrieved fitting a Normal. However, once the RV is defined as the median of the SN (cyan crosses), the standard errors are in average 10% smaller than the standard




Figure 16: Comparison between the standard errors using the bootstrap analysis for the RVs, the FWHM and the asymmetry parameter. When using RV mean SN (black circles), the standard errors are in average 60% larger than the standard errors retrieved fitting a Normal (red triangles). However, if using RV median SN (cyan crosses), the standard errors are in average 10% smaller than the standard errors coming from the Normal fit. To use as asymmetry parameter γ of the SN leads to standard errors in average 15% smaller than the standard errors related to the BIS SPAN. Note that for the asymmetry, the error in BIS SPAN is in km s^{-1} . To be able to compare the errors in γ and BIS SPAN, we multiplied the error in γ by the slope of the correlation between γ and BIS SPAN.

errors coming from the Normal fit. When looking at the correlation between the asymmetry and width parameters of the CCF (FWHM and BIS SPAN or the alternative indicators in Figueira et al. (2013) in the Normal case, and SN FWHM and γ in the SN case) with respect to the RVs (RVs in the Normal case or SN RVs in the SN case), we observe that the correlations are always stronger for the parameters of the SN. Therefore, the SN parameters are more sensitive to activity. In the case of Tau Ceti, which is at very low activity level, we find a significant correlation of 0.322 between γ and RV mean SN, while for all the other asymmetric parameterization, BIS SPAN or the alternative indicators in Figueira et al. (2013), the correlations are weaker with a maximum of 0.225.

7 Conclusion

In this paper we introduced a novel approach based on the Skew Normal (SN) distribution for deriving RVs and shape variations in the CCF of stars. When searching for small-mass exoplanets using the RV technique, it is essential to understand the shape variation of the CCF, which is a proxy for stellar activity effects. The standard approach consist at first to adjust a Normal distribution to the CCF to get the RV and FWHM, defined as the mean and the FWHM of the Normal distribution, and then to measure the asymmetry by calculating BIS SPAN. FWHM and BIS SPAN give us information on the line shape that are used to probe stellar activity signals.

In this paper we propose to conduct the analysis fitting a SN distribution to the CCF. Since the CCF presents a natural asymmetry due the convective blueshift, the SN distribution can better catch these aspects respect the Normal fit. On top of that, by using the SN distribution to fit CCFs, we can measure simultaneously the RV of the star, the width and the asymmetry of the CCF.

Starting from the simulation environment SOAP and then moving to real stars, we showed that using the SN distribution to fit CCFs brings a significant improvement in probing stellar activity. While for the Normal distribution mean and median are equivalent, using the SN fit different location parameters can be tested. While the median of the SN is more robust respect variations in the shape of the CCF, the mean of the SN is more sensible to changes in the asymmetry of the CCF. We suggest to use as parameter that defines the RV of the star the median of the SN, since the standard errors related to this parameter are 10% smaller than the standard errors retrieved using the Normal distribution. For evaluating changes in the asymmetry of the CCF, we suggest to use the mean of the SN. The correlations between RV mean SN and SN FWHM, and RV mean SN and γ (the asymmetry parameter of the SN) are much stronger than the correlations between the equivalent parameters derived using a Normal fit (RV, FWHM and BIS SPAN or the asymmetric parameters described in Figueira et al. (2013)). The precision on the asymmetry measured by γ is greater than the one on BIS SPAN by $\sim 15\%$. Therefore when searching for rotational periods in the data, or applying Gaussian Processes to account for stellar activity signals, the SN parameters should be used.

Finally, we also encourage the use of bootstrapping to estimate more realistic errors on the different parameters of the Normal or SN fitted to the CCF, mainly in the low SNR regime where a gain of 50% can be reached. This takes significantly more time, but note that 100 realization are

Star	# CCFs	R(SN γ , Bis-Span)	slope(SN γ , Bis-Span)	R(SN γ , RV mean SN)	R(Bis-Span, RV)	R(FIG BiGaussian, RV)	R(SN FWHM, RV mean SN)	R(FWHM, RV)
HD192310	1577	0.888	0.786	0.669(0.64; 0.695)	0.329(0.285; 0.373)	-0.333(-0.376; -0.289)	0.666(0.637; 0.692)	0.476(0.43670.514)
HD10700	7928	0.78	0.604	0.322(0.302; 0.342)	-0.073(-0.095; -0.0051)	0.127(0.105; 0.148)	0.421(0.403; 0.439)	0.529(0.513; 0.545)
HD215152	273	0.763	0.794	0.571(0.485; 0.646)	-0.067(-0.184; 0.052)	0.269(0.155; 0.376)	0.210(0.094; 0.321)	-0.138(-0.253; -0.020)
Corot 7	173	0.814	0.607	0.561(0.450; 0.656)	0.092(-0.058; 0.238)	-0.335(-0.228; -0.082)	-0.709(0.626; 0.776)	0.595(0.489; 0.683)

Table 2: Subset of notable correlations between the asymmetry parameter (and the FWHM) and the RVs for four stars: HD192310, HD10700, HD215152 and Corot 7.

enough to get a good estimation of errors.

8 Acknowledgements

We are grateful to all technical and scientific collaborators of the HARPS Consortium, ESO Headquarters and ESO La Silla who have contributed with their extraordinary passion and valuable work to the success of the HARPS project. XD is grateful to the Society in Science–The Branco Weiss Fellowship for its financial support.

A Appendix

In this Appendix we present the analyses conducted on other 4 stars: HD192310, HD10700, HD215152 and finally Corot-7.

[[Umberto: Add further information about the stars here presented.]]

Table 2 summarizes the results obtained by the SN fit and the some of the results based on the Normal fit. The results are all consistent with the conclusions derived by the analyses on Alpha Centauri b. The correlation between γ and RV mean SN is stronger than the correlation between the BIS SPAN and RV for all the considered stars. The correlation between SN FWHM and RV mean SN is stronger than the correlation between FWHM and RV for three of the four stars. The analyses of the correlations for the four stars are presented Fig. 17, 18, 19 and 20.

References

- G. Anglada-Escudé and R. P. Butler. The harps-terra project. i. description of the algorithms, performance, and new measurements on a few remarkable stars observed by harps. *The Astrophysical Journal Supplement Series*, 200(2):15, 2012.
- A. Azzalini. A class of distributions which includes the normal ones. *Scandinavian journal of statistics*, pages 171–178, 1985.
- A. Azzalini and A. Capitanio. The skew-normal and related families. institute of mathematical statistics monographs, 2014.
- A. Baranne, D. Queloz, M. Mayor, G. Adrianzyk, G. Knispel, D. Kohler, D. Lacroix, J.-P. Meunier, G. Rimbaud, and A. Vin. Elodie: A spectrograph for accurate radial velocity measurements. *Astronomy and Astrophysics Supplement Series*, 119(2):373–390, 1996.




Figure 17: Correlation between the asymmetry parameters and the RVs for HD 192310. The last three plots show the correlation between the FWHM and the RVs for HD 192310 using respectively the SN and the Normal fits. [[Umberto: faculae??]]




Figure 18: Correlation between the asymmetry parameters and the RVs for Tau Ceti. The last three plots show the correlation between the FWHM and the RVs for Tau Ceti using respectively the SN and the Normal fits. [\[\[Umberto: faculae??\]\]](#)




Figure 19: Correlation between the asymmetry parameters and the RVs for HD 215152. The last three plots show the correlation between the FWHM and the RVs for HD 215152 using respectively the SN and the Normal fits. [[Umberto: spot???]]




Figure 20: Correlation between the asymmetry parameters and the RVs for Corot-7. The last three plots show the correlation between the FWHM and the RVs for Corot-7 using respectively the SN and the Normal fits. [\[\[Umberto: faculae??\]\]](#)

- I. Boisse, C. Moutou, A. Vidal-Madjar, F. Bouchy, F. Pont, G. Hébrard, X. Bonfils, B. Croll, X. Delfosse, M. Desort, T. Forveille, A. Lagrange, B. Loeillet, C. Lovis, J. M. Matthews, M. Mayor, F. Pepe, C. Perrier, D. Queloz, J. F. Rowe, N. C. Santos, D. Ségransan, and S. Udry. Stellar activity of planetary host star HD 189 733. *A&A*, 495:959–966, Mar. 2009. doi: 10.1051/0004-6361/200810648.
- I. Boisse, F. Bouchy, G. Hébrard, X. Bonfils, N. Santos, and S. Vauclair. Disentangling between stellar activity and planetary signals. *A&A*, 528:A4, Apr. 2011. doi: 10.1051/0004-6361/201014354.
- M. Desort, A.-M. Lagrange, F. Galland, S. Udry, and M. Mayor. Search for exoplanets with the radial-velocity technique: quantitative diagnostics of stellar activity. *Astronomy & Astrophysics*, 473(3):983–993, 2007.
- D. Dravins, L. Lindegren, and Å. Nordlund. Solar granulation-influence of convection on spectral line asymmetries and wavelength shifts. *Astronomy and Astrophysics*, 96:345–364, 1981.
- X. Dumusque. Radial velocity fitting challenge. I. Simulating the data set including realistic stellar radial-velocity signals. *A&A*, 593:A5, Aug. 2016. doi: 10.1051/0004-6361/201628672.
- X. Dumusque, S. Udry, C. Lovis, N. C. Santos, and M. J. P. F. G. Monteiro. Planetary detection limits taking into account stellar noise. I. Observational strategies to reduce stellar oscillation and granulation effects. *A&A*, 525:A140, Jan. 2011. doi: 10.1051/0004-6361/201014097.
- X. Dumusque, F. Pepe, C. Lovis, D. Ségransan, J. Sahlmann, W. Benz, F. Bouchy, M. Mayor, D. Queloz, N. Santos, et al. An earth-mass planet orbiting [agr] centauri b. *Nature*, 491(7423):207–211, 2012.
- X. Dumusque, I. Boisse, and N. Santos. Soap 2.0: A tool to estimate the photometric and radial velocity variations induced by stellar spots and plagues. *The Astrophysical Journal*, 796(2):132, 2014.
- X. Dumusque, F. Borsa, M. Damasso, R. F. Díaz, P. C. Gregory, N. C. Hara, A. Hatzes, V. Rajpaul, M. Tuomi, S. Aigrain, G. Anglada-Escudé, A. S. Bonomo, G. Boué, F. Dauvergne, G. Frustagli, P. Giacobbe, R. D. Haywood, H. R. A. Jones, J. Laskar, M. Pinamonti, E. Poretti, M. Rainer, D. Ségransan, A. Sozzetti, and S. Udry. Radial-velocity fitting challenge. II. First results of the analysis of the data set. *A&A*, 598:A133, Feb. 2017. doi: 10.1051/0004-6361/201628671.
- F. Feng, M. Tuomi, and H. R. A. Jones. Evidence for at least three planet candidates orbiting HD20794. *ArXiv e-prints*, May 2017.
- P. Figueira, N. Santos, F. Pepe, C. Lovis, and N. Nardetto. Line-profile variations in radial-velocity measurements-two alternative indicators for planetary searches. *Astronomy & Astrophysics*, 557:A93, 2013.
- D. Fischer, C. Jurgenson, T. McCracken, D. Sawyer, R. Blackman, and A. E. Szymkowiak. EXPRES: the EXtreme PREcision Spectrograph at the Discovery Channel Telescope. In *American Astronomical Society Meeting Abstracts #229*, volume 229 of *American Astronomical Society Meeting Abstracts*, page 126.04, Jan. 2017.
- D. F. Gray. The third signature of stellar granulation. *The Astrophysical Journal*, 697(2):1032, 2009.
- A. P. Hatzes. Starspots and exoplanets. *Astronomische Nachrichten*, 323(3-4):392–394, 2002.

- R. Haywood, A. Collier Cameron, D. Queloz, S. Barros, M. Deleuil, R. Fares, M. Gillon, A. Lanza, C. Lovis, C. Moutou, et al. Planets and stellar activity: hide and seek in the corot-7 system? *Monthly notices of the royal astronomical society*, 443(3):2517–2531, 2014.
- M. Kurster, M. Endl, F. Rouesnel, S. Els, A. Kaufer, S. Brillant, A. Hatzes, S. Saar, and W. Cochran. The low-level radial velocity variability in barnard’s star (= gj 699). secular acceleration, indications for convective redshift, and planet mass limits. *ASTRONOMY AND ASTROPHYSICS-BERLIN*-, 403(3): 1077–1088, 2003.
- A.-M. Lagrange, M. Desort, and N. Meunier. Using the sun to estimate earth-like planets detection capabilities-i. impact of cold spots. *Astronomy & Astrophysics*, 512:A38, 2010.
- L. Lindegren and D. Dravins. The fundamental definition of ?radial velocity? *Astronomy & Astrophysics*, 401(3):1185–1201, 2003.
- N. Meunier, M. Desort, and A.-M. Lagrange. Using the sun to estimate earth-like planets detection capabilities-ii. impact of plages. *Astronomy & Astrophysics*, 512:A39, 2010.
- F. Pepe, M. Mayor, F. Galland, D. Naef, D. Queloz, N. Santos, S. Udry, and M. Burnet. The coralie survey for southern extra-solar planets vii-two short-period saturnian companions to hd 108147 and hd 168746. *Astronomy & Astrophysics*, 388(2):632–638, 2002.
- F. Pepe, P. Molaro, S. Cristiani, R. Rebolo, N. C. Santos, H. Dekker, D. Mégevand, F. M. Zerbi, A. Cabral, P. Di Marcantonio, M. Abreu, M. Affolter, M. Aliverti, C. Allende Prieto, M. Amate, G. Avila, V. Baldini, P. Bristow, C. Broeg, R. Cirami, J. Coelho, P. Conconi, I. Coretti, G. Cupani, V. D’Odorico, V. De Caprio, B. Delabre, R. Dorn, P. Figueira, A. Frago, S. Galeotta, L. Genolet, R. Gomes, J. I. González Hernández, I. Hughes, O. Iwert, F. Kerber, M. Landoni, J.-L. Lizon, C. Lovis, C. Maire, M. Mannetta, C. Martins, M. Monteiro, A. Oliveira, E. Poretti, J. L. Rasilla, M. Riva, S. Santana Tschudi, P. Santos, D. Sosnowska, S. Sousa, P. Spanó, F. Tenegi, G. Toso, E. Vanzella, M. Viel, and M. R. Zapatero Osorio. ESPRESSO: The next European exoplanet hunter. *ArXiv e-prints*, Jan. 2014.
- D. Queloz, G. Henry, J. Sivan, S. Baliunas, J. Beuzit, R. Donahue, M. Mayor, D. Naef, C. Perrier, and S. Udry. No planet for hd 166435. *Astronomy & Astrophysics*, 379(1):279–287, 2001.
- D. Queloz, F. Bouchy, C. Moutou, A. Hatzes, G. Hébrard, R. Alonso, M. Auvergne, A. Baglin, M. Barbieri, P. Barge, W. Benz, P. Bordé, H. J. Deeg, M. Deleuil, R. Dvorak, A. Erikson, S. Ferraz Mello, M. Fridlund, D. Gandolfi, M. Gillon, E. Guenther, T. Guillot, L. Jorda, M. Hartmann, H. Lammer, A. Léger, A. Llebaria, C. Lovis, P. Magain, M. Mayor, T. Mazeh, M. Ollivier, M. Pätzold, F. Pepe, H. Rauer, D. Rouan, J. Schneider, D. Segransan, S. Udry, and G. Wuchterl. The CoRoT-7 planetary system: two orbiting super-Earths. *A&A*, 506:303–319, Oct. 2009. doi: 10.1051/0004-6361/200913096.
- V. Rajpaul, S. Aigrain, M. A. Osborne, S. Reece, and S. Roberts. A Gaussian process framework for modelling stellar activity signals in radial velocity data. *MNRAS*, 452:2269–2291, Sept. 2015. doi: 10.1093/mnras/stv1428.
- P. Robertson, S. Mahadevan, M. Endl, and A. Roy. Stellar activity masquerading as planets in the habitable zone of the M dwarf Gliese 581. *Science*, 345:440–444, July 2014. doi: 10.1126/science.1253253.
- S. H. Saar and R. A. Donahue. Activity-related radial velocity variation in cool stars. *The Astrophysical Journal*, 485(1):319, 1997.

- A. Thompson, C. Watson, E. de Mooij, and D. Jess. The changing face of α centauri b: probing plage and stellar activity in k dwarfs. *Monthly Notices of the Royal Astronomical Society: Letters*, 468(1):L16–L20, 2017.