1. (Proof of Theorem 3.6.2,, continuous case) Let X and Y be any two random variables with joint pdf $f_{X,Y}(x,y)$ and let a and b be any two constants. Show that E(aX+bY)=aE(X)+bE(Y).

$$E(g(x)) = \int g(x) \cdot f(x) \cdot f(x) dx dy$$

$$E(ax + b+r) = \int (ax + b+r) \cdot f(x) \cdot f(x) \cdot f(x) dx dy$$

$$= a \int x \int f(x) \cdot f(x) dy dx dx + b \cdot \int y \cdot f(x) dx dy$$

$$= a \int x \int f(x) dx dx + b \cdot \int y \int f(x) dx dy$$

$$= a \int x f(x) + b \cdot E(x)$$

4. Show that

$$Cov(aX + b, cY + d) = ac Cov(X, Y)$$

for any constants a, b, c, d.

$$COV(aX+b, CY+d)$$

$$= \mathbb{E}\left(aX+b \cdot (CY+d)\right) - \mathbb{E}\left(aX+b \cdot E(CY+d)\right)$$

$$= \mathbb{E}\left(a \cdot C \cdot X \cdot Y + a \cdot d \cdot X + b \cdot c \cdot Y + b \cdot d\right) - \left(a \cdot E(X) \cdot E(Y) + a \cdot d\right)$$

$$= a \cdot C \cdot E(XY) + a \cdot d \cdot E(X) + b \cdot C \cdot E(Y) + b \cdot d - a \cdot C \cdot E(X) \cdot E(Y) - a \cdot d$$

$$= a \cdot C \cdot \left(E(XY) - E(X) \cdot E(Y)\right)$$

$$= a \cdot C \cdot \left(E(XY) - E(X) \cdot E(Y)\right)$$

$$= a \cdot C \cdot \left(E(XY) - E(X) \cdot E(Y)\right)$$

5. Suppose the length of time, in minutes, that you have to wait at a bank teller's window is uniformly distributed over the interval (0, 10). If you go to the bank four times during the next month, what is the probability that your second longest wait will be less than five minutes?

9. Let X_1, X_2, \ldots, X_n be independent and identically distributed random variables having expected value μ and variance σ^2 , and let

$$S^{2} = \sum_{i=1}^{n} \frac{(X_{i} - \overline{X})^{2}}{n-1}$$

be the sample variance, where $\overline{X} = \frac{1}{n} \sum_{i=1}^{n} X_i$. Find the $E(S^2)$.

$$S^{2} = \sum_{i=1}^{n} \frac{(x_{i} - \bar{x})^{2}}{n - 1} \qquad (n - 1) \cdot S^{2} = \sum_{i=1}^{n} (x_{i} - \bar{x})^{2}$$

$$= \sum_{i=1}^{n} (x_{i} - \bar{x})^{2} = \sum_{i=1}^{n} (x_{i} - x_{i})^{2} + (x_{i} - x_{i})^{2} + 2 \cdot (x_{i} - x_{i}) \cdot (x_{i} - \bar{x})$$

$$= \sum_{i=1}^{n} (x_{i} - x_{i})^{2} + (x_{i} - x_{i})^{2} + 2 \cdot (x_{i} - x_{i}) \cdot (x_{i} - x_{i})$$

$$= \sum_{i=1}^{n} (x_{i} - x_{i})^{2} + n \cdot (x_{i} - x_{i})^{2} + 2 \cdot (x_{i} - x_{i}) \cdot (x_{i} - x_{i})$$

$$= \sum_{i=1}^{n} (x_{i} - x_{i})^{2} + n \cdot (x_{i} - x_{i})^{2} + 2 \cdot (x_{i} - x_{i}) \cdot (x_{i} - x_{i})$$

$$\sum_{i=1}^{n} (x_i - M) = \sum_{i=1}^{n} x_i - n \cdot M$$

$$= n \cdot \sum_{i=1}^{n} x_i - n \cdot M$$

$$= n \cdot \sum_{i=1}^{n} x_i - n \cdot M$$

$$= n \cdot \sum_{i=1}^{n} x_i - n \cdot M$$

$$= n \cdot \sum_{i=1}^{n} (x_i - M)^2 - n \cdot (M - X)^2$$

$$= \sum_{i=1}^{n} (x_i - M)^2 - n \cdot (M - X)^2$$

$$\mathbb{E}\left(\frac{1}{1-1}(x_{1}-\overline{x})^{2}\right) = \mathbb{E}\left((n-1)\cdot S^{2}\right)$$

$$= \mathbb{E}\left(\frac{n}{1-1}(x_{1}-x_{1})^{2} - n\cdot (M-\overline{x})^{2}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left((x_{1}-x_{1})^{2}\right) - n\cdot \mathbb{E}\left((\overline{x}-x_{1})^{2}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left((x_{1}-x_{1})^{2}\right) - n\cdot \mathbb{E}\left((\overline{x}-x_{1})^{2}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left((x_{1}-x_{1})^{2}\right) - n\cdot \mathbb{E}\left((\overline{x}-x_{1})^{2}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left(x_{1}-x_{1}\right) - n\cdot \mathbb{E}\left((\overline{x}-x_{1})^{2}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left(x_{1}-x_{1}\right)$$

$$= \frac{n}{1-1}\mathbb{E}\left($$

$$\overline{X} = \overline{h} \cdot \Sigma X_{i}$$

$$= \overline{h} \cdot \overline{\Sigma} \cdot \overline{K}(X_{i})$$

$$= \overline{h} \cdot \overline{N} \cdot M$$

$$= M.$$

$$= M.$$

$$Var(\overline{X}) = Var(\overline{h} \Sigma X_{i})$$

$$= (\overline{h})^{2} \cdot Var(\Sigma X_{i})$$

$$= (\overline{h})^{2} \cdot \Sigma Var(X_{i})$$

$$= \frac{1}{n^{2}} \cdot n \cdot 6^{2}$$

$$= 6^{2}/n$$

 $-2n(M-X)^2$