

GSE98923 co-expression analysis

CEMiTool baseline - 54-sample balanced design

2026-09-21

Contents

1	Scope	2
2	Experimental design	3
3	Expression data	3
4	CEMiTool analysis	3
5	Diagnostic and analysis figures	4
5.1	beta_r2	4
5.2	gsea	4
5.3	hist	4
5.4	mean_k	5
5.5	mean_var	5
5.6	profile_M1	5
5.7	profile_M10	5
5.8	profile_M2	5
5.9	profile_M3	5
5.10	profile_M4	5
5.11	profile_M5	5
5.12	profile_M6	5
5.13	profile_M7	5
5.14	profile_M8	5
5.15	profile_M9	6
5.16	profile_Not_Correlated	6
5.17	qq	6

6	Exported result tables	6
6.1	beta_fit_indices.tsv	6
6.2	enrichment_es.tsv	7
6.3	enrichment_nes.tsv	7
6.4	enrichment_padj.tsv	8
6.5	module.tsv	8
6.6	parameters.tsv	9
6.7	selected_genes.txt	9
6.8	summary_eigengene.tsv	11
6.9	summary_mean.tsv	11
6.10	summary_median.tsv	12
7	Supplementary data inventory	12
8	Reproducibility	13
9	Notes for manuscript use	15
10	Post-CEMiTool module statistics	15
10.1	Network calibration and sensitivity	15
10.2	Module eigengene model	15
10.3	FDR overview by module	16
10.4	Significant Cultivar x Stage interactions	16
10.5	Cabernet Sauvignon vs Pinot noir within stage	16
10.6	Year effect	17
10.7	Interpretation boundaries and next step	17

1 Scope

This report documents the computational analysis of the balanced 54-sample subset of GEO accession **GSE98923** used in the CEMiTool baseline workflow.

The report is intended as an analysis record and potential supplementary document for a scientific manuscript. Large machine-readable matrices are not printed in full in the document; complete files are preserved separately and indexed in the supplementary manifest.

2 Experimental design

The analysis contains **54 samples**, covering two cultivars, three developmental stages, three years, and three biological replicates per cultivar-stage-year combination.

Table 1: Sample counts by cultivar, developmental stage, and year.

Cultivar	Stage	Year	Freq
Cabernet Sauvignon	FruitSet	2012	3
Pinot noir	FruitSet	2012	3
Cabernet Sauvignon	Harvest	2012	3
Pinot noir	Harvest	2012	3
Cabernet Sauvignon	Veraison	2012	3
Pinot noir	Veraison	2012	3
Cabernet Sauvignon	FruitSet	2013	3
Pinot noir	FruitSet	2013	3
Cabernet Sauvignon	Harvest	2013	3
Pinot noir	Harvest	2013	3
Cabernet Sauvignon	Veraison	2013	3
Pinot noir	Veraison	2013	3
Cabernet Sauvignon	FruitSet	2014	3
Pinot noir	FruitSet	2014	3
Cabernet Sauvignon	Harvest	2014	3
Pinot noir	Harvest	2014	3
Cabernet Sauvignon	Veraison	2014	3
Pinot noir	Veraison	2014	3

3 Expression data

The processed matrix used for CEMiTool contains **29971 genes** and **54 samples**.

Expression values were derived from the processed RPKM matrix distributed with GSE98923 and transformed as:

$$\log_2(\text{RPKM} + 1)$$

The complete matrices are available under:

- data/processed/expression_rpkм.tsv
- data/processed/expression_log2rpkм.tsv

4 CEMiTool analysis

CEMiTool identified **11 modules** in the baseline analysis.

The analysis used Pearson correlation, an unsigned network, a signed TOM, a minimum module size of 30, and a fixed random seed of 1234.

Automatic beta selection did not identify a usable value in the initial baseline run. To reproduce the established 54-sample baseline, the soft-thresholding power was fixed explicitly at **beta = 10** using **set_beta = 10** and **force_beta = FALSE**. This value is specific to the present expression matrix and should be re-evaluated if the sample set, normalization, or filtering changes. The recorded beta for this run is **10**.

Table 2: Recorded analysis parameters.

parameter	value
dataset	GSE98923
samples	54
transform	log2(RPKM+1)
correlation	pearson
network_type	unsigned
tom_type	signed
beta_strategy	manual_set_beta_beta10
set_beta	10
force_beta	FALSE
beta	10
r2	0.706374247063269
filter	TRUE
filter_pval	0.1
min_module_size	30
seed	1234
modules	11

5 Diagnostic and analysis figures

All figures generated by the analysis are stored in **results/figures/**. PDF figures are converted to PNG copies for reliable inclusion in Word and HTML reports.

5.1 beta_r2

[1] “assets/figures/beta_r2.png” attr(,“class”) [1] “knit_image_paths” “knit_asis”

5.2 gsea

[1] “assets/figures/gsea.png” attr(,“class”) [1] “knit_image_paths” “knit_asis”

5.3 hist

[1] “assets/figures/hist.png” attr(,“class”) [1] “knit_image_paths” “knit_asis”

5.4 mean_k

[1] “assets/figures/mean_k.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.5 mean_var

[1] “assets/figures/mean_var.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.6 profile_M1

[1] “assets/figures/profile_M1.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.7 profile_M10

[1] “assets/figures/profile_M10.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.8 profile_M2

[1] “assets/figures/profile_M2.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.9 profile_M3

[1] “assets/figures/profile_M3.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.10 profile_M4

[1] “assets/figures/profile_M4.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.11 profile_M5

[1] “assets/figures/profile_M5.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.12 profile_M6

[1] “assets/figures/profile_M6.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.13 profile_M7

[1] “assets/figures/profile_M7.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.14 profile_M8

[1] “assets/figures/profile_M8.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.15 profile_M9

[1] “assets/figures/profile_M9.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.16 profile_Not_Correlated

[1] “assets/figures/profile_Not_Correlated.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

5.17 qq

[1] “assets/figures/qq.png” attr(“class”) [1] “knit_image_paths” “knit_asis”

6 Exported result tables

The following section provides a compact view of every tabular result exported by CEMiTool. Complete tables remain available as machine-readable files under **results/tables/**.

6.1 beta_fit_indices.tsv

Dimensions: 15 rows x 10 columns.

Table 3: beta_fit_indices.tsv

	Power	SFT.R.sq	slope	truncated.R.sq	mean.k.	median.k.	max.k.	Density	Centralization	Heterogeneity
1	0.8394009	0.5707687	0.7946939	1561.857117	54.661221	111.98010	51225230	1.805457	0.3207976	
2	0.4232012	0.3873307	0.5948254	1055.584872	206.787817	105.72110	34620690	2.133693	0.4790910	
3	0.0123210	0.0444688	0.7254709	792.1069588	4.9284514	53.85820	25979240	2.171812	0.5801171	
4	0.0875404	-	0.8310439	627.739736	75.7017512	74.27600	20588380	2.121878	0.6544427	
		0.1174076								
5	0.2744513	-	0.8129080	514.611285	28.4269011	36.65840	16878030	3.2041506	0.7137761	
		0.2192085								
6	0.4333524	-	0.7362054	431.828944	21.5085410	26.44510	14162970	3.1951480	0.7635877	
		0.3044837								
7	0.5495793	-	0.6942176	368.657523	39.8713593	5.47990	12091100	3.1860264	0.8068294	
		0.3627214								
8	0.6217077	-	0.6648029	318.950732	77.028728	58.72380	10460830	3.1771490	0.8452677	
		0.4267637								
9	0.6650263	-	0.6414873	278.909912	28.173737	92.84740	0.091475	90.1686699	0.8800415	
		0.4781301								
10	0.7063742	-	0.6502824	246.049851	89.373377	35.54000	0.080698	50.1606465	0.9119266	
		0.5251829								
12	0.7067470	-	0.6230659	195.571181	34.480706	40.39660	0.064142	70.1459880	0.9690884	
		0.6147726								
14	0.7472853	-	0.6797739	158.919239	7.11593	564.34660	0.052121	80.1330578	1.0196729	
		0.6734742								

Power	SFT.R.sq	slope	truncated.R.sq	mean.k.	median.k.	max.k.	Density	Centralization	Heterogeneity
16	0.7496827	-	0.6990696	131.3678072	38920	502.0230	0.0430855	0.1216459	1.0654044
	0.7344417								
18	0.7495778	-	0.7234977	110.0992953	69994	449.9688	0.0361100	0.1115423	1.1073859
	0.7871557								
20	0.7605758	-	0.7500230	93.33028	40.81002	405.8351	0.0306101	0.1025615	1.1463678
	0.8268108								

6.2 enrichment_es.tsv

Dimensions: 10 rows x 7 columns.

Table 4: enrichment_es.tsv

pathway	CS_FruitSet	CS_Veraison	CS_Harvest	PN_FruitSet	PN_Veraison	PN_Harvest
M10	-0.7230231	0.5985374	0.4662943	-0.7631498	0.7632057	0.5953022
M2	-0.2702467	-0.4045229	0.1897572	0.1528338	0.4009491	0.3168241
M3	-0.4928964	-0.0775830	0.2777637	-0.4016256	0.2979030	0.5528034
M4	-0.4797068	0.7662442	-0.2166083	-0.5704144	0.7178583	-0.2388118
M5	-0.5946456	-0.6852390	0.5099956	-0.6279539	-0.5903679	0.8708191
M6	0.5981419	-0.2428080	0.3700474	0.1467911	-0.5620152	0.3800916
M7	-0.4028705	-0.2620536	-0.2969209	0.3406279	0.5067387	0.3471849
M8	-0.3513185	-0.6778466	0.5233773	0.3072853	-0.6231626	0.5444628
M9	-0.4249585	-0.1595745	0.5184045	0.3130987	-0.5479842	0.5255984
Not.Correlated	-0.4570197	-0.1856715	0.6155754	-0.4134418	-0.2873009	0.5145339

6.3 enrichment_nes.tsv

Dimensions: 10 rows x 7 columns.

Table 5: enrichment_nes.tsv

pathway	CS_FruitSet	CS_Veraison	CS_Harvest	PN_FruitSet	PN_Veraison	PN_Harvest
M10	-3.959396	2.1492587	2.615432	-4.2536734	2.728041	3.118858
M2	-2.141531	-2.2761001	1.614254	0.9429160	1.839075	2.451360
M3	-3.541564	-0.4093924	2.099724	-3.1346970	1.304286	3.856798
M4	-3.403567	3.3899077	-1.230025	-4.3302647	3.115328	-1.424578
M5	-4.156126	-3.5193848	3.767544	-4.5950476	-3.083399	5.848706
M6	2.777735	-1.1341715	2.424789	0.7430314	-2.774107	2.283311
M7	-2.394055	-1.1530162	-1.444617	1.6434296	1.941217	1.979208
M8	-2.044511	-2.9146050	3.236747	1.4406157	-2.837919	2.984385
M9	-2.405142	-0.6670530	3.002033	1.4350954	-2.385699	2.803384
Not.Correlated	-2.773385	-0.8343909	3.831869	-2.5856587	-1.359322	2.973914

6.4 enrichment_padj.tsv

Dimensions: 10 rows x 7 columns.

Table 6: enrichment_padj.tsv

pathway	CS_FruitSet	CS_Veraison	CS_Harvest	PN_FruitSet	PN_Veraison	PN_Harvest
M10	0.0000000	0.0000225	0.0000024	0.0000000	0.0000000	0.0000000
M2	0.0000000	0.0000000	0.0004024	0.6718346	0.0000035	0.0000000
M3	0.0000000	1.0000000	0.0000024	0.0000000	0.0774487	0.0000000
M4	0.0000000	0.0000000	0.1552106	0.0000000	0.0000000	0.0377113
M5	0.0000000	0.0000000	0.0000000	0.0000000	0.0000000	0.0000000
M6	0.0000000	0.3151261	0.0000022	0.8589894	0.0000000	0.0000089
M7	0.0000037	0.3151261	0.0707547	0.0165577	0.0001668	0.0011522
M8	0.0002388	0.0000000	0.0000000	0.0829238	0.0000000	0.0000000
M9	0.0000254	1.0000000	0.0000000	0.0829238	0.0000047	0.0000001
Not.Correlated	0.0000000	1.0000000	0.0000000	0.0000025	0.0732555	0.0000000

6.5 module.tsv

Dimensions: 3050 rows x 2 columns.

Table 7: module.tsv - first 40 rows shown

genes	modules
VIT_19s0014g02740	M1
VIT_08s0056g01610	M1
VIT_12s0059g00590	M1
VIT_01s0010g02030	M1
VIT_08s0056g01600	M1
VIT_05s0049g00520	M1
VIT_02s0025g03390	M1
VIT_11s0016g00210	M1
VIT_16s0022g00960	M1
VIT_07s0005g00660	M1
VIT_19s0090g01370	M1
VIT_06s0004g03240	M1
VIT_06s0004g04650	M1
VIT_05s0049g00760	M1
VIT_17s0000g03280	M4
VIT_19s0014g02730	M1
VIT_01s0011g05110	M6
VIT_03s0180g00070	M1
VIT_00s0187g00160	M1
VIT_18s0001g14270	M1
VIT_19s0090g01340	M1

genes	modules
VIT_09s0002g05180	M1
VIT_07s0031g02210	M1
VIT_19s0090g01000	M1
VIT_03s0038g01510	M1
VIT_16s0022g01770	M1
VIT_05s0094g00340	M1
VIT_18s0001g14260	M1
VIT_02s0025g01050	M1
VIT_19s0090g01350	M1
VIT_02s0033g01330	M1
VIT_02s0033g01120	M10
VIT_04s0008g05010	M1
VIT_02s0025g03600	M1
VIT_02s0025g04330	M1
VIT_13s0019g04280	M1
VIT_10s0003g00090	M1
VIT_03s0038g02520	M1
VIT_11s0016g04920	M1
VIT_12s0028g01060	M1

The complete table is provided as a supplementary machine-readable file.

6.6 parameters.tsv

Dimensions: 11 rows x 2 columns.

Table 8: parameters.tsv

Parameter	Value
filter	TRUE
apply_vst	FALSE
filter_pval	0.1
cor_method	pearson
min_ngen	30
merge_similar	TRUE
diss_thresh	0.8
r2	0.706374247063269
beta	10
phi	0.500166177070297
n_mods	11

6.7 selected_genes.txt

Dimensions: 3049 rows x 1 columns.

Table 9: selected_genes.txt - first 40 rows shown

VIT_19s0014g02740
VIT_08s0056g01610
VIT_12s0059g00590
VIT_01s0010g02030
VIT_08s0056g01600
VIT_05s0049g00520
VIT_02s0025g03390
VIT_11s0016g00210
VIT_16s0022g00960
VIT_07s0005g00660
VIT_19s0090g01370
VIT_06s0004g03240
VIT_06s0004g04650
VIT_05s0049g00760
VIT_17s0000g03280
VIT_19s0014g02730
VIT_01s0011g05110
VIT_03s0180g00070
VIT_00s0187g00160
VIT_18s0001g14270
VIT_19s0090g01340
VIT_09s0002g05180
VIT_07s0031g02210
VIT_19s0090g01000
VIT_03s0038g01510
VIT_16s0022g01770
VIT_05s0094g00340
VIT_18s0001g14260
VIT_02s0025g01050
VIT_19s0090g01350
VIT_02s0033g01330
VIT_02s0033g01120
VIT_04s0008g05010
VIT_02s0025g03600
VIT_02s0025g04330
VIT_13s0019g04280
VIT_10s0003g00090
VIT_03s0038g02520
VIT_11s0016g04920
VIT_12s0028g01060
VIT_18s0001g12830

The complete table is provided as a supplementary machine-readable file.

6.8 summary_eigengene.tsv

Dimensions: 11 rows x 55 columns.

Table 10: summary_eigengene.tsv

[illegible]

6.9 summary_mean.tsv

Dimensions: 11 rows x 55 columns.

Table 11: summary mean.tsv

[illegible]

file	bytes	md5
results/figures/beta_r2.pdf	5718	83cfaa117a752d16bf6f42a5a5f22e79
results/figures/gsea.pdf	9481	a8e0c3c482e0c878231afe9227f971dc
results/figures/hist.pdf	5105	b5dafc083d17fd9d9eafecf8f9908d19
results/figures/mean_k.pdf	5468	450d829952c18a11db0e7b08df03f015
results/figures/mean_var.pdf	173388	2b4a703d540829622eb8b7a2ff99b6cd
results/figures/profile.pdf	715657	59f97246e8cd190a556c579099cc90a5
results/figures/qq.pdf	1392120	5f384f61a0ac5630906baf5453ba76d1
results/figures/sample_tree.pdf	6798	3f42183cbcd26905c0771ac939acc8d4
results/tables/beta_fit_indices.tsv	2527	6ec616e12b6c75d209918578a4a717cb
results/tables/enrichment_es.tsv	1273	5aaa94fc1ac921d3b7ecb34dea12834d
results/tables/enrichment_nes.tsv	1213	70f2139cabd0bf1825dbb315982b389f
results/tables/enrichment_padj.tsv	1336	57ff84dcd2d32ab4d7c3616426701ebd
results/tables/module.tsv	80042	c1f78a7ab02f444690e3c6ca9778dcdb
results/tables/modules_genes.gmt	72623	76a41959c5260da575318d5dbe1621f8
results/tables/parameters.tsv	247	31188bc3cad8c270fbc0a1877f93af2
results/tables/selected_genes.txt	57950	3fbe5f3efd8f2a6f69851ea0df1e0ad7
results/tables/summary_eigengene.tsv	12093	dc0ea4013ab26493d6dc8cb8bc7bfd89
results/tables/summary_mean.tsv	10893	b9b1595a8ca08e7e003592a0c1ad22ee
results/tables/summary_median.tsv	10071	ec27d56bf7f2c816bdb63d71004c66bd

8 Reproducibility

```
## ```text
## R version 4.6.1 (2026-06-24 ucrt)
## Platform: x86_64-w64-mingw32/x64
## Running under: Windows 11 x64 (build 26200)
##
## Matrix products: default
##   LAPACK version 3.12.1
##
## locale:
## [1] LC_COLLATE=Spanish_Chile.utf8  LC_CTYPE=Spanish_Chile.utf8    LC_MONETARY=Spanish_Chile
## [4] LC_NUMERIC=C                   LC_TIME=Spanish_Chile.utf8
##
## time zone: America/Santiago
## tzcode source: internal
##
## attached base packages:
## [1] stats      graphics  grDevices  utils      datasets  methods    base
##
## other attached packages:
## [1] CEMiTool_1.36.0    GEOquery_2.80.0    Biobase_2.72.0     BiocGenerics_0.58.1 generic
##
## loaded via a namespace (and not attached):
```

```

## [1] splines_4.6.1          ggplotify_0.1.3      R.oo_1.27.1
## [5] ggpp_0.6.1             polyclip_1.10-7      preprocessCore_1.74.0
## [9] confintr_1.0.2         XML_3.99-0.24        rpart_4.1.27
## [13] httr2_1.3.0            fastcluster_1.3.0    doParallel_1.0.17
## [17] lattice_0.22-9         MASS_7.3-65          backports_1.5.1
## [21] limma_3.68.5           Hmisc_5.3-0          rmarkdown_2.32
## [25] otl_0.2.0              ggtangle_0.1.2       zip_3.0.2
## [29] DBI_1.3.0              RColorBrewer_1.1-3   abind_1.4-8
## [33] R.utils_2.13.0         purrr_1.2.2          pracma_2.4.6
## [37] nnet_7.3-20            tweenr_2.0.3         rappdirs_0.3.4
## [41] gdtools_0.5.1          IRanges_2.46.0       S4Vectors_0.50.3
## [45] ggrepel_0.9.8          tidytree_0.4.8       ggpmisc_1.0.0
## [49] MatrixModels_0.5-4     codetools_0.2-20     DelayedArray_0.38.2
## [53] DOSE_4.5.0            DT_0.34.0            xml2_1.6.0
## [57] aplot_0.3.1           farver_2.1.2         matrixStats_1.5.0
## [61] dynamicTreeCut_1.63-1 base64enc_0.1-6       Seqinfo_1.2.0
## [65] Formula_1.2-6         survival_3.8-6       iterators_1.0.14
## [69] foreach_1.5.2         tools_4.6.1          ggnewscale_0.5.2
## [73] sna_2.8               Rcpp_1.1.2           glue_1.8.1
## [77] SparseArray_1.12.2     mgcv_1.9-4           xfun_0.60
## [81] MatrixGenerics_1.24.0 ggthemes_6.0.0       dplyr_1.2.1
## [85] BiocManager_1.30.27   fastmap_1.2.0        SparseM_1.84-2
## [89] digest_0.6.39         R6_2.6.1             gridGraphics_0.5-1
## [93] GO.db_3.23.1          RSQlite_3.53.3       R.methodsS3_1.8.2
## [97] fontLiberation_0.1.0  data.table_1.18.6.1  httr_1.4.9
## [101] S4Arrays_1.12.0       scatterpie_0.2.6     pkgconfig_2.0.3
## [105] blob_1.3.0            S7_0.2.2             impute_1.86.0
## [109] clusterProfiler_4.20.0 htmltools_0.5.9      fontBitstreamVera_0.1.1
## [113] scales_1.4.0          png_0.1-9            ggfun_0.2.1
## [117] knitr_1.52            rstudioapi_0.19.0    tzdb_0.5.0
## [121] curl_8.0.0            coda_0.19-4.1        statnet.common_4.13.0
## [125] nlme_3.1-169          cachem_1.1.0         stringr_1.6.0
## [129] foreign_0.8-91        AnnotationDbi_1.74.0 pillar_1.11.1
## [133] vctrs_0.7.3           tidydr_0.0.6         cluster_2.1.8.2
## [137] evaluate_1.0.5        tinytex_0.61         readr_2.2.0
## [141] compiler_4.6.1        rlang_1.3.0          crayon_1.5.3
## [145] labeling_0.4.3        plyr_1.8.9           fs_2.1.0
## [149] stringi_1.8.9         network_1.20.0       WGCNA_1.74
## [153] Biostrings_2.80.2     lazyeval_0.2.3       fontquiver_0.2.1
## [157] quantreg_6.1          Matrix_1.7-5         hms_1.1.4
## [161] bit64_4.8.6           ggplot2_4.0.3        KEGGREST_1.52.2
## [165] SummarizedExperiment_1.42.0 igraph_2.3.3         memoise_2.0.1
## [169] fastmatch_1.1-8       bit_4.6.0            gson_0.2.1
## [173] polynom_1.4-1
##
##
##

```

9 Notes for manuscript use

This document is a computational analysis report, not a substitute for biological interpretation. Statements about biological mechanism, causal relationships, novelty, and comparison with previous literature should be supported separately with appropriate experimental evidence and verified references.

10 Post-CEMiTool module statistics

This section is an additional statistical layer performed after construction of the beta-10 CEMiTool network. It is not part of the native CEMiTool report and is not stored inside the CEMiTool object.

10.1 Network calibration and sensitivity

Beta 10 is the primary network and beta 7 is retained as a sensitivity analysis. Module labels are arbitrary across runs, so robustness is evaluated using shared genes rather than matching labels by name.

Table 14: Best beta7-to-beta10 matches based on gene membership.

Module_beta7	Module_beta10	Shared_genes	Jaccard	Percent_beta7_retained
M1	M1	2150	0.912	91.9
M2	M2	203	0.864	90.6
M3	M3	105	0.761	93.8
M4	M5	101	0.894	95.3
M5	M4	91	0.705	92.9
M6	M8	46	0.630	64.8
M7	M10	34	0.723	81.0
M8	M9	40	0.952	100.0
Not.Correlated	Not.Correlated	17	0.298	100.0

10.2 Module eigengene model

For each beta-10 biological module, PC1 of standardized module-gene expression was used as the module eigengene. PC1 sign was oriented to correlate positively with mean module expression.

Model:

```
Eigengene ~ Cultivar * Stage + Year
```

Type III ANOVA used sum-to-zero contrasts and Benjamini-Hochberg FDR.

10.3 FDR overview by module

Table 15: FDR by module and model effect.

	Module	FDR_Cultivar	FDR_Stage	FDR_Year	FDR_Cultivar:Stage
1	M1	0.4943000	0.0000000	0.0031450	0.039650
5	M10	0.0000028	0.0000000	0.1165000	0.001052
9	M2	0.0000000	0.0002871	0.0052300	0.019510
13	M3	0.0136800	0.0070290	0.0000000	0.019510
17	M4	0.1448000	0.0000000	0.7054000	0.884000
21	M5	0.0000000	0.0000000	0.0011460	0.000000
25	M6	0.0401300	0.0000000	0.0031450	0.100400
29	M7	0.0732200	0.0000001	0.0009819	0.540700
33	M8	0.5370000	0.0005303	0.0006119	0.972900
37	M9	0.6657000	0.0954400	0.0205000	0.567300

10.4 Significant Cultivar x Stage interactions

5 of 10 modules passed $FDR < 0.05$. Strongest: **M5** ($FDR=5.29e-22$).

Table 16: Significant Cultivar x Stage interactions.

Module	F value	p_value	FDR
M5	190.926286	5.29e-23	5.29e-22
M10	10.235546	2.10e-04	1.05e-03
M2	5.562952	6.86e-03	1.95e-02
M3	5.402947	7.81e-03	1.95e-02
M1	4.274767	1.98e-02	3.97e-02

A significant interaction means that the Cabernet-versus-Pinot difference changes with developmental stage. It does not by itself demonstrate a causal relationship with skin thickness.

10.5 Cabernet Sauvignon vs Pinot noir within stage

The contrast is Cabernet Sauvignon minus Pinot noir. Negative estimates therefore indicate a lower module eigengene in Cabernet Sauvignon.

Table 17: Significant cultivar contrasts within stage.

Module	Stage	estimate	Direction	p.value	FDR_global
M5	Harvest	-22.279	Cabernet Sauvignon < Pinot noir	2.85e-28	8.56e-27
M2	Veraison	-25.367	Cabernet Sauvignon < Pinot noir	7.98e-16	1.20e-14
M2	Harvest	-21.832	Cabernet Sauvignon < Pinot noir	1.28e-13	1.28e-12
M2	FruitSet	-15.558	Cabernet Sauvignon < Pinot noir	2.40e-09	1.80e-08

Module	Stage	estimate	Direction	p.value	FDR_global
M10	Harvest	-3.356	Cabernet Sauvignon < Pinot noir	1.79e-07	1.07e-06
M3	Harvest	-6.989	Cabernet Sauvignon < Pinot noir	7.62e-05	3.81e-04
M10	Veraison	-2.128	Cabernet Sauvignon < Pinot noir	3.18e-04	1.36e-03
M6	FruitSet	6.146	Cabernet Sauvignon > Pinot noir	2.40e-03	8.99e-03
M5	Veraison	-2.458	Cabernet Sauvignon < Pinot noir	8.78e-03	2.93e-02

10.6 Year effect

Year passed FDR < 0.05 in **8 of 10 modules**.

Table 18: Modules with significant Year effect.

	Module	p_value	FDR
3	M1	1.89e-03	3.14e-03
11	M2	3.66e-03	5.23e-03
15	M3	1.75e-21	1.75e-20
23	M5	4.59e-04	1.15e-03
27	M6	1.64e-03	3.14e-03
31	M7	2.95e-04	9.82e-04
35	M8	1.22e-04	6.12e-04
39	M9	1.64e-02	2.05e-02

This supports keeping 2012, 2013 and 2014 explicit. The current additive model adjusts for Year but does not yet test whether the Cultivar x Stage pattern itself is stable across years.

10.7 Interpretation boundaries and next step

- GSE98923 is berry pericarp, not isolated skin.
- There is no direct skin-thickness phenotype in this dataset.
- Only two cultivars are compared.
- Several modules have a significant Year effect.
- The baseline uses the historical GEO RPKM matrix.

The next planned statistical test is:

```
Eigengene ~ Cultivar * Stage * Year
```

followed by Cabernet-versus-Pinot contrasts within each Stage x Year combination. Modules that remain robust can then proceed to functional annotation, hub-gene prioritization and skin-only validation.