

# Higgs boson production at the Large Hadron Collider: accurate theoretical predictions at higher orders in QCD

Jesús Urtasun Elizari

PhD presentation - Milan, February 25th, 2022



UNIVERSITÀ  
DEGLI STUDI  
DI MILANO



European  
Research  
Council

This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.

# Outline

## ① QCD and collider physics

- The strong interactions
- Asymptotic freedom and pQCD
- QCD Factorization
- Phenomenology at the LHC

## ② All order perturbative resummation

- Higher order radiative corrections
- Resummation of large logarithmic corrections
- Resummed component, asymptotic and fixed-order

## ③ HTurbo numerical implementation

- Higgs production at the LHC
- HTurbo numerical implementation
- $N^3LL$  implementation

## ④ Results & Conclusions

# Part I

## QCD and collider physics

# Introduction

## QCD and the strong interactions

- QCD is the theory of the strong interactions
- Part of the Standard describing fundamental interactions at the TeV scale
- Fundamental objects described as irreducible representations of Lorentz group
- Particles as local excitations of fields with quantum mechanical behavior

# Introduction

## QCD and the strong interactions

How to explore proton's inner structure?




- At different scales, hadrons show different behavior
- From point-like to complex internal dynamics
- Scattering experiments (DIS) and hadronic physics (LHC)

"A way of describing high energy collisions is to consider any hadron as a composite object of point-like constituents → **partons**" R.Feynman, 1969

# QCD and collider physics

## Asymptotic freedom and pQCD



- Parton model as LO approximation to QCD
- Real QCD coupling strength changes with energy
- At high energies the hadron involves extremely complex internal dynamics

QCD is strongly coupled at large distances / low energies  $\longrightarrow$  confinement

Non-perturbative physics

# QCD and collider physics

## Asymptotic freedom and pQCD


- Running coupling given by Renormalization Group Equation (RGE)

$$\mu \frac{d\alpha_s(\mu)}{d\mu} = \beta(\alpha_s(\mu)) = - \sum_{n=0}^{\infty} \beta_n \left( \frac{\alpha_s}{\pi} \right)^{n+1}$$

- Coupling  $\alpha_s$  evolves with scale  $\mu$  as given by RGE  $\rightarrow$  LO behavior driven by  $\beta_0$
- $\beta_0^{\text{QED}} < 0 \implies$  strongly coupled at large energies, UV divergent
- $\beta_0^{\text{QCD}} > 0 \implies$  weakly coupled at large energies, IR divergent

# QCD and collider physics

## Asymptotic freedom and pQCD



- Running coupling given by Renormalization Group Equation (RGE)

$$\alpha_s(\mu) = \frac{1}{\beta_0 \log\left(\frac{\mu^2}{\Lambda_{\text{QCD}}^2}\right)}$$

- $\beta_0$  LO of the  $\beta$  function, is  $> 0$
- $\Lambda_{\text{QCD}}$ , parameter that defines value of the coupling at large scales


QCD is weakly coupled for  $\mu \gg \Lambda_{\text{QCD}} \rightarrow$  asymptotically free

Perturbative Quantum Chromodynamics (pQCD)



# QCD and collider physics


## Hadronic processes and factorization



- LHC physics rely on hadronic collisions  $\rightarrow$  pQCD
- Compute **cross section**  $\sigma^F \rightarrow$  probability for a given process

# QCD and collider physics

## Hadronic processes and factorization




Compute hadronic cross sections is a **hard problem**  $\rightarrow$  **QCD Factorization**

$$\sigma^F(p_1, p_2) = \int_0^1 dx_1 dx_2 f_\alpha(x_1, \mu_F^2) * f_\beta(x_2, \mu_F^2) * \hat{\sigma}_{\alpha\beta}^F(x_1 p_1, x_2 p_2, \alpha_s(\mu_R^2), \mu_F^2)$$

# QCD and collider physics

## Hadronic processes and factorization




- Parton densities (PDFs)  $f_a(x_i, \mu_F^2)$ : non perturbative but universal
- Partonic cross section  $\hat{\sigma}_{\alpha\beta}^F$ : process dependent but computable as perturbative series in  $\alpha_s$

# QCD and collider physics

## Parton densities

Parton Distribution Functions: probability distribution of finding a particular parton ( $u$ ,  $d$ , ...,  $g$ ) carrying a fraction  $x$  of the proton's momentum




- Each parton has a different PDF  $\rightarrow u(x), d(x), \dots, g(x)$
- PDFs can not predicted and yet can not measured  $\rightarrow$  extracted from data (MSTW, CTEQ, NNPDF collaborations)
- The N3PDF project: Machine Learning for PDFs determination  
[Urtasun-Elizari et al.] ref. at [1910.07049](#)

# QCD and collider physics

## Partonic cross section and pQCD

- Born cross section is the leading-order (LO) term of the perturbative series
- $\sigma^{(1)}, \sigma^{(2)}, \sigma^{(3)}$  are the NLO, NNLO, N<sup>3</sup>LO corrections



$$\hat{\sigma} = \sigma^{\text{Born}} \left( 1 + \alpha_s \sigma^{(1)} + \alpha_s^2 \sigma^{(2)} + \alpha_s^3 \sigma^{(3)} + \dots \right)$$

Lower order predictions strongly depend on the auxiliary / unphysical scales


Need higher order corrections to increase theoretical accuracy!

# QCD and collider physics

## LHC phenomenology

Standard Model Total Production Cross Section Measurements

Status: July 2021




## Part II

### All order resummation


# Resummation in QCD

Higher order corrections - need for resummation


- 1 Calculation of higher order corrections is **not an easy task** due to **infrared (IR) soft and collinear singularities**
- 2 Final state singularities **cancel** by combining real and virtual contributions  $\rightarrow$  KLN theorem
- 3 Initial state collinear singularities **factorized** inside the PDFs




a



b



c



d


Cancellation only works in completely inclusive final states!



# Resummation in QCD

## $q_\perp$ resummation

- Describing exclusive final states
- Study the differential  $q_\perp$  distribution  
 $h_1(p_1) + h_2(p_2) \longrightarrow F(M, q_\perp) + X$



$$\int_0^{Q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}}{dq_\perp^2} \sim c_0 + \alpha_s(c_{12}L^2 + c_{11}L + c_{10}) + \dots, \quad \text{where} \quad L = \ln(M^2/q_\perp^2)$$

$\alpha_s L^2$	$\alpha_s L$	$\dots$	$\mathcal{O}(\alpha_s)$
$\alpha_s^2 L^4$	$\alpha_s^2 L^3$	$\dots$	$\mathcal{O}(\alpha_s^2)$
$\dots$	$\dots$	$\dots$	$\dots$
$\alpha_s^n L^{2n}$	$\alpha_s^n L^{2n-1}$	$\dots$	$\mathcal{O}(\alpha_s^n)$
dominant logs	$\dots$	$\dots$	$\dots$

Truncated fixed-order predictions  $\rightarrow$  enhanced  $\alpha_s^n \ln^m(M^2/q_\perp^2)$  appear

# Resummation in QCD

## $q_\perp$ resummation

Separate partonic  $q_\perp$  distribution as follows

$$\frac{d\hat{\sigma}_{ab}}{dq_\perp^2} = \left[ \frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_\perp^2} \right]_{\text{l.a.}} + \left[ \frac{d\hat{\sigma}_{ab}^{(\text{fin.})}}{dq_\perp^2} \right]_{\text{f.o.}}, \quad \text{such that}$$

$$\int_0^{q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_\perp^2} \sim \sum \alpha_s^n \log^m \left( \frac{M^2}{q_\perp^2} \right) \quad \text{for } q_\perp \rightarrow 0$$
$$\lim_{q_\perp \rightarrow 0} \int_0^{q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}_{ab}^{(\text{fin.})}}{dq_\perp^2} = 0$$

Resummed and finite components can be matched (LL+LO, NLL+NLO, NNLO+NNLL, ...) to have uniform accuracy in a wide range of  $q_\perp$

# Resummation in QCD

## Resummed component

Resummation holds in impact parameter space  $b$

$$\frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_{\perp}^2} = \frac{M^2}{\hat{s}} \int db \frac{b}{2} J_0(bq_{\perp}) \mathcal{W}_{ab}(b, M)$$

with  $\mathcal{W}_{ab}$  also expressed in Mellin space (with respect to  $z = M^2/\hat{s}$ )

$$\mathcal{W}_N(b, M) = \mathcal{H}_N(\alpha_s) \times \exp\{\mathcal{G}_N(\alpha_s, L)\} \quad \text{being} \quad L \equiv \log(M^2 b^2)$$

- Large logarithms exponentiated in the universal Sudakov form factor  $\mathcal{G}_N(\alpha_s, L)$
- Constant (b-independent) terms factorized in the process dependent hard factor  $\mathcal{H}_N(\alpha_s)$

# Resummation in QCD

## Resummed component

Sudakov factor  $\mathcal{G}_N$  and hard coefficient  $\mathcal{H}_N$  can be expanded as perturbative series in  $\alpha_s$

$$\mathcal{G}_N(\alpha_s, L) = L g^{(1)}(\alpha_s L) + g^{(2)}(\alpha_s L) + \frac{\alpha_s}{\pi} g^{(3)}(\alpha_s L) + \dots$$

$$\mathcal{H}_N(\alpha_s) = 1 + \alpha_s \mathcal{H}^{(1)} + \alpha_s^2 \mathcal{H}^{(2)} + \dots$$

For each new order implement a factor of  $\mathcal{G}_N$  and Hard  $\mathcal{H}_N$

$$\text{LL}(\sim \alpha_s^n L^{n+1}) : g^{(1)}, \hat{\sigma}^{(0)}$$

$$\text{NLL}(\sim \alpha_s^n L^n) : g^{(2)}, \mathcal{H}^{(1)}$$

$$\text{NNLL}(\sim \alpha_s^n L^{n-1}) : g^{(3)}, \mathcal{H}^{(2)}$$

Each term  $g^{(i)}$  and  $\mathcal{H}^{(i)}$  in the series becomes increasingly complicated

Current codes able to produce only up to NNLL predictions!

## Part III

### HTurbo numerical implementation

# HTurbo


## Resummation for Higgs differential distribution

- Fast and accurate predictions for Higgs boson production cross section
- Predictions for differential cross section  $d\sigma^H/dq_\perp^2$
- Numerical implementation of resummed and finite components

$$d\sigma_{(N)NLL+(N)LO}^H = d\sigma_{(N)NLL}^{(\text{res.})} - d\sigma_{(N)LO}^{(\text{asy.})} + d\sigma_{(N)LO}^{(\text{f.o.})}$$

$$d\sigma_{(N)NLL}^{(\text{res.})} = \hat{\sigma}_{LO}^H \times \mathcal{H}_{(N)LO} \times \exp \mathcal{G}_{(N)NLL}$$


$$d\sigma_{(N)LO}^{(\text{asy.})} = \hat{\sigma}_{LO}^H \times \Sigma_{(N)LO}$$



# HTurbo

## Predictions for Higgs $q_{\perp}$ distribution

- $q_{\perp}$  resummation implemented in numerical codes **HRes**, **HRes**, **HNNLO** [Catani, de Florian, Ferrera, Grazzini, Tommasini]
- Higher order accuracy require **high computation times**
- NNLL predictions can take up to 48h  
→ need for **fast numerical implementations**



Codes producing fast and accurate predictions are needed for precision era of the LHC

# HTurbo

Starting point: DYTurbo

Numerical code **DYTurbo** [Camarda et al.] ref. at [1910.07049](#), fast and precise  $q_\perp$  resummation and several improvements for Drell-Yan ( $h_1 h_2 \rightarrow V + X \rightarrow l^+ l^- + X$ )

**First goal:** set up a numerical code for Higgs boson production starting from **DYTurbo**

- Set LO amplitude  $gg \rightarrow H$
- Set Sudakov and Hard coefficients for resummed component
- Set  $\Sigma$  coefficients for asymptotic term
- Implement MC producing the LO and NLO H+jet cross sections
- Compare with **HRes** and **HqT**

**Final goal:** extend theoretical accuracy up to  $N^3\text{LL}+N^3\text{LO}$



# HTurbo

## Code optimization

Optimized reimplementation of **HqT**, **HRes** and **HNNLO** for  $q_T$ -resummation



- **C++** structure with **Fortran** interfaces → Multi-threading
- Optimization in the integration routines / integral transforms
  - Factorize boson and decay kinematics
  - Gauss-Legendre quadrature rules (1-dim.)
  - Vegas/Cuhre through **Cuba** (multi-dim.)

Comparison **HRes** and **HTurbo** - speed performance

Predictions	<b>HRes</b>	<b>HTurbo</b>
resummed NNLL	10h	10'
combined NNLO+NNLL	48h	2h

# Results



## Comparison HTurbo and HRes - NLL resummed



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nlo\_as\_0118 PDF set

# Results



## Comparison HTurbo and HRes - NNLL resummed



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nnlo\_as\_0118 PDF set

# Results



## Comparison HTurbo and HRes - LO asymptotic



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nlo\_as\_0118 PDF set

# Results



## Comparison HTurbo and HRes - NLO asymptotic



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nnlo\_as\_0118 PDF set

# Results



## Comparison HTurbo and HRes - LO fixed-order



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nlo\_as\_0118 PDF set

# Results

## Comparison HTurbo and HRes - NLO fixed-order



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- CM energy  $\sqrt{s} = 13$  GeV and PDF set NNPDF31\_nnlo\_as\_0118 PDF set

# Results

## N<sup>3</sup>LL implementation

Sudakov factor  $\mathcal{G}_N$  and hard coefficient  $\mathcal{H}_N$  can be expanded as perturbative series in  $\alpha_s$

$$\mathcal{G}_N(\alpha_s, L) = L g^{(1)}(\alpha_s L) + g^{(2)}(\alpha_s L) + \frac{\alpha_s}{\pi} g^{(3)}(\alpha_s L) + \frac{\alpha_s}{\pi} g^{(4)}(\alpha_s L) + \dots$$

$$\mathcal{H}_N(\alpha_s) = 1 + \alpha_s \mathcal{H}^{(1)} + \alpha_s^2 \mathcal{H}^{(2)} + \alpha_s^2 \mathcal{H}^{(3)} + \dots$$

For each new order implement a factor of  $\mathcal{G}_N$  and Hard  $\mathcal{H}_N$

$$\text{LL}(\sim \alpha_s^n L^{n+1}) : g^{(1)}, \hat{\sigma}^{(0)}$$

$$\text{NLL}(\sim \alpha_s^n L^n) : g^{(2)}, \mathcal{H}^{(1)}$$

$$\text{NNLL}(\sim \alpha_s^n L^{n-1}) : g^{(3)}, \mathcal{H}^{(2)}$$



$$\text{N}^3\text{LL}(\sim \alpha_s^n L^{n-2}) : g^{(4)}, \mathcal{H}^{(3)}$$

Start by building predictions up to NNLO+NNLL, then add **N<sup>3</sup>LO+N<sup>3</sup>LL**



# Results

## $N^3\text{LL}$ implementation



- Cross section for fully inclusive (LHS) and fiducial (RHS) phase space ✓
- Implementation of  $N^3\text{LL}$  factors following [Li - Zhu], 1604.01404 [Von Manteuffel et al.], 2002.04617

# Summary & Conclusions

- ① Fast and accurate predictions are needed towards the precision era of the LHC
- ② Developing a novel numerical code, **HTurbo**, which implements  $q_\perp$  resummation for Higgs boson production
- ③ HTurbo is faster than any of the existing codes
- ④ Outlook of thesis work:
  - Add  $N^3\text{LO}+N^3\text{LL}$  prediction
  - Perform phenomenological studies comparing with LHC data

# Discussion & next steps

- ① Fast and accurate predictions are needed towards the precision era of the LHC
- ② Developing a novel numerical code, **HTurbo**, which implements  $q_{\perp}$  resummation for Higgs boson production
- ③ HTurbo is faster than any of the existing codes
- ④ Outlook of thesis work:
  - Add  $N^3\text{LO}+N^3\text{LL}$  prediction
  - Perform phenomenological studies comparing with LHC data

Thank you!



This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.

# Back up

# Back up

# Back up