

Higgs boson production at the Large Hadron Collider: accurate theoretical predictions at higher orders in QCD

Jesús Urtasun Elizari

PhD seminar - Milan, September 2021



UNIVERSITÀ
DEGLI STUDI
DI MILANO



European
Research
Council

This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.

Outline

- ① QCD and collider physics
 - QCD Factorization
 - Partonic cross section and perturbative QCD
- ② All order perturbative resummation
 - Higher order radiative corrections
 - Resummation of large logarithmic corrections
- ③ Precise and fast predictions for Higgs boson physics
 - Higgs production at the LHC
 - HTurbo numerical code
 - Preliminary results & Conclusions


QCD and collider physics

QCD and collider physics

LHC physics


Standard Model Production Cross Section Measurements

Status: July 2018



QCD and collider physics

Factorization theorem




Compute hadronic cross sections is a **hard problem** \rightarrow QCD Factorization

$$\sigma^F(p_1, p_2) = \int_0^1 dx_1 dx_2 f_\alpha(x_1, \mu_F^2) * f_\beta(x_2, \mu_F^2) * \hat{\sigma}_{\alpha\beta}^F(x_1 p_1, x_2 p_2, \alpha_s(\mu_R^2), \mu_F^2)$$

QCD and collider physics


Partonic cross section



- Parton densities (PDFs) $f_{\alpha}(x_i, \mu_F^2)$: non perturbative but universal
- Partonic cross section $\hat{\sigma}_{\alpha\beta}^F$: process dependent but computable as perturbative series in α_s

QCD and collider physics

QCD coupling



- Running coupling given by Renormalization Group Equation (RGE)

$$\alpha_s(\mu) = \frac{1}{\beta_0 \log\left(\frac{\mu^2}{\Lambda_{\text{QCD}}^2}\right)}$$

- β_0 LO of the β function, is > 0
- Λ_{QCD} , parameter that defines value of the coupling at large scales


QCD is weakly coupled for $\mu \gg \Lambda_{\text{QCD}} \rightarrow$ asymptotically free

Perturbative Quantum Chromodynamics (pQCD)

QCD and collider physics

Perturbative QCD

- Born cross section is the leading-order (LO) term of the perturbative series
- $\sigma^{(1)}, \sigma^{(2)}, \sigma^{(3)}$ are the NLO, NNLO, N3LO corrections



$$\hat{\sigma} = \sigma^{\text{Born}} \left(1 + \alpha_s \sigma^{(1)} + \alpha_s^2 \sigma^{(2)} + \alpha_s^3 \sigma^{(3)} + \dots \right)$$


Lower order predictions strongly depend on the auxiliary and unphysical renormalization and factorization scales \rightarrow **Need higher order corrections to increase theoretical accuracy!**

All order perturbative resummation

Resummation in QCD

Higher order corrections

- 1 Calculation of higher order corrections is **not an easy task** due to **infrared (IR) soft and collinear singularities**
- 2 Final state singularities **cancel** by combining real and virtual contributions
- 3 Initial state collinear singularities **factorized** inside the PDFs




Resummation in QCD

q_\perp resummation

Study the differential q_\perp distribution

$$h_1(p_1) + h_2(p_2) \longrightarrow F(M, \mathbf{q}_\perp) + X$$



$$\int_0^{Q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}}{dq_\perp^2} \sim c_0 + \alpha_s (c_{12} L^2 + c_{11} L + c_{10}) + \dots, \quad \text{where} \quad L = \ln(q_\perp/M^2)$$

$\alpha_S L^2$	$\alpha_S L$	\dots	$\mathcal{O}(\alpha_S)$
$\alpha_S^2 L^4$	$\alpha_S^2 L^3$	\dots	$\mathcal{O}(\alpha_S^2)$
\dots	\dots	\dots	\dots
$\alpha_S^n L^{2n}$	$\alpha_S^n L^{2n-1}$	\dots	$\mathcal{O}(\alpha_S^n)$
dominant logs	\dots	\dots	\dots

Truncated fixed order predictions \rightarrow enhanced $\alpha_s^n \ln^m(M^2/q_\perp^2)$ appear

Resummation in QCD

q_\perp resummation

Separate partonic q_\perp distribution as follows

$$\frac{d\hat{\sigma}_{ab}}{dq_\perp^2} = \left[\frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_\perp^2} \right]_{\text{l.a.}} + \left[\frac{d\hat{\sigma}_{ab}^{(\text{fin.})}}{dq_\perp^2} \right]_{\text{f.o.}}, \quad \text{such that}$$

$$\int_0^{q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_\perp^2} \sim \sum \alpha_s^n \log^m \frac{M^2}{q_\perp^2} \quad \text{for } q_\perp \rightarrow 0$$
$$\lim_{q_\perp \rightarrow 0} \int_0^{q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}_{ab}^{(\text{fin.})}}{dq_\perp^2} = 0$$

Resummed and finite components can be matched (LL+LO, NLL+NLO, NNLO+NNLL, ...) to have uniform accuracy in a wide range of q_\perp

Resummation in QCD

q_\perp resummation

Resummation holds in impact parameter space b

$$\frac{d\hat{\sigma}_{ab}^{(\text{res.})}}{dq_\perp^2} = \frac{M^2}{\hat{s}} \int db \frac{b}{2} J_0(bq_\perp) \mathcal{W}_{ab}(b, M)$$

with \mathcal{W}_{ab} also expressed in Mellin space (with respect to $z = M^2/\hat{s}$)

$$\mathcal{W}_N(b, M) = \mathcal{H}_N(\alpha_s) \times \exp\{\mathcal{G}_N(\alpha_s, L)\} \quad \text{being} \quad L \equiv \log(M^2 b^2)$$


- Large logarithms exponentiated in the universal Sudakov form factor $\mathcal{G}_N(\alpha_s, L)$
- Constant (b -independent) terms factorized in the process dependent hard factor $\mathcal{H}_N(\alpha_s)$

Accurate predictions for Higgs boson at the LHC

HqT and HRes

Predictions for Higgs q_{\perp} distribution

- q_{\perp} resummation implemented in numerical codes HqT and HRes [Catani, de Florian, Ferrera, Grazzini, Tommasini]
- Higher order accuracy require **high computation times**
- Codes producing fast and accurate predictions are needed for precision era of the LHC



HTurbo

Starting point DYTurbo

Numerical code **DYTurbo** [Camarda et al.] ref. at [1910.07049](#), fast and precise q_{\perp} resummation and several improvements for Drell-Yan ($h_1 h_2 \rightarrow V + X \rightarrow l^+ l^- + X$)

- **First goal**: set up a numerical code for Higgs boson production starting from **DYTurbo**
- Set LO amplitude $gg \rightarrow H$
- Set Sudakov and Hard coefficients for Higgs production
- Compare with **HRes** and **HqT**

Final goal: extend theoretical accuracy up to $N^3\text{LL}+N^3\text{LO}$

HTurbo

Starting point DYTurbo

Both Sudakov factor \mathcal{G}_N and hard coefficient \mathcal{H}_N can be expanded as perturbative series in α_s

$$\mathcal{G}_N(\alpha_s, L) = L g^{(1)}(\alpha_s L) + g^{(2)}(\alpha_s L) + \frac{\alpha_s}{\pi} g^{(3)}(\alpha_s L) + \dots$$

$$\mathcal{H}_N(\alpha_s) = 1 + \alpha_s \mathcal{H}^{(1)} + \alpha_s^2 \mathcal{H}^{(2)} + \dots$$

For each new order implement a factor of \mathcal{G}_N and Hard \mathcal{H}_N

$$\text{LL}(\sim \alpha_s^n L^{n+1}) : g^{(1)}, \hat{\sigma}^{(0)}$$

$$\text{NLL}(\sim \alpha_s^n L^n) : g^{(2)}, \mathcal{H}^{(1)}$$

$$\text{NNLL}(\sim \alpha_s^n L^{n-1}) : g^{(3)}, \mathcal{H}^{(2)}$$

Start by building predictions up to NNLO+NNLL, then add
N³LO+N³LL

Reimplementation of **HqT** and **HRes** for q_T -resummation


- **C++** structure with **Fortran** interfaces \rightarrow Multi-threading
- Optimization in the integration routines / integral transforms
 - Factorize boson and decay kinematics
 - Gauss-Legendre quadrature rules (1-dim.)
 - Vegas/Cuhre through **Cuba** (multi-dim.)

Comparison **HRes** and **HTurbo** - speed performance

Predictions	HRes	HTurbo
resummed NNLL	10h	10'
combined NNLO+NNLL	20h	1h

Results


Comparison HTurbo and HqT - LL



- HTurbo q_{\perp} distribution vs HRes and HqT at LL
- Excellent numerical agreement up to the 0.1% level

Results


Comparison HTurbo and HqT - NLL



- HTurbo q_{\perp} distribution vs HRes and HqT at NLL
- Excellent numerical agreement up to the 0.1% level

Results



Comparison HTurbo and HqT - NNLL



- HTurbo q_{\perp} distribution vs HRes and HqT at NNLL
- Excellent numerical agreement up to the 0.1% level

Results


Comparison HTurbo and HRes - resummed



- Represent full (LHS) and fiducial (RHS) phase space ✓
- Agreement up to resummed NNLL \rightarrow ready for N^3 LL

Results


Comparison HTurbo and HRes - fixed order



- Represent fiducial (RHS) phase space - HRes vs HTurbo ✓
- Agreement up to fixed order NLO \rightarrow ready for N³LL

Results

Comparison HTurbo and HRes - finite



- Represent fiducial phase space - HRes vs HTurbo ✓
- Agreement up to finite NNLL+NLO \longrightarrow ready for $N^3\text{LL}$

Summary & Conclusions

- ① Fast and accurate predictions are needed towards the precision era of the LHC
- ② Developing a novel numerical code, **HTurbo**, which implements q_{\perp} resummation for Higgs boson production
- ③ HTurbo is faster than any of the existing codes
- ④ Outlook of thesis work:
 - Add $N^3\text{LO}+N^3\text{LL}$ prediction
 - Perform phenomenological studies comparing with LHC data

Thank you!



This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.