

# High precision perturbative QCD predictions for LHC physics

Jesús Urtasun Elizari

Milan, February 2021



UNIVERSITÀ  
DEGLI STUDI  
DI MILANO



European  
Research  
Council

This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.

# Outline


- ① QCD factorization in a nutshell
  - QCD factorization
  - Parton Densities and partonic cross section
- ② The N3PDF project
  - Machine Learning for PDFs
- ③ Fast predictions for resummed distributions
  - Higher order corrections and all orders resummation
  - Higgs production at the LHC
  - **HTurbo** numerical code
- ④ Conclusions

## QCD factorization in a nutshell

# QCD factorization in a nutshell

## Factorization theorem

$$h_1(p_1) + h_2(p_2) \rightarrow F + X$$



Factorize process as **PDFs** and **partonic (hard) interaction**


$$\sigma^F(p_1, p_2) = \sum_{\alpha, \beta} \int_0^1 dx_1 dx_2 f_{\alpha/h_1}(x_1, \mu_F^2) * f_{\beta/h_2}(x_2, \mu_F^2) * \hat{\sigma}_{\alpha\beta}^F(x_1 p_1, x_2 p_2, \alpha_s(\mu_R^2), \mu_F^2)$$

# QCD factorization in a nutshell

## Perturbative QCD

Precise QCD predictions require precise knowledge of **PDFs** and **partonic cross section  $\hat{\sigma}$**

- Born cross section is the leading-order (LO) term of the perturbative series
- $\sigma^{(1)}, \sigma^{(2)}, \sigma^{(3)}$  are the NLO, NNLO, N<sup>3</sup>LO corrections



$$\hat{\sigma} = \sigma^{\text{Born}} \left( 1 + \alpha_s \sigma^{(1)} + \alpha_s^2 \sigma^{(2)} + \alpha_s^3 \sigma^{(3)} + \dots \right)$$

LO predictions strongly depend on unphysical renormalization and factorization scales

**Need higher order corrections to increase theoretical accuracy!**

## The N3PDF project

# The N3PDF project

## General structure of n3fit


Parton Distribution Functions (PDFs) can not be predicted or measured

PDFs need to be extracted from data!



- Use TensorFlow and Keras to determine the PDFs
- Use Stochastic Gradient Descent **n3fit** replacing primitive genetic algorithms
- See paper by S.Carraza - J.Cruz-Martinez  
"Towards a new generation of parton densities with deep learning models",  
<https://arxiv.org/abs/1907.05075>


## Neural Network model for PDF fits $\rightarrow$ n3fit



- Build a NN model to compute  $\sigma_{pred}$  observables from a grid  $x_i$
- Perform  $\chi^2$  minimization comparing with data
- Update values of PDF  $\rightarrow$  Fit



## Neural Network model for PDF fits $\rightarrow$ n3fit



- 1 TF relies in symbolic computation  $\rightarrow$  High memory usage
- 2 Implement c++ operator replacing the convolution
- 3 Further details in Urtasun-Elizari et al.

"Towards hardware acceleration for parton densities estimation",  
<https://arxiv.org/abs/1909.10547>


## HTurbo: Fast predictions for resummed distributions

# HTurbo

All order  $q_\perp$  resummation

Study the differential  $q_\perp$  distribution

$$h_1(p_1) + h_2(p_2) \longrightarrow F(M, \mathbf{q}_\perp) + X$$



$$\int_0^{Q_\perp^2} dq_\perp^2 \frac{d\hat{\sigma}}{dq_\perp^2} \sim c_0 + \alpha_s (c_{12} L^2 + c_{11} L + c_{10}) + \dots, \quad \text{being} \quad L = \ln(q_\perp/M^2)$$

$\alpha_S L^2$	$\alpha_S L$	$\dots$	$\mathcal{O}(\alpha_S)$
$\alpha_S^2 L^4$	$\alpha_S^2 L^3$	$\dots$	$\mathcal{O}(\alpha_S^2)$
$\dots$	$\dots$	$\dots$	$\dots$
$\alpha_S^n L^{2n}$	$\alpha_S^n L^{2n-1}$	$\dots$	$\mathcal{O}(\alpha_S^n)$
dominant logs	$\dots$	$\dots$	$\dots$

Truncated fixed order predictions lead to **logarithmic enhancement**  $\alpha_s^n \ln^m(M^2/q_\perp^2)$

**All order resummation is needed**

# HTurbo

Starting point: DYTurbo

$q_{\perp}$  resummation implemented in numerical codes HqT and HRes [Catani et al.]  
Higher order accuracy require **high computation times**


Numerical code **DYTurbo** [Camarda et al. ATLAS collaboration, 1910.07049],  
fast and precise  $q_{\perp}$  resummation and several improvements for Drell-Yan  
( $h_1 + h_2 \rightarrow V + X \rightarrow l^+ l^- + X$ )

- **Goal:** set up a numerical code generalizing **DYTurbo** for Higgs boson production
- **Goal:** extend theoretical accuracy up to  $N^3\text{LL}+N^3\text{LO}$  and to other processes

Preliminary results to be seen in [Ferrera, Urtasun-Elizari.]

# Results


## Comparison with HRes and HqT - LL



- HTurbo  $q_{\perp}$  distribution vs HRes and HqT at LL
- Excellent numerical agreement up to the 0.1% level ✓

# Results


## Comparison with HTurbo and HqT - NLL



- HTurbo  $q_{\perp}$  distribution vs HRes and HqT at NLL
- Excellent numerical agreement up to the 0.1% level ✓

# Results


## Comparison with HRes and HqT - NNLL



- HTurbo  $q_{\perp}$  distribution vs HRes and HqT at NNLL
- Excellent numerical agreement up to the 0.1% level ✓

# Results

## Comparison HRes and HqT - all orders



- Higher orders lead to more accurate predictions ✓
- **HRes** needs 3 days to produce NNLL distribution → 3 minutes with **HTurbo!** ✓
- Agreement up to NNLL → ready for  $N^3$ LL



# Summary & Conclusions

- ① Precise knowledge of PDFs and partonic cross sections are required towards the precision era of the LHC
- ② Machine Learning models provide a robust way for PDFs determination optimized through **operator implementation in TF**
- ③ We develop a numerical code **HTurbo**, implementing  $q_\perp$  resummation for Higgs boson production, which is **faster than any of the existing codes**
- ④ Next steps:
  - Validate results at NNLO
  - Include full **N<sup>3</sup>LO** prediction
  - Perform phenomenological studies comparing with LHC data

Thank you!



This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 740006.