

QCD and Monte Carlo event generators

Monte Carlo course seminar - Milan, February 2021



UNIVERSITÀ
DEGLI STUDI
DI MILANO




European
Research
Council

- ① Hadron collisions and strong interactions
 - Hadron collisions and strong interactions
 - Renormalization group
 - QCD factorization
- ② MC and Parton Showers
 - Collinear factorization
 - Final state radiation
 - Initial state radiation
- ③ Hadronization: some basics
 - Large number of colors approximation
 - Hadronization models

Strong interactions

QCD from e^+e^- annihilation

Quantum Chromodynamics (QCD) \rightarrow theory describing the interaction between quarks and gluons (strong interactions)



QCD arises already from e^+e^- annihilation $\rightarrow R_0$ ratio

$$R_0 = \frac{\sigma(\gamma^* \rightarrow \text{hadrons})}{\sigma(\gamma^* \rightarrow \mu^+ \mu^-)} = 3 \sum_f c_f^2$$

- ❶ Color factor (3 color for each quark)
- ❷ Sum over charges of different flavors
- ❸ Threshold and higher order corrections

Strong interactions

Renormalization group


- Running coupling given by Renormalization Group Equation (RGE)

$$\mu \frac{d\alpha_s(\mu)}{d\mu} = \beta(\alpha_s(\mu)) = - \sum_{n=0}^{\infty} \beta_n \left(\frac{\alpha_s}{\pi} \right)^{n+1}$$

- Coupling α_s evolves with scale μ as given by RGE \rightarrow LO behavior driven by β_0
- $\beta_0^{\text{QCD}} > 0 \implies$ weakly coupled at large energies, asymptotic freedom
- $\beta_0^{\text{QED}} < 0 \implies$ strongly coupled at large energies, UV divergent!

Strong interactions

Renormalization group



- Running coupling given by Renormalization Group Equation (RGE)

$$\alpha_s(\mu) = \frac{1}{\beta_0 \log\left(\frac{\mu^2}{\Lambda_s^2}\right)}$$

- β_0 LO of the β function, is > 0
- Λ_s , parameter that defines value of the coupling at large scales


QCD is weakly coupled for $\mu \gg \Lambda_s \rightarrow$ asymptotically free

Perturbative Quantum Chromodynamics (pQCD)

Factorization theorem

QCD factorization

LHC processes $H_1 + H_2 \rightarrow F$



Separate process **PDFs** and **partonic (hard) interaction**

$$\sigma^F(p_1, p_2) = \sum_{\alpha, \beta} \int_0^1 dx_1 dx_2 f_{\alpha}(x_1, \mu_F^2) * f_{\beta}(x_2, \mu_F^2) * \hat{\sigma}_{\alpha\beta}^F(x_1 p_1, x_2 p_2, \alpha_s(\mu_R^2), \mu_F^2)$$

Parton showers

MC Parton showers

Partons in the initial and final state emit radiation. Initial state Radiation (ISR) and Final State Radiation (FSR) model by Monte Carlo (MC) shower algorithms


Shower Monte Carlo programs (HERWIG, PYTHIA)

- Libraries for computing SM and BSM cross sections
- Shower algorithms produce a number of enhanced coloured parton emissions to be added to the hard process
- Hadronization models, underlying event, decays of unstable hadrons, etc

Parton showers

Collinear limit

- QCD emission processes are enhanced in the collinear limit (θ small)
- σ dominated by collinear splittings $q \rightarrow qg, g \rightarrow gg, g \rightarrow q\bar{q}$




Collinear factorization \longrightarrow The cross section factorizes into the product of a tree-level cross section and a splitting factor out tree level amplitude and splitting

$$|M_{n+1}|^2 d\Phi_{n+1} \Rightarrow |M_n|^2 d\Phi_n \frac{\alpha_S}{2\pi} \frac{dt}{t} P_{q,qg}(z) dz \frac{d\phi}{2\pi}.$$

$$d\Phi_n = (2\pi)^4 \delta^4 \left(\sum_i^n k_i - q \right) \prod_i^n \frac{d^3 k_i}{(2\pi)^3 2k_i^0}$$

Parton showers

Kinematics of splitting



Kinematics of splitting given by (t, z, ϕ)

- t : parameter with dimensions of energy that vanish in the collinear limit
 - Virtuality $t = (k + l)^2 \approx z(1 - z)E^2\theta^2$
 - Transverse momentum $t = k_{\perp}^2 = l_{\perp}^2 \approx z^2(1 - z)^2E^2\theta^2$
 - Hardness $t = E^2\theta^2$
- z : fraction of energy of radiated parton $z = \frac{k^0}{k^0 + l^0}$
- ϕ represents azimuth of the k, l plane

Parton showers

AP splitting functions


Factorization holds for small angles \rightarrow small t variable

Difference in the splitting \rightarrow Altarelli-Parisi splitting functions (singular in $z \rightarrow 0, 1$)

$$P_{q,qg}(z) = C_F \frac{1+z^2}{1-z}$$

$$P_{g,gg}(z) = C_A \left(\frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right)$$

$$P_{g,q\bar{q}}(z) = T_F(z^2 + (1-z)^2)$$



We can proceed in an iterative way

$$|M_{n+2}|^2 d\Phi_{n+2} = |M_n|^2 d\Phi_n \frac{\alpha_s(t')}{2\pi} P_{q,qg}(z') \frac{dt'}{t'} dz' \frac{d\phi'}{2\pi} \frac{\alpha_s(t)}{2\pi} P_{q,qg}(z) \frac{dt}{t} dz \frac{d\phi}{2\pi}$$

Angles become small, maintaining a strong ordering relation $\theta' \gg \theta \rightarrow 0$

Parton showers

Exclusive final state

To describe exclusive final state \rightarrow sum perturbative expansion to all orders in α_s

$$\sigma_0 \alpha_s^n \int \frac{dt_1}{t_1} \dots \frac{dt_n}{t_n} \theta(Q^2 > t_1^2 > \dots > t_n^2 > \Lambda_S^2) = \sigma_0 \frac{\alpha_s^n}{n!} \log^n \left(\frac{Q^2}{\Lambda_S^2} \right)$$

Limit to the most singular terms, in ordered sequence of angles

Collinear approximation \rightarrow Leading log approximation

Final state radiation MC

General structure

Approximated description of a hadronic final state

Model a given hard scattering with arbitrary number of enhanced radiations

- Choose hard interaction with specified Born kinematics
- Consider all possible tree-level splittings for each coloured parton
- Assign the variables (t, z, ϕ) at each splitting vertex, t ordered in decreasing way
- At each splitting vertex assign the weight

$$\frac{\alpha_S(t)}{2\pi} P_{i,jl}(z) \frac{dt}{t} dz \frac{d\phi}{2\pi}$$

- Each line has a weight known as Sudakov factor

$$\Delta_i(t', t'') = \exp \left[- \sum_{(jl)} \int_{t''}^{t'} \frac{dt}{t} \int_0^1 dz \frac{\alpha_S(t)}{2\pi} P_{i,jl}(z) \right]$$

Final state radiation MC

Formal representation of a shower

Graphical notation for the representation of a shower

$$S_i(t, E) = \text{diagram of a blue circle with a line from the left labeled } i \text{ and horizontal lines on the right labeled } t, E,$$

Ensemble of all possible branchings from parton i at scale t

$$\text{diagram of a blue circle with a line from the left labeled } i \text{ and horizontal lines on the right labeled } t, E = \text{diagram of a green circle with a line from the left labeled } i \text{ and horizontal lines on the right labeled } t, t_0 + \text{diagram of a red circle with a line from the left labeled } i \text{ and horizontal lines on the right labeled } t, t', \text{ branching into two blue circles labeled } j \text{ and } l \text{ with horizontal lines labeled } t', zE \text{ and } t', (1-z)E$$

Forward evolution equation \rightarrow recursive structure

$$S_i(t, E) = \Delta_i(t, t_0) S_i(t_0, E) + \sum_{jl} \int_{t_0}^t \frac{dt'}{t'} \int_0^1 dz \int_0^{2\pi} \frac{d\phi}{2\pi} \frac{a_S(t')}{2\pi} \Delta_i(t', t_0) S_j(t', zE) S_l(t', (1-z)E)$$

Final state radiation MC

Probabilistic interpretation

$$\frac{\alpha_s(t')}{2\pi} P_{i,jl}(z') \frac{dt'}{t'} dz' \frac{d\phi'}{2\pi}$$

Probability of branching in the infinitesimal volume $dt' dz d\phi$

$$dP_{br} = \frac{\alpha_s(t')}{2\pi} \frac{dt'}{t'} \int_0^1 dz' P_{i,jl}(z') \int_0^{2\pi} \frac{d\phi'}{2\pi}$$

Probability of branching in the interval dt'

$$dP_{nobr} = 1 - dP_{br} = 1 - \frac{\alpha_s(t')}{2\pi} \frac{dt'}{t'} \int_0^1 dz' P_{i,jl}(z') \int_0^{2\pi} \frac{d\phi'}{2\pi}$$

Probability of no branching in the interval dt'

$$\Delta_i(t, t') = 1 - dP_{br} = \prod_i \left(1 - \frac{\alpha_s(t_i)}{2\pi} \frac{\delta t}{t_i} \int_0^1 dz' \int_0^{2\pi} P_{i,jl}(z') \frac{d\phi'}{2\pi} \right)$$

Sudakov form factor

$$dP_{fbr} = \Delta_i(t, t') \frac{\alpha_s(t')}{2\pi} P_{i,jl}(z') \frac{dt'}{t'} dz' \frac{d\phi'}{2\pi}$$

Probability of, starting at t , first branching in the phase space element $dt' dz d\phi$

Final state radiation MC

Shower algorithm

Generate hard process with probability proportional to its parton level cross section.
For each final state colored parton:

- ❶ Set scale $t = Q$, hard scale of the process.
- ❷ Generate random number $0 < r < 1$.
- ❸ Solve $r = \Delta_i(t, t')$ for t' .
- ❹ i) if $t' < t_0$, no further branching and stop shower.
- ❺ ii) if $t' \geq t_0$, generate j, l with energies

$$E_j = zE_i \quad \text{and} \quad E_l = (1 - z)E_i,$$


with z following a distribution given by $P_{i,jl}(z)$ and with azimuth ϕ uniformly distributed in the interval $[0, 2\pi]$.

- ❻ For each branched partons set $t = t'$ and start from (2).

Initial state radiation MC

General structure

- QCD coupling much larger than QED
- Coupling grows for small momentum transfer \rightarrow we can never neglect QCD ISR




Radiation from some initial state leading to some hard collision

ISR showers are spacelike \rightarrow initial parton is on shell

$$d\sigma_j^{\text{ISR}}(p, \dots) = \frac{\alpha_S}{2\pi} \frac{dt}{t} dz P_{ij}(z) d\sigma_i(zp, \dots)$$

Initial state radiation MC

Formal representation



$\delta x S_i(m, x, t, E)$ is ensemble of all possible states containing a spacelike parton m with energy between $x E$ and $(x + \delta x E)$, scale t

$$S_i(m, x, t, E) = \frac{t_0, E}{i} \text{S} \begin{array}{c} m, t, xE \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

- Lines between t_1 and t_2 , t_2 and hard scattering are spacelike
- Difference from FSR in Sudakov factors and Splitting functions arise at NLO

Forward evolution equation


$$\frac{t_0, E}{i} \text{S} \begin{array}{c} m, t, xE \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \frac{t_0}{i} \frac{t}{\delta_{im} \delta(1-x)} \begin{array}{c} t', zE \\ \text{S} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} j \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} t', (1-z)E \\ \text{S} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} + \frac{t_0}{i} \frac{t'}{i} \begin{array}{c} t', zE \\ \text{S} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} j \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} t', (1-z)E \\ \text{S} \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Initial state radiation MC

Backwards evolution equation

Great amount of computation time to generate configurations leading to the hard scattering that we want

Moderns MC programs \rightarrow recursive procedure starting at the large scale



$$\sum_{\mathcal{F}} S_i(m, t, x, E) = f_m^{(i)}(x, t)$$

- The blob l at the splitting vertex is given by the inclusive splitting kernel P_{jm}
- Backwards evolution equation (scale dependent parton density)

$$f_m^{(i)}(x, t) = \delta_{im} \delta(1-x) \Delta_m(t, t_0) + \int_{t_0}^t \frac{dt'}{t'} \int_x^1 \frac{dz}{z} \sum_j f_j^{(i)}(z, t') \frac{\alpha_S(t')}{2\pi} \hat{P}_{jm} \left(\frac{x}{z} \right) \Delta_m(t, t')$$

Initial state radiation MC

Shower algorithm

Generate hard process with probability proportional to its parton level cross section. For each final state colored parton:

- 1 Set scale $t = Q$, hard scale of the process
- 2 Generate random number $0 < r < 1$
- 3 Solve

$$r = \frac{f_i^{(h)}(t', x) \Delta_i(t, t')}{f_i^{(h)}(t, x)} \quad \text{for } t'$$

- 4 i) if $t' < t_0$, no further branching and stop shower
- 5 ii) if $t' \geq t_0$, generate j, l with energies

$$E_j = zE_i \quad \text{and} \quad E_l = (1 - z)E_i,$$

with z following a distribution given by $P_{i,jl}(z)$ and with azimuth ϕ uniformly distributed in the interval $[0, 2\pi]$


- 6 For parton j set $t = t'$ and start from (2). For parton l generate a timelike parton shower according to the algorithm shown previously

Hadronization

Some basics

A parton becomes a measurable hadron through the emission of a partonic shower
Large number of color approximation

- Color indices ranging from 1 to N_c and keep only dominant contribution
- Each parton identified by a unique label



Hadronization

Some basics

Hadronization models

- Cluster models
 - Gluons are forced to split in quark-antiquark pair
 - Then decay each color connected quark-antiquark pair independently
 - Preconfinement, assuming subsystems of color singlet partons with universal invariant mass distribution (power suppressed at high masses)
- Lund string model
 - Color connected partons collected in a system consisting of a quark, several intermediate gluons, and an antiquark
 - Intermediate gluons are transverse kicks of a continuum medium

Fragmentation models are one of the most complex aspects of Shower Monte Carlo
These models have unavoidably a large number of parameters

Summary

- LHC processes factorization in perturbative and non-perturbative components
- Perturbative QCD applied only at high energies
- Monte Carlo shower algorithms describe the enhanced emissions produced by initial and final state partons
- Hadronization models describing the non-perturbative physics