

EE2023 Signals & Systems Quiz

Semester 1 AY2014/15

Date : 2 October 2014

Time Allowed : 1.5 hours

Instructions :

1. Answer all 4 questions. Each question carries 10 marks.
2. This is a closed book quiz.
3. Tables of Fourier transforms and trigonometric identities are given on Pages 11 and 12.
4. No programmable or graphic calculator is allowed.
5. Write your answers in the spaces indicated in this question paper. Attachment is not allowed.
6. Write your name, matric number and lecture group in the spaces indicated below.

Name : _____

Matric # : _____

Class Group # : _____

For your information :

Group 1 : A/Prof Loh Ai Poh

Group 2 : A/Prof Ng Chun Sum

Group 3 : A/Prof Tan Woei Wan

Group 4 : Prof Lawrence Wong

Question #	Marks
1	
2	
3	
4	
Total Marks	

Q.1 ANSWER ~ continued

[illegible]

Q.2 A signal $x(t)$ is modeled by $x(t) = \sum_n \text{sinc}(t - 3n)$.

- (a) Express $x(t)$ in the form of a convolution and then derive its Fourier transform $X(f)$. (4 marks)

- (b) Find the inverse Fourier transform of $X(f)$.

[Your answer for this part should not contain the summation \sum operator.]

- (3 marks)

- (c) Using the result of Part (b), or otherwise, determine the Fourier series coefficients, X_k , of $x(t)$.

- (3 marks)

Q.2 ANSWER

[illegible]

Q.2 ANSWER ~ continued

[illegible]

Q.3 ANSWER ~ continued

[illegible]

Q.4. (a) Determine the Fourier transform of the signal $x(t)$ that comprises two rectified cycles of a sinusoid as shown in Figure Q.4(a).




Figure Q.4(a)

(4 marks)

(b) Using the Dirac- δ replication property, derive an expression that shows the relationship between $x(t)$ and the periodic signal $x_p(t)$ shown in Figure Q.4(b).




Figure Q.4(b)

(2 marks)

(c) Determine the Fourier transform and the Fourier Series coefficients of the periodic signal $x_p(t)$.

(4 marks)

Q.4 ANSWER

[illegible]

Q.4 ANSWER ~ continued

[illegible]

This page is intentionally left blank to facilitate detachment of the formula sheet for easy reference. Anything written on this page will not be graded.

Fourier Series:
$$\begin{cases} X_k = \frac{1}{T} \int_{\tilde{t}}^{\tilde{t}+T} x(t) \exp(-j2\pi k t/T) dt \\ x(t) = \sum_{k=-\infty}^{\infty} X_k \exp(j2\pi k t/T) \end{cases}$$

Fourier Transform:
$$\begin{cases} X(f) = \int_{-\infty}^{\infty} x(t) \exp(-j2\pi f t) dt \\ x(t) = \int_{-\infty}^{\infty} X(f) \exp(j2\pi f t) df \end{cases}$$

FOURIER TRANSFORMS OF BASIC FUNCTIONS		
	$x(t)$	$X(f)$
Constant	K	$K\delta(f)$
Unit Impulse	$\delta(t)$	1
Unit Step	$u(t)$	$\frac{1}{2} \left[\delta(f) + \frac{1}{j\pi f} \right]$
Sign (or Signum)	$\text{sgn}(t)$	$\frac{1}{j\pi f}$
Rectangle	$\text{rect}\left(\frac{t}{T}\right)$	$T \text{sinc}(fT)$
Triangle	$\text{tri}\left(\frac{t}{T}\right)$	$T \text{sinc}^2(fT)$
Sine Cardinal	$\text{sinc}\left(\frac{t}{T}\right)$	$T \text{rect}(fT)$
Complex Exponential	$\exp(j2\pi f_o t)$	$\delta(f - f_o)$
Cosine	$\cos(2\pi f_o t)$	$\frac{1}{2} [\delta(f - f_o) + \delta(f + f_o)]$
Sine	$\sin(2\pi f_o t)$	$-\frac{j}{2} [\delta(f - f_o) - \delta(f + f_o)]$
Gaussian	$\exp\left(-\frac{t^2}{\alpha^2}\right)$	$\alpha \pi^{0.5} \exp(-\alpha^2 \pi^2 f^2)$
Comb	$\sum_{m=-\infty}^{\infty} \delta(t - mT)$	$\frac{1}{T} \sum_{k=-\infty}^{\infty} \delta\left(f - \frac{k}{T}\right)$

FOURIER TRANSFORM PROPERTIES		
	Time-domain	Frequency-domain
Linearity	$\alpha x_1(t) + \beta x_2(t)$	$\alpha X_1(f) + \beta X_2(f)$
Time scaling	$x(\beta t)$	$\frac{1}{ \beta } X\left(\frac{f}{\beta}\right)$
Duality	$X(t)$	$x(-f)$
Time shifting	$x(t - t_o)$	$X(f) \exp(-j2\pi f t_o)$
Frequency shifting (Modulation)	$x(t) \exp(j2\pi f_o t)$	$X(f - f_o)$
Differentiation in the time-domain	$\frac{d^n}{dt^n} x(t)$	$(j2\pi f)^n X(f)$
Multiplication in the time-domain	$x_1(t) x_2(t)$	$\int_{-\infty}^{\infty} X_1(\zeta) X_2(f - \zeta) d\zeta$ or $X_1(f) * X_2(f)$
Convolution in the time-domain	$\int_{-\infty}^{\infty} x_1(\zeta) x_2(t - \zeta) d\zeta$ or $x_1(t) * x_2(t)$	$X_1(f) X_2(f)$
Integration in the time-domain	$\int_{-\infty}^t x(\tau) d\tau$	$\frac{1}{j2\pi f} X(f) + \frac{1}{2} X(0) \delta(f)$
		$\frac{1}{j2\pi f} X(f)$ if $X(0) = 0$

TRIGONOMETRIC IDENTITIES	
$\exp(\pm j\theta) = \cos(\theta) \pm j \sin(\theta)$	$\sin(\alpha \pm \beta) = \sin(\alpha)\cos(\beta) \pm \cos(\alpha)\sin(\beta)$
$\cos(\theta) = \frac{1}{2}[\exp(j\theta) + \exp(-j\theta)]$	$\cos(\alpha \pm \beta) = \cos(\alpha)\cos(\beta) \mp \sin(\alpha)\sin(\beta)$
$\sin(\theta) = \frac{1}{j2}[\exp(j\theta) - \exp(-j\theta)]$	$\tan(\alpha \pm \beta) = \frac{\tan(\alpha) \pm \tan(\beta)}{1 \mp \tan(\alpha)\tan(\beta)}$
$\sin^2(\theta) + \cos^2(\theta) = 1$	
$\sin(2\theta) = 2\sin(\theta)\cos(\theta)$	$\sin(\alpha)\sin(\beta) = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$
$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta)$	$\cos(\alpha)\cos(\beta) = \frac{1}{2}[\cos(\alpha - \beta) + \cos(\alpha + \beta)]$
$\sin^2(\theta) = \frac{1}{2}[1 - \cos(2\theta)]$	$\sin(\alpha)\cos(\beta) = \frac{1}{2}[\sin(\alpha - \beta) + \sin(\alpha + \beta)]$
$\cos^2(\theta) = \frac{1}{2}[1 + \cos(2\theta)]$	$\mathbf{C}\cos(\theta) - \mathbf{S}\sin(\theta) = \sqrt{\mathbf{C}^2 + \mathbf{S}^2} \cos\left[\theta + \tan^{-1}\left(\frac{\mathbf{S}}{\mathbf{C}}\right)\right]$