

EE2023 TUTORIAL 2 (SOLUTIONS)

Solution to Q.1

- (a) $x(t)$ has an average (or DC) value of 2 ∴ Zero-frequency component has value 2
- (b) $x(t)$ is a POWER SIGNAL ∴ $\begin{cases} \text{Spectrum is defined only at discrete} \\ \text{frequency points (sum of sinusoids)} \end{cases}$
- (c) $x(t)$ is APERIODIC ∴ $\{\pi, \pi^2, \pi^3\} \dots$ has no common factor
- Therefore, $x(t)$ does not have a Fourier series expansion.
-

Solution to Q.2

- (a) $x(t) = \cos(3.2t) + \sin(1.6t) + \exp(j2.8t) \dots \begin{cases} \cos(3.2t) & \text{has a frequency of } 3.2 \text{ rad/s} \\ \sin(1.6t) & \text{has a frequency of } 1.6 \text{ rad/s} \\ \exp(j2.8t) & \text{has a frequency of } 2.8 \text{ rad/s} \end{cases}$

Highest common factor (HCF) of $\{3.2, 1.6, 2.8\}$ exists and is equal to 0.4. Thus, $x(t)$ is periodic with a fundamental frequency of $\omega_p = 0.4 \text{ rad/s}$. The period of $x(t)$ is $T_p = \frac{2\pi}{0.4} = 5\pi \text{ s}$.

REMARKS: Although $x(t)$ is periodic with a fundamental frequency of 0.4 rad/s , it does not contain the fundamental frequency component itself.

- (b) $x(t) = \cos(4t) + \sin(\pi t) \dots \begin{cases} \cos(4t) & \text{has a frequency of } 4 \text{ rad/s} \\ \sin(\pi t) & \text{has a frequency of } \pi \text{ rad/s} \end{cases}$

The set of frequencies $\{4, \pi\}$ has no common factor. Thus, $x(t)$ is not periodic.

REMARKS: Summing sinusoids does not necessarily lead to a periodic signal unless the frequencies of the sinusoids are harmonics of a common fundamental frequency.

Solution to Q.3

- (a) The fundamental frequency of $x(t) = 6\sin(12\pi t) + 4\exp\left(j\left(8\pi t + \frac{\pi}{4}\right)\right) + 2$ is $\begin{cases} f_p = \text{HCF}\{6, 4\} = 2 \\ T_p = 0.5 \end{cases}$.

Re-write $x(t)$ as a sum of weighted zero-phase complex exponentials and arrange the terms in ascending frequency order:


$$\begin{aligned} x(t) &= \frac{6}{j2} [\exp(j12\pi t) - \exp(-j12\pi t)] + 4\exp(j\pi/4)\exp(j8\pi t) + 2 \\ &= j3\exp(-j12\pi t) + 2 + 4\exp(j\pi/4)\exp(j8\pi t) - j3\exp(j12\pi t) \end{aligned} \quad (1)$$

Express $x(t)$ as a complex exponential Fourier series:

$$\begin{aligned} x(t) &= \sum_{k=-\infty}^{\infty} c_k \exp\left(j2\pi \frac{k}{T_p} t\right) = \sum_{k=-\infty}^{\infty} c_k \exp(j4\pi k t) \\ &= \left(\begin{aligned} &\cdots + c_{-3}\exp(-j12\pi t) + c_{-2}\exp(-j8\pi t) + c_{-1}\exp(-j4\pi t) \\ &\quad + c_0 \\ &+ c_1\exp(j4\pi t) + c_2\exp(j8\pi t) + c_3\exp(j12\pi t) + \cdots \end{aligned} \right) \end{aligned} \quad (2)$$


Comparing coefficients of complex exponential terms in (1) and (2), we conclude that:

$$c_{-3} = j3, \quad c_0 = 2, \quad c_2 = 4\exp(j\pi/4), \quad c_3 = -j3 \quad \text{and} \quad [c_k = 0; k \neq 0, 2, \pm 3].$$



Remarks: If a periodic signal is given as a sum of sinusoids, then its Fourier series coefficients can be evaluated using the above method without the need to perform any integration.

- (b) $x(t) = \frac{1}{2}(|\sin(\pi t)| + \sin(\pi t))$: Half-wave rectification of $\sin(\pi t)$.




Period: $T = 2$ s

Fundamental frequency: $f_1 = \frac{1}{T} = 0.5$ Hz

4th harmonic frequency: $f_4 = 4f_1 = 2$ Hz

Solution to Q.4

Graphically, we observe that $x(t) = \sum_{n=-\infty}^{\infty} 2p(t-1.6n) = 1.8$.




By Deduction:


- $x(t)$ has a zero-frequency component of value 1.8, which implies that $c_0 = 1.8$.
- $x(t)$ has no non-zero frequency components, which implies that $c_k = 0$; $k \neq 0$.

By Derivation:

Since $x(t)$ is a constant (or a DC signal), it may be treated as a periodic signal of arbitrary period T , where $0 < T < \infty$. Its Fourier series coefficients can thus be computed as


$$\begin{aligned} c_k &= \frac{1}{T} \int_{-T/2}^{T/2} 1.8 \exp\left(-j2\pi \frac{k}{T} t\right) dt = \frac{1.8}{T} \left[\frac{\exp(-j2\pi kt/T)}{-j2\pi k/T} \right]_{-T/2}^{T/2} \\ &= \frac{1.8}{T} \left[\frac{\exp(-j\pi k)}{-j2\pi k/T} - \frac{\exp(j\pi k)}{-j2\pi k/T} \right] \\ &= 1.8 \frac{\sin(\pi k)}{\pi k} = 1.8 \operatorname{sinc}(k) = \begin{cases} 1.8; & k = 0 \\ 0; & k \neq 0 \end{cases} \end{aligned}$$



Solution to Q.5

Solution to S.1

(a)



(b)

$$\begin{aligned}
 c_k &= \frac{1}{1} \int_0^1 \sin(\pi t) \exp(-j2\pi kt) dt \\
 &= \frac{1}{j2} \int_0^1 [\exp(j\pi t) - \exp(-j\pi t)] \exp(-j2\pi kt) dt \\
 &= \frac{1}{j2} \int_0^1 \exp[j\pi(1-2k)t] - \exp[-j\pi(1+2k)t] dt \\
 &= \frac{1}{j2} \left[\frac{\exp[j\pi(1-2k)t]}{j\pi(1-2k)} - \frac{\exp[-j\pi(1+2k)t]}{-j\pi(1+2k)} \right]_0^1 \\
 &= \frac{1}{j2} \left[\frac{\exp[j\pi(1-2k)] - 1}{j\pi(1-2k)} - \frac{\exp[-j\pi(1+2k)] - 1}{-j\pi(1+2k)} \right] \\
 &= \frac{1}{\pi(2k+1)} - \frac{1}{\pi(2k-1)} = -\frac{2}{\pi} \cdot \frac{1}{4k^2 - 1}
 \end{aligned}$$

$$x(t) = \sum_{k=-\infty}^{\infty} c_k \exp(j2\pi kt) = -\frac{2}{\pi} \sum_{k=-\infty}^{\infty} \frac{1}{4k^2 - 1} \exp(j2\pi kt)$$

(c)

$$a_k = \frac{c_{-k} + c_k}{2} = -\frac{2}{\pi} \cdot \frac{1}{4k^2 - 1}$$

$$b_k = \frac{c_{-k} - c_k}{j2} = 0$$

$$x(t) = a_0 + 2 \sum_{k=1}^{\infty} a_k \cos(2\pi kt) + b_k \sin(2\pi kt) = \frac{2}{\pi} - \frac{4}{\pi} \sum_{k=1}^{\infty} \frac{1}{4k^2 - 1} \cos(2\pi kt)$$

Solution to S.2

$$\begin{aligned}
c_k &= \frac{1}{2\pi} \int_{-\pi}^{\pi} t^2 \exp(-jkt) dt = \frac{1}{2\pi} \left(\left[t^2 \frac{\exp(-jkt)}{-jk} \right]_{-\pi}^{\pi} - \int_{-\pi}^{\pi} 2t \frac{\exp(-jkt)}{-jk} dt \right) \\
&= \frac{1}{2\pi} \left(\left[t^2 \frac{\exp(-jkt)}{-jk} \right]_{-\pi}^{\pi} - \left[2t \frac{\exp(-jkt)}{(-jk)^2} \right]_{-\pi}^{\pi} + \int_{-\pi}^{\pi} 2 \frac{\exp(-jkt)}{(-jk)^2} dt \right) \\
&= \frac{1}{2\pi} \left(\left[t^2 \frac{\exp(-jkt)}{-jk} \right]_{-\pi}^{\pi} - \left[2t \frac{\exp(-jkt)}{(-jk)^2} \right]_{-\pi}^{\pi} + \left[2 \frac{\exp(-jkt)}{(-jk)^3} \right]_{-\pi}^{\pi} \right) \\
&= \pi \left[\frac{\sin(\pi k)}{k} \right] + \left[\frac{2 \cos(\pi k)}{k^2} \right] - \frac{2}{\pi} \left[\frac{\sin(\pi k)}{k^3} \right] \\
&= \pi^2 \left[\frac{\sin(\pi k)}{\pi k} \right] + \left[\frac{2\pi k \cos(\pi k) - 2 \sin(\pi k)}{\pi k^3} \right] = \begin{cases} 2(-1)^k / k^2; & k \neq 0 \\ \pi^2 / 3; & k = 0 \end{cases}
\end{aligned}$$

$$x(t) = \sum_{k=-\infty}^{\infty} c_k \exp(jkt) = \frac{\pi^2}{3} + \sum_{\substack{k=-\infty \\ k \neq 0}}^{\infty} \frac{2(-1)^k}{k^2} \exp(jkt) = \frac{\pi^2}{3} + 4 \sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} \cos(kt)$$

Solution to S.3

$$\begin{aligned}
x(t) &= h_0 + \sum_{k=1}^{\infty} h_k \cos\left(2\pi \frac{k}{T_0} t - \theta_k\right) \\
&= \underbrace{h_0}_{a_0} + 2 \sum_{k=1}^{\infty} \underbrace{0.5 h_k \cos(\theta_k)}_{a_k} \cos\left(2\pi \frac{k}{T_0} t\right) + \underbrace{0.5 h_k \sin(\theta_k)}_{b_k} \sin\left(2\pi \frac{k}{T_0} t\right)
\end{aligned}$$

$$\left. \begin{aligned} c_{-k} &= c_k^* \quad (\because x(t) \text{ is real}) \\ a_k &= \frac{c_{-k} + c_k}{2} = \operatorname{Re}[c_k] \\ b_k &= \frac{c_{-k} - c_k}{j2} = -\operatorname{Im}[c_k] \end{aligned} \right\} \rightarrow \begin{cases} h_0 = a_0 = c_0 \\ h_k = 2\sqrt{a_k^2 + b_k^2} = \sqrt{\operatorname{Re}^2[c_k] + \operatorname{Im}^2[c_k]} = 2|c_k| \\ \theta_k = \tan^{-1}\left(\frac{b_k}{a_k}\right) = -\tan^{-1}\left(\frac{\operatorname{Im}[c_k]}{\operatorname{Re}[c_k]}\right) \end{cases}$$
