ST2334 (2020/2021 Semester 1) Solutions to Questions in Tutorial 7

Question 1

$f_{X, Y}(x, y)$		3	r	
		2	4	$f_{y}(y)$
	1	0.10	0.15	0.25
у	3	0.20	0.30	0.50
	5	0.10	0.15	0.25
$f_x(x)$		0.40	0.60	1

(a) It can be verified directly that

$$f_{X,Y}(x,y) = f_X(x)f_Y(y)$$
 for $x = 2, 4$ and $y = 1, 3, 5$.

Hence, *X* and *Y* are independent

(b)

y 1		3	5		
$f_{Y X}(y 2)$	0.10/0.40 = 1/4	0.20/0.40 = 2/4	0.10/0.40 = 1/4		

$$E(Y|X=2) = 1(1/4) + 3(2/4) + 5(1/4) = 3.$$

Alternatively, as X and Y are independent, then E(Y|X=2)=E(Y)=1(0.25)+3(0.5)+5(0.25)=3.

(c)

X	2	4			
$f_{X Y}(x 3)$	0.20/0.50 = 2/5	0.30/0.50 = 3/5			

$$E(X|Y=3) = 2(2/5) + 4(3/5) = 16/5 = 3.2$$

Alternatively, as X and Y are independent, then E(X|Y=3)=E(X)=2(0.4)+4(0.6)=3.2.

- (d) E(X) = 2(0.40) + 4(0.60) = 3.2 E(Y) = 1(0.25) + 3(0.50) + 5(0.25) = 3E(2X - 3Y) = 2E(X) - 3E(Y) = 2(3.2) - 3(3) = -2.6. We used the results in parts (b) and (c) in the computation.
- (e) E(XY) = (2)(1)(0.10) + (2)(3)(0.10) + (2)(5)(0.10) + (4)(1)(0.10) + (4)(3)(0.10) + (4)(5)(0.10) = 9.6.

Alternatively, as X and Y are independent, then E(XY) = E(X)E(Y) = 3.2(3) = 9.6.

(f) $E(X^2) = 2^2(0.4) + 4^2(0.6) = 11.2$. Hence, $V(X) = E(X^2) - [E(X)]^2 = 11.2 - (3.2)^2 = 0.96$

Similarly, $E(Y^2) = 1^2(0.25) + 3^2(0.5) + 5^2(0.25) = 11$. Hence, $V(Y) = E(Y^2) - [E(Y)]^2 = 11 - (3)^2 = 2$

(g) $\sigma_{X,Y} = 0$ as X and Y are independent.

Alternatively, Cov(X, Y) = E(XY) - E(X)E(Y) = 9.6 - 3.2(3) = 0 $\rho_{XY} = 0$ as $\sigma_{XY} = 0$. Alternatively, $\rho_{XY} = 0$ as X and Y are independent.

Question 2

By the definition of X and Y, Profit = 8X + 3Y - 10; and the marginal distributions of X and Y are given in the following table.

$f_{X, Y}(x, y)$					
		0	1	2	$f_{y}(y)$
	0	0.01	0.01	0.03	0.05
y	1	0.03	0.08	0.07	0.18
2		0.03	0.06	0.06	0.15
	3	0.07	0.07	0.13	0.27
	4	0.12	0.04	0.03	0.19
	5	0.08	0.06	0.02	0.16
$f_x(x)$		0.34	0.32	0.34	1

$$E(X) = \sum x f_X(x) = 1. \ E(X^2) = \sum x^2 f_X(x) = 1.68. \ V(X) = E(X^2) - [E(X)]^2 = 0.68.$$

$$E(Y) = \sum y f_Y(y) = 2.85. \ E(Y^2) = \sum y^2 f_Y(y) = 10.25. \ V(Y) = E(Y^2) - [E(Y)]^2 = 2.1275.$$

$$\sigma_{X,Y} = E(XY) - E(X)E(Y) = (2.47) - (1)(2.85) = -0.38$$

Profit = $8X + 3Y - 10$. $E(Profit) = 8E(X) + 3E(Y) - 10 = 6.55$
 $V(profit) = V[8X + 3Y - 10] = 8^2V(X) + 3^2E(Y) + 2(8)(3)Cov(X,Y) = 44.4275$.

Question 3

We first compute the marginal probability density functions of X and Y. They are also needed for parts (b) to (d).

$$f_X(x) = \int_0^1 \frac{2}{3}(x+2y) \, dy = \frac{2}{3} \left[xy + \frac{2y^2}{2} \right]_0^1 = \frac{2}{3}(x+1)$$
, for $0 \le x \le 1$; and 0 otherwise $f_Y(y) = \int_0^1 \frac{2}{3}(x+2y) \, dx = \frac{2}{3} \left[\frac{x^2}{2} + 2xy \right]_0^1 = \frac{2}{3} \left(\frac{1}{2} + 2y \right)$, for $0 \le y \le 1$; and 0 otherwise,

(a) Since $f_{X,Y}(x,y) \neq f_X(x)f_Y(y)$, X and Y are dependent.

(b)
$$E(X) = \frac{2}{3} \int_0^1 x(x+1) dx = \frac{2}{3} \left[\frac{x^3}{3} + \frac{x^2}{2} \right]_0^1 = \frac{2}{3} \left(\frac{1}{3} + \frac{1}{2} \right) = \frac{5}{9} = 0.55556.$$

 $E(X^2) = \frac{2}{3} \int_0^1 x^2(x+1) dx = \frac{2}{3} \left[\frac{x^4}{4} + \frac{x^3}{3} \right]_0^1 = \frac{2}{3} \left(\frac{1}{4} + \frac{1}{3} \right) = \frac{7}{18} = 0.38889.$
 $V(X) = E(X^2) - [E(X)]^2 = 13/162 = 0.08024.$

(c)
$$E(Y) = \frac{2}{3} \int_0^1 y \left(\frac{1}{2} + 2y\right) dy = \frac{2}{3} \left[\frac{y^2}{4} + \frac{2y^3}{3}\right]_0^1 = \frac{2}{3} \left(\frac{1}{4} + \frac{2}{3}\right) = \frac{11}{18} = 0.61111.$$

$$E(Y^2) = \frac{2}{3} \int_0^1 y^2 \left(\frac{1}{2} + 2y\right) dx = \frac{2}{3} \left[\frac{y^3}{6} + \frac{y^4}{2}\right]_0^1 = \frac{2}{3} \left(\frac{1}{6} + \frac{1}{2}\right) = \frac{4}{9} = 0.44444.$$

$$V(Y) = E(Y^2) - [E(Y)]^2 = 23/324 = 0.07099.$$

(d)
$$E(XY) = \frac{2}{3} \int_0^1 \int_0^1 xy(x+2y) \, dx dy = \frac{2}{3} \int_0^1 \int_0^1 x^2 y + 2xy^2 \, dx dy$$

 $= \frac{2}{3} \int_0^1 \left[\frac{x^3 y}{3} + \frac{2x^2 y^2}{2} \right]_0^1 \, dy = \frac{2}{3} \int_0^1 \left(\frac{y}{3} + y^2 \right) \, dy = \frac{2}{3} \left[\frac{y^2}{6} + \frac{y^3}{3} \right]_0^1 = \frac{2}{3} \left(\frac{1}{6} + \frac{1}{3} \right) = \frac{1}{3} = 0.333333.$
Hence, $\sigma_{X,Y} = E(XY) - E(X)E(Y) = 1/3 - (5/9)(11/18) = -1/162 = -0.00617.$

Question 4

$$f_X(x) = \int_0^1 \frac{3}{2} (x^2 + y^2) dy = \frac{3}{2} \left[x^2 y + \frac{y^3}{3} \right]_0^1 = \frac{3}{2} (x^2 + \frac{1}{3})$$
, for $0 \le x \le 1$; and 0 otherwise.

$$f_Y(y) = \int_0^1 \frac{3}{2} (x^2 + y^2) dx = \frac{3}{2} \left[\frac{x^3}{3} + xy^2 \right]_0^1 = \frac{3}{2} \left(\frac{1}{3} + y^2 \right)$$
, for $0 \le y \le 1$; and 0 otherwise.

(a) Since $f_{X,Y}(x,y) \neq f_X(x)f_Y(y)$, X and Y are dependent

(b)
$$E(X) = \frac{3}{2} \int_0^1 x \left(x^2 + \frac{1}{3} \right) dx = \frac{3}{2} \left[\frac{x^4}{4} + \frac{x^2}{6} \right]_0^1 = \frac{3}{2} \left(\frac{1}{4} + \frac{1}{6} \right) = \frac{5}{8} = 0.625.$$

 $E(X^2) = \frac{3}{2} \int_0^1 x^2 \left(x^2 + \frac{1}{3} \right) dx = \frac{3}{2} \left[\frac{x^5}{5} + \frac{x^3}{9} \right]_0^1 = \frac{3}{2} \left(\frac{1}{5} + \frac{1}{9} \right) = \frac{7}{15} = 0.46667.$
 $V(X) = E(X^2) - [E(X)]^2 = 73/960 = 0.07604.$

(c) Method 1: Repeat the work in part (b)

$$E(Y) = \frac{3}{2} \int_0^1 y \left(y^2 + \frac{1}{3} \right) dy = \frac{3}{2} \left[\frac{y^4}{4} + \frac{y^2}{6} \right]_0^1 = \frac{3}{2} \left(\frac{1}{4} + \frac{1}{6} \right) = \frac{5}{8} = 0.625.$$

$$E(Y^2) = \frac{3}{2} \int_0^1 y^2 \left(y^2 + \frac{1}{3} \right) dy = \frac{3}{2} \left[\frac{y^5}{5} + \frac{y^3}{9} \right]_0^1 = \frac{3}{2} \left(\frac{1}{5} + \frac{1}{9} \right) = \frac{7}{15} = 0.46667.$$

$$V(Y) = E(Y^2) - [E(Y)]^2 = 73/960 = 0.07604.$$

Method 2: Observe that the joint p.d.f. is symmetric in x and y (i.e. $f_{(X,Y)}(x,y) = f_{X,Y}(y,x)$), so E(Y) = E(X) = 5/8 = 0.625 and V(Y) = V(X) = 73/960 = 0.07604.

(d)
$$E(XY) = \frac{3}{2} \int_0^1 \int_0^1 xy(x^2 + y^2) dxdy = \frac{3}{2} \int_0^1 \int_0^1 x^3 y + xy^3 dxdy$$

 $= \frac{3}{2} \int_0^1 \left[\frac{x^4 y}{4} + \frac{x^2 y^3}{2} \right]_0^1 dy = \frac{3}{2} \int_0^1 \left(\frac{y}{4} + \frac{y^3}{2} \right) dy = \frac{3}{2} \left[\frac{y^2}{8} + \frac{y^4}{8} \right]_0^1 = \frac{3}{2} \left(\frac{1}{8} + \frac{1}{8} \right) = \frac{3}{8} = 0.375.$
 $\sigma_{X,Y} = E(XY) - E(X)E(Y) = (3/8) - (5/8)(5/8) = -1/64 = -0.01563.$

- (e) E(X + Y) = E(X) + E(Y) = 5/8 + 5/8 = 5/4 = 1.25
- (f) $V(X+Y) = V(X) + V(Y) + 2\left(\sigma_{X,Y}\right) = \frac{73}{960} + \frac{73}{960} + 2\left(\frac{-1}{64}\right) = \frac{29}{240} = 0.12083.$

Question 5

(a)
$$f_X(x) = \int_0^1 (x+y) \, dy = \left[xy + \frac{y^2}{2} \right]_0^1 = x + \frac{1}{2}, \text{ for } 0 \le x \le 1; \text{ and } f_X(x) = 0 \text{ otherwise.}$$

$$E(X) = \int_0^1 x \left(x + \frac{1}{2} \right) \, dx = \left[\frac{x^3}{3} + \frac{x^2}{4} \right]_0^1 = \left(\frac{1}{3} + \frac{1}{4} \right) = \frac{7}{12} = 0.58333.$$

$$f_Y(y) = \int_0^1 (x+y) \, dx = \left[\frac{x^2}{2} + xy \right]_0^1 = y + \frac{1}{2}, \text{ for } 0 \le y \le 1; \text{ and } f_Y(y) = 0 \text{ otherwise.}$$

$$E(Y) = \int_0^1 y \left(y + \frac{1}{2} \right) \, dy = \left[\frac{y^3}{3} + \frac{y^2}{4} \right]_0^1 = \left(\frac{1}{3} + \frac{1}{4} \right) = \frac{7}{12} = 0.58333.$$

$$E(XY) = \int_0^1 \int_0^1 xy(x+y,) \, dxdy = \int_0^1 \int_0^1 x^2y + xy^2 \, dxdy = \int_0^1 \left[\frac{x^3y}{3} + \frac{x^2y^2}{2} \right]_0^1 \, dy$$

$$= \int_0^1 \left(\frac{y}{3} + \frac{y^2}{2} \right) \, dy = \left[\frac{y^2}{6} + \frac{y^3}{6} \right]_0^1 = \frac{1}{3} = 0.33333.$$

$$\sigma_{X,Y} = E(XY) - E(X)E(Y) = 1/3 - (7/12)(7/12) = -1/144 = -0.00694.$$

(b) To compute E(Y|X=0.2), we need to compute $f_{Y|X}(y|0.2)$. We shall restrict ourselves to $0 \le y \le 1$ because the conditional probability density function is 0 for $y \notin [0,1]$. We have $f_{Y|X}(y|x=0.2) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \frac{x+y}{x+1/2} = \frac{0.2+y}{0.7} = \frac{2+10y}{7}$, for $0 \le y \le 1$; and $f_{Y|X}(y|x=0.2) = 0$ otherwise.

$$E(Y|X=0.2) = \int_0^1 y\left(\frac{2+10y}{7}\right) dy = \frac{1}{7} \left[y^2 + \frac{10y^3}{3}\right]_0^1 = \frac{1}{7} \left(1 + \frac{10}{3}\right) = \frac{13}{21} = 0.61905.$$

(c) Similarly, $f_{X|Y}(x|y=0.5) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{x+y}{y+1/2} = \frac{0.5+x}{1} = x + \frac{1}{2}$, for $0 \le x \le 1$; and $f_{X|Y}(x|y=0.5) = 0$ otherwise.

$$E(X|Y=0.5) = \int_0^1 x \left(x + \frac{1}{2}\right) dx = \left[\frac{x^3}{3} + \frac{x^2}{4}\right]_0^1 = \left(\frac{1}{3} + \frac{1}{4}\right) = \frac{7}{12} = 0.58333.$$

Question 6

$$V(Z) = V(-2X + 4Y) = (-2)^2 V(X) + 4^2 V(Y) + 2(-2)(4) Cov(X, Y)$$

(a) If X and Y are independent, then Cov(X,Y) = 0, hence,

$$V(Z) = V[-2X + 4Y - 3] = (-2)^{2}V(X) + (4)^{2}V(Y) = 4(5) + 16(3) = 68.$$

(b) If Cov(X,Y) = 1, then

$$V(Z) = V[-2X + 4Y - 3] = (-2)^{2}V(X) + (4)^{2}V(Y) + 2(-2)(4)Cov(X,Y)$$

$$= 4(5) + 16(3) + 2(-2)(4)(1) = 52.$$
(c) From part (b), $\rho_{X,Y} = \frac{Cov(X,Y)}{\sqrt{V(X)}\sqrt{V(Y)}} = \frac{1}{\sqrt{5}\sqrt{3}} = 0.2582.$

Ouestion 7

 $X \sim discrete uniform$

(a) You can give your answer either in a table

Ī	\overline{x}	1	2	3	1	5	6	7	8	Q	10
	$f_X(x)$										

or

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{10}, & x = 1, 2, \dots, 10; \\ 0, & \text{otherwise}. \end{cases}$$

(b)
$$\Pr(X < 4) = \sum_{x=1}^{3} f(x) = \frac{3}{10} = 0.3$$

(c)
$$\mu = \sum_{x=1}^{10} x \left(\frac{1}{10}\right) = 5.5. \ \sigma^2 = \sum_{x=1}^{10} (x - 5.5)^2 \frac{1}{10} = 8.25$$

Alternatively,
$$E(X^2) = \sum_{x=1}^{10} x^2 \left(\frac{1}{10}\right) = 38.5$$
. $V(X) = 38.5 - 5.5^2 = 8.25$.