

ST2334 PROBABILITY AND STATISTICS
SEMESTER I, AY 2022/2023

Example Questions for the Final Examination
--

- The format of the exam is open book (hard copies or/and soft copies physically stored on the laptop for exam are allowed), onsite through **exam software: Exemplify**. A laptop with the software installed and solid battery that can last for at least 2 hours is needed.
- Calculators of any kind are allowed.
- All chapters covered during the semester, i.e., Chapters 1 – 7.

1. **Multiple choice question: choose the unique correct answer.**

A random variable X has the following probability function.

$$f_X(x) = \frac{x}{4}, \quad \text{for } x = 0.4, 0.6, 0.9, 2.1;$$

and $f_X(x) = 0$ elsewhere.

What type of random variable is X ?

- (a) Continuous
- (b) Discrete
- (c) Unable to determine
- (d) None of the given options

2. **True/False**

Suppose $X \sim N(0, 1)$ and $Y \sim N(0, 1.5^2)$. $P(X < 1)$ is larger than $P(Y < 1)$.

- TRUE
- FALSE

3. **Multiple choice question: choose the unique correct answer.**

A professor receives, on average, 21.7 emails from students the day before the midterm exam. To compute the probability of receiving at least 10 emails on such day, what type of probability distribution will he use?

- (a) Binomial distribution.
- (b) Poisson distribution.
- (c) Normal distribution.
- (d) Negative Binomial distribution.

4. **Multiple choice question: choose the unique correct answer.**

Which of the following would be an appropriate null hypothesis?

- (a) The mean of a population is equal to 55.
- (b) The mean of a sample is equal to 55.
- (c) The mean of a population is greater than 55.
- (d) None of the given options

5. **Multiple choice question: choose the unique correct answer.**

Let A, B be events in sample space S . Which of the following may **NOT** be true?

- (a) $A \cap A' = \emptyset$
- (b) $A \cup A' = S$.
- (c) $(A \cup B)' = A' \cup B'$
- (d) $A \cup B = A \cup (B \cap A')$

6. **Fill in the blank.**

Let X and Y be independent random variables such that $E(X) = 1$, $E(Y) = 2$, $V(X) = 3$, $V(Y) = 4$. Compute $V(2X - Y)$.

Answer: _____

7. **Fill in the blank.**

We toss a fair die until the outcome “6” appears twice. Find the probability that it takes 5 tosses.

Answer: _____

8. **True/False**

Under the usual random sampling setup, we can halve the standard deviation of the sample mean by doubling the sample size.

- TRUE
- FALSE

9. **True/False**

Let X be a discrete random variable; then $E(X)$ always exists.

- TRUE
- FALSE

10. **True/False**

Let $f_X(x)$ be the probability function for random variable X . If $f_X(x) = 0$ for every $x \in (0, 10)$, then $P(X \leq 0 \text{ or } X \geq 10) = 1$.

- TRUE
- FALSE

11. **Multiple choice question: choose the unique correct answer.**

Which of the following is a valid cumulative distribution function?

$$(a) F(x) = \begin{cases} 0 & x \leq -1 \\ 0.3 & -1 < x \leq 1 \\ 0.7 & 1 < x \leq 10 \\ 1 & \text{elsewhere} \end{cases}$$

$$(b) F(x) = \begin{cases} 0 & x < -1 \\ 0.5 & -1 < x < 1 \\ 0.7 & 1 < x < 10 \\ 1 & \text{elsewhere} \end{cases}$$

$$(c) F(x) = \begin{cases} 0 & x < -1 \\ 0.6 & 1 < x \leq 10 \\ 0.7 & -1 < x \leq 1 \\ 1 & \text{elsewhere} \end{cases}$$

$$(d) F(x) = \begin{cases} 0 & x < -1 \\ 0.6 & -1 \leq x < 1 \\ 0.7 & 1 \leq x < 10 \\ 1 & \text{elsewhere} \end{cases}$$

12. **True/False**

Let $\hat{\mu}_1$ and $\hat{\mu}_2$ be two estimators for the population mean μ based on the random sample X_1, X_2, \dots, X_n . If $V(\hat{\mu}_1) < V(\hat{\mu}_2)$, then $\hat{\mu}_1$ must be a better estimator than $\hat{\mu}_2$.

- TRUE
- FALSE

13. **Multiple choice question: choose the unique correct answer.**

The life time (in years) of a certain brand of light bulb follows an exponential distribution with the probability density function: $\frac{1}{2} \exp(-x/2)$. What is the probability that the bulb will last for more than 5 years, given that it has been working for 3 years?

- (a) $\frac{1}{2} \exp(-5/2)$
- (b) $\exp(-5/2)$
- (c) $\frac{1}{2} \exp(-1)$
- (d) $\exp(-1)$

14. **Multiple choice question: choose the unique correct answer.**

Which of the following is a random sample:

- (a) In order to estimate the probability of getting heads for a biased coin, flip the coin 100 times and collect the results for all flips.
- (b) In order to study the scores of undergraduate students of Singapore, 1000 registered undergraduates in NUS were randomly sampled.
- (c) In order to study the average studying hours of students in NUS every week. A random survey with size 1000 was conducted at all libraries of NUS.
- (d) In order to study the average life time of a brand of bulbs, all bulbs' life times in LT32 over a whole semester were recorded.

15. **Multiple choice question: choose the unique correct answer.**

The Central Limit Theorem is important in statistics because

- (a) for a large sample size, n , it says the population is approximately normal.
- (b) for any population, it says the sampling distribution of the sample mean is approximately normal, regardless of the sample size.
- (c) for a large sample size, n , it says the sampling distribution of the sample mean is approximately normal, regardless of the shape of the population.

- (d) for any sized sample, it says the sampling distribution of the sample mean is approximately normal.

16. **Fill in the blank.**

John rolls a fair die 6 times independently. What is the probability that he will get numbers more than 2 at least twice?

Answer: _____

17. **Fill in the blank.**

Jill's bowling scores are approximately normally distributed with mean 170 and standard deviation 20, while Jack's bowling scores are approximately normally distributed with mean 160 and standard deviation 15. If Jack and Jill each bowl one game, then assuming that their scores are independent random variables, the probability that the sum of the scores is higher than 340 is approximately equal to _____.

Answer: _____

18. **Fill in the blank.**

An experiment was carried out to test whether mean weight gain for pigs fed ration A is higher than those fed ration B. Eight pairs of pigs were used. The rations were assigned at random to the two animals within each pair. The gain (in kilograms) after 45 days, assuming normally distributed, are given as follows.

Pairs	1	2	3	4	5	6	7	8	mean	sd
Ration A	30	17	18	21	22	30	24	27	23.625	5.0409
Ration B	26	18	15	20	21	25	27	23	21.875	4.1555
Difference, A - B	4	-1	3	1	1	5	-3	4	1.75	2.7646

Suppose that the pigs within each pair were littermates. What is the observed value of the test statistic in testing the alternative hypothesis that ration A is better, in terms of mean weight gain, than ration B at a 5% significance level?

Answer: _____

19. **Fill in the blanks.**

The mean lifetime of 100 randomly selected pumps made by a particular factory was 200 days. Assuming it is known that the population standard deviation $\sigma = 40$, find a 95% confidence interval for the mean lifetime of pumps made by the factory.

Answer: (_____, _____) (Round to 1 decimal place).

Note: $z_{0.025} = 1.96$; $z_{0.05} = 1.64$.

20. **Fill in the blanks.**

We roll a fair die 3 times. Find the probability that the sum is equal to 5.

Answer: _____

21. **True/False**

Consider the Z -test for $H_0 : \mu = 0$ based on X_1, \dots, X_n i.i.d. $N(\mu, \sigma^2)$. It turns out that $\bar{X} = 2.3$. The p -value for the one-sided test ($H_1 : \mu > 0$) is half of that for the two-sided test ($H_1 : \mu \neq 0$).

- TRUE
- FALSE

22. **Fill in the blanks.**

A new COVID rapid test is able to correctly diagnose that you do not have the virus 90% of the time. However, if you do have the virus, it fails to detect it 25% of the time. Given that the overall COVID infection rate at a particular worker dorm is 20%, what is the probability of a worker being infected if his rapid test does not detect the virus?

Answer: _____

23. **Multiple responses question: choose all that apply.**

Which of the following can happen if the hypothesis is rejected.

- (a) $p\text{-value} > \alpha$;
- (b) test statistic falls in the rejection region;
- (c) type I error occurs;
- (d) type II error occurs.

24. **Fill in the blanks.**

Let X_1, X_2, \dots, X_{100} be independent and identically distributed continuous random variables with $E(X_i) = 5$ and $V(X_i) = 4$. Compute approximately $P\left(\sum_{i=1}^{100} X_i > 510\right)$.

Answer: _____

25. **Multiple choice question: choose the unique correct answer.**

Which of the following statements about probability is **INCORRECT**?

- (a) If A and B are two events and $P(A \cap B) = P(A)$, then $P(A \cap B') = 0$.
- (b) Let S be the sample space and let A be an event. If there exists an $x \in S$ but $x \notin A$, then $P(A) < 1$.
- (c) Let A and B be two events; then $P(A \cup B) \leq P(A) + P(B)$.
- (d) Let A and B be independent events. If $P(A) > 0$ and $P(B) > 0$, then A and B are not mutually exclusive.

26. **Fill in the blanks.**

Consider the following game:

- First round: the gamer flips a fair coin. If he gets a head, he loses; otherwise he wins the round.

- Second round: the gamer flips two fair coins independently. If he gets two heads, he loses; otherwise he wins the round.
- Third round: the gamer flips three fair coins independently. If he gets three heads, he loses; otherwise he wins the round.
- And so on.

What is the probability that the gamer will make his first win in the 4th round?

Note: you can assume that from rounds to rounds, the flips are independently conducted.

Answer: _____

27. Multiple choice question: choose the unique correct answer.

Let $\{X_1, X_2, \dots, X_{25}\}$ be a random sample from the $N(\mu, 2^2)$ distribution. Consider the hypotheses: $H_0 : \mu = 0$ versus $H_1 : \mu \neq 0$, and the test statistic $Z = \frac{\bar{X}}{2/\sqrt{25}}$. Suppose that we reject H_0 if $|z_{obs}| > 2$; and do not reject H_0 otherwise. What is the probability that we will not reject H_0 given that the true value of μ is equal to 2?

Note: $\Phi(z)$ denotes the c.d.f. of the standard normal distribution: $\Phi(z) = Pr(\tilde{Z} \leq z)$ with $\tilde{Z} \sim N(0, 1)$.

- (a) $\Phi(-3) - \Phi(-7)$
- (b) $\Phi(2) - \Phi(-2)$
- (c) $\Phi(-2) - \Phi(-6)$
- (d) $\Phi(-3) - \Phi(-8)$

28. True/False

For any $\theta \in \mathbb{R}$, the function

$$f(x) = \begin{cases} \theta - x & \text{if } \theta - 1 \leq x < \theta \\ x - \theta & \text{if } \theta \leq x \leq \theta + 1 \\ 0 & \text{elsewhere} \end{cases}$$

can serve as a probability density function of some distribution, whose population mean is equal to θ .

- TRUE
- FALSE

29. Multiple choice question: choose the unique correct answer.

In a course, students are graded based on a “normal curve”. For example, students within 0.5 standard deviation from the mean receive a *C*; between 0.5 and 1.0 standard deviation above the mean receive a *C+*; between 1.0 and 1.5 standard deviation above the mean receive a *B*; between 1.5 and 2.0 standard deviation receive a *B+*, etc. The class average in an exam was 60 with a standard deviation of 10. What are the bounds for a *B* grade and the percentage of students who will receive a *B* grade?

- (a) (65, 75), 24.17%

- (b) (65, 75), 12.08%
- (c) (70, 75), 18.38%
- (d) (70, 75), 9.19%

30. **Multiple choice question: choose the unique correct answer.**

Flip an unfair coin. If a head shows, roll a fair die and report the number; otherwise, roll a fair die twice and report the summation minus 1. Then $P(6 \text{ is reported}) = ?$

- (a) 1/6
- (b) 1/4
- (c) 1/2
- (d) can not tell

31. **Fill in the blanks.**

Suppose that X_1 is Poisson with expectation 1, X_2 is Poisson with expectation 1, and X_3 is Poisson with expectation 2 and assume that the three random variables are independent. Let $Y_1 = X_1 + X_2$ and let $Y_2 = X_2 + X_3$. The conditional probability that $Y_1 = 1$, given that $Y_2 = 2$, is equal to _____.

Answer: _____ (Leave your answer to 3 decimal places.)

32. **Fill in the blanks.**

An urn contains 3 red balls and 5 white balls. 4 balls are drawn uniformly at random without replacement from the urn. Let X be the random variable for the number of red balls drawn. If $X \leq 1$, you win $\$X$. If $X > 1$, you flip a fair coin. If the coin comes up heads, you double your winnings and win a total of $\$2X$. If the coin comes up tails, you still win $\$X$. Let W be the random variable for your winnings. Find the probability that W is odd.

Answer: _____ (Leave your answer to 4 decimal places.)

33. **Multiple choice question: choose the unique correct answer.**

Let X be a random variable with density function

$$f(x) = \begin{cases} k\sqrt{x} & , \text{ for } 0 \leq x \leq 1 \\ ke^{\frac{1-x}{2}} & , \text{ for } x > 1 \\ 0 & , \text{ otherwise} \end{cases}$$

where k is a constant. What is the value of k ?

- (a) 1/2
- (b) 2/5
- (c) 3/8
- (d) 4/9

34. **Multiple responses question: choose all that apply.**

Let X_1, X_2, \dots, X_{n_1} be independent and identically distributed (i.i.d.) random variables with population mean μ_1 ; let Y_1, Y_2, \dots, Y_{n_2} be i.i.d. random variables with population mean μ_2 ; let U_1, U_2, \dots, U_{n_3} be i.i.d. random variables with population mean 4. All these random variables have the unknown but common variance σ^2 . Which of the following is/are unbiased estimator(s) for σ^2 ?

- (a) $\frac{\sum_{i=1}^{n_1} (X_i - \bar{X})^2 + \sum_{j=1}^{n_2} (Y_j - \bar{Y})^2 + \sum_{k=1}^{n_3} (U_k - 4)^2}{n_1 + n_2 + n_3 - 2}.$
- (b) $\frac{\sum_{i=1}^{n_1} (X_i - \bar{X})^2 + \sum_{j=1}^{n_2} (Y_j - \bar{Y})^2 + \sum_{k=1}^{n_3} (U_k - 4)^2}{n_1 + n_2 + n_3 - 3}.$
- (c) $\frac{\sum_{i=1}^{n_1} (X_i - \bar{X})^2 + \sum_{j=1}^{n_2} (Y_j - \bar{Y})^2 + \sum_{k=1}^{n_3} (U_k - \bar{U})^2}{n_1 + n_2 + n_3 - 2}.$
- (d) $\frac{\sum_{i=1}^{n_1} (X_i - \bar{X})^2 + \sum_{j=1}^{n_2} (Y_j - \bar{Y})^2 + \sum_{k=1}^{n_3} (U_k - \bar{U})^2}{n_1 + n_2 + n_3 - 3}.$

35. **Fill in the blank.**

Let X_1, X_2, \dots, X_{10} be independent and identically distributed random variables having the exponential distribution $\text{Exp}(1)$. Let $T = \min\{X_1, X_2, \dots, X_{10}\}$. Find $E(T)$.

Answer: _____