AN AXIOMATIC DERIVATION OF THE FEYNMAN PATH SUM AT THE LIGHT-CONE KERNEL WIDTH AND CLOSURE ORDER $\alpha = 2$ OF THE CAUSAL ACCUMULATION LAW

SAM LEIZERMAN

ABSTRACT. We take the Causal Accumulation Law (CAL) at closure order $\alpha = 2$ and give its memory the light-cone width: at lag τ the spatial profile of the kernel is the light front of radius $c\tau$. Four results follow.

Lorentz invariance selects the order. The symbol of the order-α light-cone memory is $(\tau_c R)^{-\alpha} \sin((\alpha-1)\varphi)/((\alpha-1)\sin\varphi)$, with $s = R\cos\varphi$ and $c|k| = R\sin\varphi$. It depends on $s^2 + c^2k^2$ alone exactly when $\alpha = 2$ (and trivially at $\alpha = 0$); the leading defect is $\alpha(2-\alpha)\varphi^2/6$.

- At $\alpha = 2$ the kernel is the retarded Green's function of the wave operator: Kirchhoff's shell in $3+1$ dimensions.
- In $1+1$ dimensions it is the product of one order-one accumulation along each null leg. The order two is one plus one.

The balanced $\alpha = 2$ law is Klein–Gordon. In the free, flat sector the transport chain is the identity, so the law is $T = 2J + K_2 J$.

- A carrier whose present balances its light-cone past, $T = 0$, obeys $(\Box + \mu^2)J = 0$ with $\mu\tau_c = 1/\sqrt{2}$.
- The kernel's lag scale is the only scale, so the mass is the carrier's own rate, and only ratios between carriers are physical. No coupling constant and no double scaling are needed.
- The phase belongs to the source. The double accumulation supplies the half turn by itself, and no factor is put on the memory; an added factor -1 would make the mass tachyonic.
- The law is conservative exactly at $\alpha = 2$. Just below, both signs of frequency decay at $(\pi/4)(2-\alpha)\mu^2/\omega$; just above, the causal response runs away at the same rate.

The field equation is a path sum. Its Neumann series converges absolutely for every bounded source with zero past. It is a sum over piecewise-null histories: $n+1$ light-cone legs, with the real amplitude $-\mu^2$ at each of the n corners. It resums to the retarded massive propagator.

- In $1+1$ dimensions the n-th term equals the $(2n+1)$-corner amplitude of Feynman's checkerboard divided by $2ic\mu$, because $(-\mu^2)^n = (i\mu)^{2n}$.
- In $3+1$ dimensions, by descent, the histories without corners live on the cone, and the corners alone fill its interior. The Bessel tail of the massive propagator is the sum over histories with at least one corner.

Unitarity, the readout, and Feynman's kernel.

- Each sign of frequency evolves by a unitary group generated by $\sqrt{m^2c^4 - \hbar^2c^2\nabla^2}$.
- On the positive-frequency sector the trace of the carrier closure, $\int JJ^\dagger$, is conserved, so the trace readout keeps its normalization in every frame. On mixed frequencies it does not.
- Nothing near $\alpha = 2$ selects a sign of frequency. As at the edge, the sign is inherited from the source's past.
- Removing the rest phase and letting $c \to \infty$, the positive-frequency part of the light-cone path sum converges to Feynman's free kernel with error $O(1/c^2)$, and its composition law is Feynman's time slicing.

The $\alpha = 2$ dynamics is real. The i of Feynman's kernel enters only through the sign of frequency inherited from the source.

Numerical runs confirm each statement.

ORCID: 0009-0000-0133-2291. Companion to *The Causal Accumulation Law: The Four Corners of α, Algebraic Closure, and the Standard Model* [1], to *Schrödinger evolution from the bare Causal Accumulation Law at $\alpha \to 1^-$* [2], and to *An Axiomatic Derivation of the Born Rule at the δ-Kernel Limit and Closure Order $\alpha = 2$* [3]. This version follows the revised edge paper: no coupling constant, no time unit beyond the kernel's own lag scale, and no phase factor on the memory. Dated November 13, 2026.

Contents

1. Scope and claim 2
2. Definitions 3
3. Axioms and the field equation 4
4. The light-cone memory: symbol and Lorentz invariance 4
5. The balanced $\alpha = 2$ law is Klein–Gordon 6
6. The path sum 7
7. Composition, unitarity, and the readout 8
8. The nonrelativistic limit: Feynman's free kernel 10
9. Numerical verification 11
10. Remarks on scope 12
11. Conclusion 13

Appendix A. Notation 14

Appendix B. Numerical methods 14

References 14

1. Scope and claim

The companion paper [2] obtains the free Schrödinger equation at the $\alpha \to 1^-$ edge of the CAL. There the causal structure is time ordering alone, the memory is a one-way accumulator, and nothing interacts. The Lorentz signature, and with it the light cone, belongs to the quadratic closure $JJ^\dagger$ at $\alpha = 2$ [1, 2]. This paper takes the CAL at $\alpha = 2$, with the memory spread over the light cone, and asks what a carrier whose present balances that past obeys. It follows the revised edge paper: no coupling constant, no time unit beyond the kernel's own lag scale, and no phase factor on the memory.

> *Give the order-α memory the light-cone width. Its symbol is Lorentz invariant exactly at $\alpha = 2$ (Theorem 4.2), where the kernel is the retarded Green's function of $\Box$ (Corollary 4.3). A carrier that balances its light-cone past obeys the Klein–Gordon equation, with the kernel's lag scale as the only scale (Theorem 5.1). The law is conservative exactly at $\alpha = 2$: below it every mode decays, and above it the causal response runs away (Theorem 5.2). Its Neumann series is a sum over piecewise-null histories with amplitude $-\mu^2$ per corner. It converges absolutely and resums to the retarded massive propagator (Theorem 6.1). In $1+1$ dimensions it is Feynman's checkerboard term by term (Proposition 6.2). In $3+1$ dimensions the interior of the cone is filled by the corners alone (Proposition 6.4). The past enters only through the pair $(J, \partial_t J)$ (Proposition 7.1). Each sign of frequency is unitary, and on positive frequency the carrier trace is conserved (Theorem 7.3). As $c \to \infty$ the positive-frequency path sum converges to Feynman's free kernel (Theorem 8.1, Corollary 8.2).*

Remark 1.1 (Bookkeeping). *Inputs:*

- the CAL (Axiom 1);
- closure order $\alpha = 2$ with the light-cone profile, in the free, flat sector where the transport chain is the identity (Axiom 2);
- the balance reading with a source of zero past (Axiom 3);
- the phase as a property of the source (Remark 3.1);
- for the readout only, one sign of frequency, inherited from the source's past (Theorem 7.3).

Derived: every proposition, theorem and corollary of §§4–8.

Not claimed:

- interaction between two carriers at $\alpha = 2$;
- spin: each entry of J obeys the scalar Klein–Gordon equation here, and a first-order (Dirac) form is not derived;
- a mass ratio between different carriers (every carrier with the same kernel has the same rate here), or the relation to the edge's rest rotation [2];
- a frame-independent position readout.

The $1 + 1$ checkerboard and its nonrelativistic limit are known [5, 6, 7], and so are chain sums for the retarded propagator on causal sets [8]. What is new here is the source: the history sum is the memory series of the CAL at $\alpha = 2$, and the light-cone width is selected by Lorentz invariance of the memory symbol.

2. Definitions

Throughout, $d \in \{1, 2, 3\}$ is the spatial dimension, s is a complex frequency with $\operatorname{Re} s > 0$, $k \in \mathbb{R}^d$ is a wave vector, τ_c is the kernel's lag scale, and $E_c = \hbar/\tau_c$. It is the only scale in the problem. Time is relational: only ratios such as $\mu\tau_c$ are physical. Complex powers take the principal branch.

Definition 2.1 (Light-cone profile)**.** For $\tau > 0$, the light-cone profile ν_τ is the probability measure on $\mathbb{R}^d$ that is rotation invariant, supported in $|y| \le c\tau$, and has Fourier transform

$$\hat{\nu}_\tau(k) = \int e^{-ik\cdot y}\, \nu_\tau(dy) = \operatorname{sinc}(c|k|\tau), \qquad \operatorname{sinc} z = \frac{\sin z}{z}.$$

Explicitly:

- $d = 3$: the uniform measure on the sphere $|y| = c\tau$, i.e. $\delta(|y| - c\tau)/(4\pi c^2\tau^2)$;
- $d = 2$: the density $\theta(c\tau - |y|)/\big(2\pi c\tau\sqrt{c^2\tau^2 - |y|^2}\big)$;
- $d = 1$: the uniform density $\theta(c\tau - |y|)/(2c\tau)$.

The profiles for $d = 2$ and $d = 1$ are the orthogonal projections of the sphere in $d = 3$. For $d = 1$ this is Archimedes' hat-box theorem. Projection restricts the Fourier transform to a subspace, so all three have the same transform.

The *light-cone width* is the statement that the profile at lag τ has radius $c\tau$. By contrast, the profile at the edge [2, Def. 2.2] does not depend on the lag.

Definition 2.2 (Order-α light-cone memory)**.** For $\alpha > 0$ and f a function on spacetime,

$$(K_\alpha f)(t, x) = \tau_c^{-\alpha} \int_0^\infty \frac{\tau^{\alpha-1}}{\Gamma(\alpha)} \int_{\mathbb{R}^d} f(t - \tau, x - y)\, \nu_\tau(dy)\, d\tau.$$

It integrates over the closed past light cone $J^-(t, x) = \{(t', x') : |x - x'| \le c(t - t')\}$, and in $d = 3$ over the cone itself.

A function f has *zero past* if $f = 0$ for $t < t_0$, for some t_0. For bounded f with zero past, $K_\alpha f$ is finite and

$$\sup_x |K_\alpha f(t, x)| \le \tau_c^{-\alpha} \frac{(t - t_0)^\alpha}{\Gamma(\alpha + 1)} \sup|f|,$$

because ν_τ is a probability measure.

Definition 2.3 (Symbol)**.** On $e^{st+ik\cdot x}$ with $\operatorname{Re} s > 0$, K_α acts by multiplication by

$$m_\alpha(s, k) = \tau_c^{-\alpha} \int_0^\infty \frac{\tau^{\alpha-1}}{\Gamma(\alpha)}\, e^{-s\tau}\, \operatorname{sinc}(c|k|\tau)\, d\tau.$$

On a carrier $e^{-iEt/\hbar+ik\cdot x}$ with real E, the value is the limit $\operatorname{Re} s \downarrow 0$ at $s=-iE/\hbar$, as for the $\alpha=1$ value in [2, Def. 2.1].

For real $s>0$ and $k\neq 0$ write

$$s=R\cos\varphi, \qquad c|k|=R\sin\varphi, \qquad R=\sqrt{s^2+c^2k^2}, \quad \varphi\in(0,\tfrac{\pi}{2}).$$

The wave operator is $\Box=\partial_t^2-c^2\nabla^2$, with symbol $s^2+c^2k^2$.

3. Axioms and the field equation

Axiom 1 (Causal accumulation law; C8 of [1])**.**

$$T(x)=\alpha J(x)-(1-\alpha)\int_{J^-(x)} k(\tau;x,x')\,\mathcal{P}(x,x')\,J(x')\,d^4x'.$$

The kernel k and the transport $\mathcal{P}$ are real, and J takes values in $\mathbb{C}\otimes\mathbb{H}\cong M_2(\mathbb{C})$.

Axiom 2 (Closure order two at the light-cone width)**.** $\alpha=2$. The kernel is $k(\tau,y)=\tau_c^{-2}\,\tau\,\nu_\tau(y)$ (Definitions 2.1 and 2.2). In the free, flat sector the bitensor coupling, the parallel transport and the internal fiber transport are the identity, $\Pi P\Phi=\mathbf{1}$, so the memory term is K_2J.

At $\alpha=2$ the prefactor $-(1-\alpha)$ is $+1$, so the law reads $T=2J+K_2J$.

Axiom 3 (Balance)**.** For a bounded source S with zero past,

$$T=2J+K_2J=S. \tag{1}$$

T is the new state, and it vanishes only when the present balances the accumulated past. A free carrier is such a balanced configuration: $T=0$ wherever $S=0$. A source is a nonzero imbalance. A field is started by a source acting on a zero past, and no initial-value data are used.

The edge's update rule [2, Axiom 3] is not used here: at $\alpha=2$ the term $2J$ alone would double the current at every update.

Remark 3.1 (The phase belongs to the source). The channel angle is what an order-α accumulation does to the source's phase channel, $\theta=\alpha\pi/2$ [2, Axiom 2]. At $\alpha=2$ the double accumulation supplies the half turn by itself: on timelike modes the symbol $1/(s^2+c^2k^2)$ is negative. No factor is put on the memory.

An added factor -1 on the memory would flip the sign of μ^2. That is what the edge's former prefactor $e^{-i\sigma\alpha\pi/2}$ gives at $\alpha=2$. The dispersion would become $E^2=\hbar^2c^2k^2-m^2c^4$, a tachyon, and modes with $c|k|<\mu$ would grow at rate $\sqrt{\mu^2-c^2k^2}$. Table 2 shows both.

4. The light-cone memory: symbol and Lorentz invariance

Proposition 4.1 (Symbol)**.** *For $\alpha>0$, $\operatorname{Re} s>0$ and $k\neq 0$, with $b=c|k|$,*

$$\tau_c^{\alpha}\,m_\alpha(s,k)=\frac{(s-ib)^{1-\alpha}-(s+ib)^{1-\alpha}}{2ib\,(\alpha-1)} \quad (\alpha\neq 1), \qquad \tau_c\,m_1(s,k)=\frac{\arctan(b/s)}{b}.$$

For real $s>0$, $\tau_c^{\alpha}m_\alpha=R^{-\alpha}h_\alpha(\varphi)$ with

$$h_\alpha(\varphi)=\frac{\sin((\alpha-1)\varphi)}{(\alpha-1)\sin\varphi} \quad (\alpha\neq 1), \qquad h_1(\varphi)=\frac{\varphi}{\sin\varphi}.$$

At $k=0$ the symbol is $(\tau_c s)^{-\alpha}$, the Weyl symbol of [2, Def. 2.1]. *The symbol does not depend on d.*

Proof. Write $\operatorname{sinc}(b\tau) = \sin(b\tau)/(b\tau)$ and use $\int_0^\infty \tau^{\beta-1} e^{-z\tau} d\tau = \Gamma(\beta) z^{-\beta}$ for $\operatorname{Re} z > 0$, with $\beta = \alpha - 1$ and $z = s \mp ib$. The integral converges at $\tau = 0$ for $\alpha > 0$, because $\tau^{\alpha-2} \sin(b\tau) = O(\tau^{\alpha-1})$. Then $\Gamma(\alpha-1)/\Gamma(\alpha) = 1/(\alpha-1)$ gives the first formula, and its limit $\alpha \to 1$ gives the second. For real s, $s \mp ib = Re^{\mp i\varphi}$. The transform $\hat{\nu}_\tau$ is the same for every d (Definition 2.1). □

Theorem 4.2 (Lorentz invariance selects $\alpha = 2$)**.** *For $\alpha > 0$ the following are equivalent:*

(1) *$m_\alpha(s,k)$ depends on (s,k) only through $s^2 + c^2k^2$;*
(2) *$\alpha = 2$.*

Then $m_2(s,k) = \tau_c^{-2}/(s^2 + c^2k^2)$. In the limit $\alpha \to 0^+$ the memory becomes the identity, which is trivially invariant. For small φ,

$$h_\alpha(\varphi) = 1 + \frac{\alpha(2-\alpha)}{6}\,\varphi^2 + O(\varphi^4),$$

so the Lorentz defect is positive for $0 < \alpha < 2$ and negative for $\alpha > 2$. The function h_α is symmetric under $\alpha \mapsto 2 - \alpha$, about the edge $\alpha = 1$.

Proof. For real $s > 0$ and $k \neq 0$, (R, φ) is a bijection onto $(0,\infty) \times (0, \pi/2)$. Statement (1) holds on this set exactly when h_α is constant. Both sides of (1) are analytic in s for $\operatorname{Re} s > 0$, so the real half-line determines the whole half-plane.

Expanding, $\sin((\alpha-1)\varphi) = (\alpha-1)\varphi\,[1 - (\alpha-1)^2\varphi^2/6 + O(\varphi^4)]$ and $\sin\varphi = \varphi\,[1 - \varphi^2/6 + O(\varphi^4)]$. Their quotient gives the expansion, which also holds at $\alpha = 1$. A constant h_α needs $\alpha(2-\alpha) = 0$, so $\alpha = 2$ for $\alpha > 0$. Conversely $h_2 \equiv 1$. Symmetry: h_α depends on $\alpha - 1$ only through $\sin(\beta\varphi)/\beta$, which is even in β. □

On a carrier, $s = -iE/\hbar + 0$, and $s^2 + c^2k^2 = -(E^2 - \hbar^2c^2k^2)/\hbar^2$ with the retarded prescription. This is the Minkowski square of the four-momentum: the $\alpha = 2$ memory is a function of p^2 and of the sign of E, both invariant under orthochronous Lorentz transformations.

Corollary 4.3 (The order-2 kernel is the retarded Green's function of $\Box$)**.** *$\tau_c^2 K_2 f = G * f$ with $G(t,\cdot) = \theta(t)\, t\, \nu_t$, and $\Box(G * f) = f$ for f with zero past. Explicitly,*

$$G^{(3)} = \frac{\theta(t)\,\delta(|x| - ct)}{4\pi c^2 t}, \qquad G^{(2)} = \frac{\theta(ct - |x|)}{2\pi c\sqrt{c^2t^2 - |x|^2}}, \qquad G^{(1)} = \frac{\theta(ct - |x|)}{2c}.$$

$G^{(3)}$ is Kirchhoff's shell [9].

Proof. By Theorem 4.2 the symbol of G is $1/(s^2 + c^2k^2)$, the inverse of the symbol of $\Box$. Both G and f vanish in the past, so $G * f$ is the retarded solution. The explicit forms are $t\,\nu_t$ from Definition 2.1. □

Proposition 4.4 ($\alpha = 2$ is 1+1: two null accumulations)**.** *In $d = 1$ let $u = ct - x$ and $v = ct + x$. Then $\Box = 4c^2 \partial_u \partial_v$ and*

$$(G * f)(u,v) = \frac{1}{4c^2} \int_{-\infty}^{u} \int_{-\infty}^{v} f(u', v')\, dv'\, du'.$$

So $G = (4c^2)^{-1} W_u^1 W_v^1$, the product of one order-one causal accumulation along each null direction. The order-two memory is two order-one accumulators of the kind found at the edge [2, Prop. 6.1]*, one per null leg.*

Proof. $\partial_t = c(\partial_u + \partial_v)$ and $\partial_x = \partial_v - \partial_u$, so $\partial_t^2 - c^2\partial_x^2 = 4c^2\partial_u\partial_v$. The Jacobian is $du\,dv = 2c\,dt\,dx$, and $G^{(1)} = \theta(u)\theta(v)/(2c)$. □

Proposition 4.5 (Descent)**.** *For an even function f_1 on $\mathbb{R}$, define the radial function $(\mathcal{D}f_1)(y) = -(2\pi|y|)^{-1} f_1'(|y|)$ on $\mathbb{R}^3$. Then $\widehat{\mathcal{D}f_1}(k) = \hat{f}_1(|k|)$. In particular $G^{(3)*n}(t,\cdot) = \mathcal{D}\, G^{(1)*n}(t,\cdot)$ for every $n \geq 1$, and the same holds for any convergent series in powers of G.*

Proof. For a radial function, $\hat{f}_3(\kappa) = (4\pi/\kappa)\int_0^\infty r f_3(r)\sin(\kappa r)\,dr$. With $f_3 = \mathcal{D}f_1$, an integration by parts gives $2\int_0^\infty f_1(r)\cos(\kappa r)\,dr = \hat{f}_1(\kappa)$; for distributions it holds weakly. The spatial transform of $G^{*n}(t,\cdot)$ is the n-fold time convolution of $\theta(t)\sin(c|k|t)/(c|k|)$. That is the same function of $|k|$ for $d = 1$ and $d = 3$. This is the classical method of descent [9], read on symbols. □

5. The balanced $\alpha = 2$ law is Klein–Gordon

Theorem 5.1 (Klein–Gordon)**.** *Set*

$$\mu^2 = \frac{1}{2\tau_c^2}, \qquad mc^2 = \hbar\mu = \frac{\hbar}{\sqrt{2}\,\tau_c}.$$

(1) *For every bounded source S with zero past, (1) has exactly one locally bounded solution with zero past. It is*

$$J = \tfrac{1}{2}S - \tfrac{1}{2}\mu^2\, G_\mu * S,$$

where G_μ is the retarded fundamental solution of $\Box + \mu^2$.

(2) *$(\Box+\mu^2)J = \frac{1}{2}\Box S$ in the sense of distributions. On every open set where $S = 0$, J solves the Klein–Gordon equation $(\Box + \mu^2)J = 0$.*

(3) *A plane wave $e^{-iEt/\hbar + ik\cdot x}$ solves $T = 0$, with the memory valued as in Definition 2.3, if and only if $E^2 = \hbar^2c^2k^2 + m^2c^4$.*

The mass is the kernel's own rate, $\mu\tau_c = 1/\sqrt{2}$. With no external time unit this ratio is the physical content, and every carrier with the same kernel has the same rate. No coupling constant and no double scaling are needed.

Proof. By Corollary 4.3, $K_2 = \tau_c^{-2}G* = 2\mu^2 G*$, so (1) is $J = \frac{1}{2}S - \mu^2 G * J$. Existence and uniqueness are Theorem 6.1(1)–(2), and the resummation is Theorem 6.1(3).

(2) $\Box G_\mu * S = S - \mu^2 G_\mu * S$, so $(\Box + \mu^2)J = \frac{1}{2}\Box S + \frac{1}{2}\mu^2 S - \frac{1}{2}\mu^2 S$.

(3) The symbol of T is $2 + \tau_c^{-2}/(s^2 + c^2k^2)$. It vanishes if and only if $s^2 + c^2k^2 = -\mu^2$. At $s = -iE/\hbar$ this is $E^2/\hbar^2 = c^2k^2 + \mu^2$. On shell $E^2 \neq \hbar^2c^2k^2$, so the memory value is finite. □

Theorem 5.2 (Conservative exactly at $\alpha = 2$)**.** *For α near 2, take the order-α light-cone law $T = \alpha J - (1-\alpha)K_\alpha J$, with the same kernel family and no factor on the memory. Its balanced modes satisfy $\alpha + (\alpha - 1)\, m_\alpha(-iE/\hbar, k) = 0$. Write $\omega = E/\hbar$, and $\mu^2 = \omega^2 - c^2k^2$ at the root.*

(1) *At $\alpha = 2$ the roots $\omega = \pm\sqrt{c^2k^2 + \mu^2}$ are real: both signs of frequency are conserved.*

(2) *For $\alpha = 2 + \delta$, both roots move off the real axis by*

$$\operatorname{Im}\omega = \frac{\pi}{4}\,\delta\,\frac{\mu^2}{|\omega|}\,\big(1 + O(\delta)\big).$$

So below 2 both signs decay at the same rate, and nothing selects a sign. Above 2 both grow: the causal response runs away, as for the Abraham–Lorentz–Dirac equation [10].

Proof (first order in δ). By Proposition 4.1, on real timelike ω the symbol is $m_\alpha = |m_\alpha|\, e^{i\,\mathrm{sgn}(\omega)\,\alpha\pi/2}$. At $\alpha = 2 + \delta$ the phase is $-e^{i\,\mathrm{sgn}(\omega)\delta\pi/2}$. With $|m_2| = \tau_c^{-2}/\mu^2$ at the root, the symbol of T changes by $-i\pi\delta\,\mathrm{sgn}(\omega)$, and its ω-derivative is $4\omega/\mu^2$. Hence $\Delta\omega = i\pi\delta\mu^2/(4|\omega|)$.

Roots in the upper half-plane are reached by the convergent Laplace integral, so they are genuine growing solutions of the sourced problem with zero past. Table 3 checks the rate, the count of growing modes, and a time-stepped run. □

6. The path sum

Theorem 6.1 (The Neumann series is a sum over light-cone histories)**.** *Let S be bounded with $S = 0$ for $t < t_0$.*

(1) *The series*

$$J = \tfrac{1}{2}\sum_{n\ge 0}(-\mu^2)^n\, G^{*n} * S \qquad (G^{*0} * S = S)$$

converges absolutely and uniformly on every slab $t \le t_0 + t_1$, with

$$\sup_x \big|G^{*n} * S(t,x)\big| \le \frac{(t-t_0)^{2n}}{(2n)!}\,\sup|S|, \qquad \sup_x|J(t,x)| \le \tfrac{1}{2}\cosh\big(\mu(t-t_0)\big)\sup|S|.$$

(2) *Its sum is the only locally bounded solution of (1) with zero past.*

(3) *$G_\mu \equiv \sum_{n\ge 0}(-\mu^2)^n G^{*(n+1)}$ is the retarded fundamental solution of $\Box + \mu^2$, with symbol $1/(s^2 + c^2k^2 + \mu^2)$, and $J = \frac{1}{2}S - \frac{1}{2}\mu^2 G_\mu * S$.*

(4) *In $d = 3$,*

$$G^{*(n+1)}(t,x) = \int \delta\Big(t - \sum_{i=0}^{n}\tau_i\Big)\,\delta^3\Big(x - c\sum_{i=0}^{n}\tau_i\hat\omega_i\Big)\prod_{i=0}^{n}\tau_i\,d\tau_i\,\frac{d\Omega_i}{4\pi}.$$

Each term is an integral over histories made of $n+1$ null legs, leg i of duration τ_i in direction $\hat\omega_i$, with weight τ_i per leg. So G_μ is the sum over piecewise-null histories with amplitude $-\mu^2$ at each corner.

Proof. (1) Let $F(t) = \sup_x|f(t,\cdot)|$. Because ν_τ is a probability measure, $\sup_x|G * f(t,\cdot)| \le \int_0^{t-t_0}\tau F(t-\tau)\,d\tau$, which is the Riemann–Liouville integral of order two of F from t_0. By induction, the n-fold iterate is bounded by the order-$2n$ integral of the constant $\sup|S|$, which is $(t-t_0)^{2n}/(2n)!\,\sup|S|$. Summing $\mu^{2n}(t-t_0)^{2n}/(2n)!$ gives the cosh.

(2) Each partial sum J_N satisfies $J_N = \frac{1}{2}S - \mu^2 G * J_{N-1}$; let $N \to \infty$. If two locally bounded solutions have zero past, their difference Δ satisfies $\Delta = -\mu^2 G * \Delta$. Iterating n times gives $\sup_x|\Delta(t)| \le \mu^{2n}(t-t_0)^{2n}/(2n)!\,\sup_{[t_0,t]}|\Delta| \to 0$.

(3) $\Box G^{*(n+1)} = G^{*n}$ with $G^{*0} = \delta$, so $(\Box+\mu^2)G_\mu$ telescopes to δ. The support is in the forward cone, and the symbol is the geometric series $\sum(-\mu^2)^n(s^2+c^2k^2)^{-n-1}$. Absolute convergence of the kernel series is shown in Propositions 6.2 and 6.4. Then $\frac{1}{2}\sum_{n\ge1}(-\mu^2)^n G^{*n} * S = -\frac{1}{2}\mu^2 G_\mu * S$.

(4) $G(\tau,\cdot) = \tau\,\nu_\tau$, and ν_τ is the uniform measure $d\Omega/4\pi$ on the sphere of radius $c\tau$. Convolving $n+1$ times gives the formula. □

Proposition 6.2 ($1+1$: explicit terms and Feynman's checkerboard)**.** *Let $d = 1$, $u = ct - x$, $v = ct + x$.*

(1)

$$G^{*(n+1)} = \frac{\theta(u)\theta(v)}{2c}\,\frac{1}{(n!)^2}\Big(\frac{uv}{4c^2}\Big)^n, \qquad G_\mu = \frac{\theta(u)\theta(v)}{2c}\,J_0\Big(\mu\sqrt{t^2 - x^2/c^2}\Big),$$

and $\sum_n \mu^{2n}|G^{*(n+1)}| = \frac{\theta(u)\theta(v)}{2c} I_0\big(\mu\sqrt{t^2 - x^2/c^2}\big)$.

(2) *Let $\Phi_{2n+1}(t,x)$ be the continuum checkerboard amplitude of null zigzags from $(0,0)$ to (t,x) that start along one null direction and end along the other, with $2n+1$ corners and amplitude $i\mu$ per corner per unit time. Then*

$$(-\mu^2)^n\, G^{*(n+1)} = \frac{\Phi_{2n+1}}{2ic\mu} \qquad \text{for every } n \ge 0,$$

and $G_\mu = (2ic\mu)^{-1}\sum_n \Phi_{2n+1}$.

(3) *On a lattice of time step ε, with Feynman's amplitude $i\varepsilon\mu$ per corner* [5]*, the corner-odd amplitude $L_\varepsilon(t,x)$ satisfies $L_\varepsilon/\varepsilon \to \sum_n \Phi_{2n+1} = i\mu J_0\big(\mu\sqrt{t^2-x^2/c^2}\big)$, uniformly on compact subsets of the open cone.*

Proof. (1) By Proposition 4.4, $G*$ is $(4c^2)^{-1}$ times one integration in u and one in v. Starting from $\theta(u)\theta(v)/(2c)$, n steps give $(4c^2)^{-n}(u^n/n!)(v^n/n!)/(2c)$. The series is that of $J_0(z) = \sum_n (-1)^n (z/2)^{2n}/(n!)^2$ with $(z/2)^2 = \mu^2 uv/(4c^2)$, and $uv = c^2t^2 - x^2$. With all signs positive it is I_0.

(2) Along a right-moving leg u is constant and v grows at rate $2c$; along a left-moving leg the roles swap. A zigzag with $2n+1$ corners has $n+1$ legs of each kind. Their durations add to $v/2c$ and $u/2c$, so the set of such zigzags has volume $(v/2c)^n/n! \cdot (u/2c)^n/n!$. Hence $\Phi_{2n+1} = (i\mu)^{2n+1}(uv/4c^2)^n/(n!)^2$, and $(i\mu)^{2n+1}/(2ic\mu) = (-\mu^2)^n/(2c)$.

(3) With $c=1$, $N = t/\varepsilon$ steps, $N_R = (t+x)/2\varepsilon$ right steps and $N_L = (t-x)/2\varepsilon$ left steps,

$$L_\varepsilon = \sum_n (i\varepsilon\mu)^{2n+1} \binom{N_R-1}{n}\binom{N_L-1}{n}.$$

Here $\varepsilon^n\binom{N_R-1}{n} \to (v/2)^n/n!$ and $\varepsilon^n\binom{N_R-1}{n} \le (v/2)^n/n!$, so dominated convergence gives the limit. The error is $O(\varepsilon)$ (Table 5). □

Remark 6.3 (Reading the checkerboard). The CAL insertion is real: $-\mu^2$ per corner of a light-cone history. Feynman's corner amplitude is imaginary: $i\mu$ per corner of a null zigzag. They agree term by term because $(-\mu^2)^n = (i\mu)^{2n}$. A history with n insertions consists of $n+1$ steps of G. By Proposition 4.4 each step is one u-leg and one v-leg, so the history has $2n+2$ legs and $2n+1$ corners: exactly the zigzags counted by Φ_{2n+1}. The fixed factor $1/(2ic\mu)$ converts the corner-odd entry of the first-order checkerboard [5, 6] into the second-order propagator G_μ. In the chiral basis that entry is the mass term $\mu\, G_\mu$ of the $1+1$ Dirac propagator.

Proposition 6.4 ($3+1$: the corners fill the cone)**.** *Let $d=3$, $r=|x|$ and $w = c^2t^2 - r^2$. Then*

$$G^{*1} = \frac{\delta(r-ct)}{4\pi cr}, \qquad G^{*(n+1)} = \theta(ct-r)\,\frac{n\,w^{n-1}}{2\pi c\,(4c^2)^n (n!)^2} \quad (n \ge 1),$$

and

$$G_\mu = \frac{\delta(r-ct)}{4\pi cr} - \theta(ct-r)\,\frac{\mu}{4\pi c^2}\,\frac{J_1(\mu\sqrt{w}/c)}{\sqrt{w}}.$$

The histories without corners lie on the cone (Huygens). Every history with $n \ge 1$ corners lands strictly inside. So the Bessel tail of the massive propagator is the sum over histories with at least one corner. The tail series converges absolutely, with $\sum_{n\ge1} \mu^{2n}|G^{*(n+1)}| = \frac{\mu}{4\pi c^2}\, I_1(\mu\sqrt{w}/c)/\sqrt{w}$.

Proof. Apply $\mathcal{D}$ (Proposition 4.5) to Proposition 6.2(1) with $x \to r$. For $n=0$, $-\partial_r\theta(ct-r) = \delta(r-ct)$. For $n \ge 1$, $\partial_r w^n = -2rnw^{n-1}$, and w^n vanishes on the cone, so no shell term appears. For the sum, $J_0' = -J_1$ and $\partial_r\sqrt{w} = -r/\sqrt{w}$. The absolute bound uses $\sum_{n\ge1} nx^{n-1}/(n!)^2 = I_1(2\sqrt{x})/\sqrt{x}$. A history with legs $\tau_i\hat{\omega}_i$ reaches $|x| = c\,|\sum \tau_i\hat{\omega}_i| < c\sum\tau_i$ unless all directions coincide, a set of measure zero for $n \ge 1$. □

7. Composition, unitarity, and the readout

Proposition 7.1 (The past enters through the pair)**.** *Let $S=0$ for $t \ge t_1$. For $t \ge t_1$, each spatial Fourier mode of the solution of Theorem 5.1 evolves by*

$$\begin{pmatrix}\hat{J}\\ \partial_t\hat{J}\end{pmatrix}(t) = M_k(t-t_1)\begin{pmatrix}\hat{J}\\ \partial_t\hat{J}\end{pmatrix}(t_1), \qquad M_k(t) = \begin{pmatrix}\cos\omega t & \omega^{-1}\sin\omega t\\ -\omega\sin\omega t & \cos\omega t\end{pmatrix},$$

with $\omega(k) = \sqrt{c^2k^2+\mu^2}$, and $M_k(t+t') = M_k(t)M_k(t')$.

Proof. For $t > t_1$, J solves $(\Box+\mu^2)J = 0$ (Theorem 5.1(2)), whose solutions are fixed by $(J, \partial_t J)$ on one slice. M_k is the flow of $\ddot{\hat{J}} = -\omega^2 \hat{J}$. □

Remark 7.2 (Not the faithful limit). The $\alpha = 2$ memory reaches over the whole past light cone, yet the past enters only through two accumulated quantities. At the edge the state was J alone, kept by the fundamental theorem of calculus [2, Prop. 6.1]. At $\alpha = 2$ it is the pair $(J, \partial_t J)$: in $1+1$ dimensions, one accumulator per null leg (Proposition 4.4). No records are kept, so this is not the faithful Markov limit, and the statement of [2] that a closed system cannot reach that limit is untouched.

Write $\mathcal{H} = L^2(\mathbb{R}^d) \otimes M_2(\mathbb{C})$, with the Hilbert–Schmidt product on $M_2(\mathbb{C})$. Let $\hat{H} = \sqrt{m^2c^4 - \hbar^2c^2\nabla^2}$, the self-adjoint operator of multiplication by $\hbar\omega(k)$ in Fourier space. A source-free solution has *positive frequency* if every Fourier mode is proportional to $e^{-i\omega(k)t}$, and *negative frequency* if every mode is proportional to $e^{+i\omega(k)t}$.

Theorem 7.3 (Unitarity on each sign of frequency; the trace readout). (1) *Positive-frequency solutions evolve by $J(t) = e^{-i\hat{H}t/\hbar}J(0)$, a strongly continuous unitary group on $\mathcal{H}$ [12, 13]. Negative-frequency solutions evolve by $e^{+i\hat{H}t/\hbar}$.*

(2) *On positive frequency the closure $Q(t) = \int JJ^\dagger\, d^dx \in M_2(\mathbb{C})$ is constant. So are $\operatorname{Tr} Q = \|J\|^2$, the readout $p_m = \operatorname{Tr}(P_m Q)/\operatorname{Tr} Q$ for projectors P_m on the left factor, and the normalization of the position readout $\int_\Delta \operatorname{tr}(JJ^\dagger)\, d^dx/\operatorname{Tr} Q$.*

(3) *On a solution with both signs of frequency, $\|J(t)\|^2$ is not constant: the cross terms oscillate at frequency 2ω.*

(4) *The Klein–Gordon charge*

$$q(J) = \frac{i\hbar}{2mc^2}\int \operatorname{tr}\left(J^\dagger \partial_t J - \partial_t J^\dagger J\right) d^dx$$

is conserved on every solution. It is positive on positive frequency, where $q(J) = \int(\hbar\omega/mc^2)\operatorname{tr}(\hat{J}^\dagger\hat{J})\, d^dk$, and negative on negative frequency. On positive frequency,

$$0 \le q(J) - \|J\|^2 \le \frac{\hbar^2}{2m^2c^2}\|\nabla J\|^2.$$

Proof. (1) Each mode has the phase $e^{\mp i\omega t}$, and $\hbar\omega(k)$ is real, so this is Stone's theorem for the multiplication operator $\hat{H}$.

(2) $Q_{ij} = \sum_l \langle J_{jl}, J_{il}\rangle_{L^2}$, and inner products are preserved by a unitary group that acts on the spatial factor only. The left projectors commute with that action.

(3) With $\hat{J} = a_+e^{-i\omega t} + a_-e^{i\omega t}$, $|\hat{J}|^2$ contains $2\mathrm{Re}\big(\bar{a}_+a_-e^{2i\omega t}\big)$.

(4) The charge is that of the conserved current of $(\Box+\mu^2)J = 0$. In Fourier space the cross terms of the two signs cancel in q, which leaves $\int(\hbar\omega/mc^2)(|a_+|^2 - |a_-|^2)$. Finally, $1 \le \hbar\omega/mc^2 = \sqrt{1+\hbar^2k^2/m^2c^2} \le 1 + \hbar^2k^2/2m^2c^2$. □

Remark 7.4 (Two conserved norms). On positive frequency, two positive quantities are conserved: the carrier trace $\|J\|^2 = \operatorname{Tr}\int JJ^\dagger$, and the Klein–Gordon charge q.

- q is the charge of a local conserved current, and its value is the same in every inertial frame.
- $\|J\|^2$ is conserved in each frame, but it is not the charge of a local current, and its value depends on the frame.

They differ at order v^2/c^2. So the trace readout of the closure $JJ^\dagger$ is a conserved, normalized Born readout in every fixed frame, on one sign of frequency. This is consistent with [2, §7.1]: the trace is a frame-dependent density, and the Lorentz-invariant structure sits in the determinant. A frame-independent position readout is the Newton–Wigner problem [11] and is not claimed.

8. The nonrelativistic limit: Feynman's free kernel

Fix m and $\hbar$, and let $c \to \infty$; then $\mu = mc^2/\hbar \to \infty$, and the kernel's lag scale $\tau_c = 1/(\sqrt{2}\mu)$ shrinks relative to any fixed comparison clock. Write a positive-frequency solution as $J(t) = e^{-imc^2 t/\hbar}\psi_c(t)$, and let

$$K_F(t) = e^{i\hbar t\nabla^2/2m}, \qquad K_F(t;x,x') = \Big(\frac{m}{2\pi i\hbar t}\Big)^{d/2} \exp\Big(\frac{im|x-x'|^2}{2\hbar t}\Big) \quad (t>0),$$

be Feynman's free kernel [4, 5].

Theorem 8.1 (Nonrelativistic limit)**.** *For $\psi \in H^4(\mathbb{R}^d) \otimes M_2(\mathbb{C})$,*

$$\big\| e^{-i(\hat{H}-mc^2)t/\hbar}\psi - K_F(t)\psi \big\| \le \frac{|t|\hbar^3}{8m^3c^2}\,\|\Delta^2\psi\|.$$

Both evolutions are unitary for every c.

Proof. With $x = \hbar^2k^2/m^2c^2 \ge 0$, $-x^2/8 \le \sqrt{1+x} - 1 - x/2 \le 0$. Hence $|\hbar\omega - mc^2 - \hbar^2k^2/2m| \le \hbar^4k^4/(8m^3c^2)$. Then use $|e^{ia} - e^{ib}| \le |a-b|$ mode by mode, and Plancherel. □

Corollary 8.2 (Feynman's kernel from the light-cone path sum)**.** *Let $[G_\mu]^+$ be the positive-frequency part of G_μ. In Fourier space $[\hat{G}_\mu]^+(t,k) = \theta(t)\,\frac{i}{2\omega}e^{-i\omega t}$, and*

$$\frac{2mc^2}{i\hbar}\, e^{imc^2t/\hbar}\,[G_\mu]^+(t) \;\longrightarrow\; K_F(t) \qquad (c\to\infty),$$

strongly on L^2. For $\psi \in H^4$ the error is at most $\frac{\hbar^2}{2m^2c^2}\|\Delta\psi\| + \frac{|t|\hbar^3}{8m^3c^2}\|\Delta^2\psi\|$. With Theorem 6.1, Feynman's free kernel is the limit of the sum over piecewise-null histories with amplitude $-\mu^2$ per corner,

$$K_F(t) = \lim_{c\to\infty} \frac{2mc^2}{i\hbar}\, e^{imc^2t/\hbar}\Big[\sum_{n\ge 0}(-\mu^2)^n\, G^{*(n+1)}(t)\Big]^+.$$

The composition $K_F(t+t') = K_F(t)K_F(t')$, which is Feynman's time slicing, is the limit of the group law of Theorem 7.3(1).

Proof. $\theta(t)\sin(\omega t)/\omega = \theta(t)\,(e^{i\omega t} - e^{-i\omega t})/2i\omega$, and its $e^{-i\omega t}$ part is $\theta(t)\,(i/2\omega)e^{-i\omega t}$. So the left side is the multiplier $(mc^2/\hbar\omega)\,e^{-i(\omega - mc^2/\hbar)t}$. Use $0 \le 1 - mc^2/\hbar\omega \le \hbar^2k^2/2m^2c^2$ and Theorem 8.1. □

Remark 8.3 (Where the i comes from). At the edge the i of the Schrödinger equation is the rotation produced by negative feedback on the accumulated past [2, Thm. 5.1]. At $\alpha = 2$ the dynamics is real: G, the insertion $-\mu^2$ and J_0 are all real. The i of K_F enters only through the sign of frequency. That sign is inherited from the source's past, as at the edge [2, Prop. 5.4], and nothing near $\alpha = 2$ selects it (Theorem 5.2). In $1+1$ dimensions it is also the corner amplitude whose square is the real insertion (Remark 6.3).

Remark 8.4 (Two routes to the Schrödinger equation). There are two routes:

- *The edge* [2]: the CAL iterated as a transducer at $\alpha \to 1^-$. The current rotates by $\sqrt{1-\alpha}$ per update, with $\hbar/2m = \langle r^2\rangle/12$ in units of the rest rotation. It is Galilean from the start, with no cone and no interaction.
- $\alpha = 2$: the balanced carrier at the light-cone width, on one sign of frequency, as $c \to \infty$ with $mc^2 = \hbar/(\sqrt{2}\,\tau_c)$. As $c \to \infty$ the past light cone opens to the whole time-ordered past, which is the causal structure of the edge [2, §7.1].

Both routes give the free Schrödinger equation. Whether the two masses coincide is not claimed. The condition that would tie them, a kernel width of $\sqrt{6}$ Compton wavelengths at the edge, is the equivalence condition stated as a forward pointer in [2, §9].

9. Numerical verification

All numbers come from `scripts/cal_feynman_alpha2.py` (numpy only, about 25 seconds). Units: $\hbar = c = 1$ and $\tau_c = 1/\sqrt{2}$, so that $\mu = 1$, unless stated.

Symbol and Lorentz invariance (Proposition 4.1, Theorem 4.2). Table 1 compares direct integration with the closed form at $(s, c|k|) = (0.8, 0.6)$, where $R = 1$, and lists h_α at four angles. The maximal Lorentz defect $\max_\varphi |h_\alpha(\varphi) - 1|$ over $(0, \pi/2]$ is

α	0	0.5	1.5	1.9	2	2.1	3
defect	0	4.1×10^{-1}	4.1×10^{-1}	9.7×10^{-2}	0	1.0×10^{-1}	1.0

The defect is zero only at $\alpha = 0$ and $\alpha = 2$, and is symmetric about $\alpha = 1$.

α	numeric	closed form	$h_\alpha(0.2)$	$h_\alpha(0.6)$	$h_\alpha(1.0)$	$h_\alpha(1.4)$
1.5	1.054093	1.054093	1.0050	1.0468	1.1395	1.3075
2.0	1.000000	1.000000	1.0000	1.0000	1.0000	1.0000
2.5	0.913547	0.913547	0.9917	0.9249	0.7903	0.5840
3.0	0.800000	0.800000	0.9801	0.8253	0.5403	0.1700

Table 1. The light-cone memory symbol: direct integration against the closed form, and the angular factor.

Dispersion (Theorem 5.1(3), Remark 3.1). With $\tau_c = 1/\sqrt{2}$, so $\mu^2 = 1/(2\tau_c^2) = 1$, the zero of the retarded symbol of T is found by bisection.

k	root E	$\sqrt{k^2 + \mu^2}$	E^2 with an added factor -1
0.0	1.0000000000	1.0000000000	-1.00
0.5	1.1180339887	1.1180339887	-0.75
1.0	1.4142135624	1.4142135624	0.00
2.0	2.2360679775	2.2360679775	3.00
4.0	4.1231056256	4.1231056256	15.00

Table 2. Klein–Gordon dispersion of the balanced $\alpha = 2$ law. An added factor -1 on the memory, which is not in the CAL, would make the mass tachyonic.

Near $\alpha = 2$ (Theorem 5.2). The roots are found by Newton iteration on the continued symbol, at $k = 1$. Growing modes are counted by the winding number of the symbol around the rectangle $\operatorname{Im}\omega \in [0.002, 12]$.

Path sum in $1 + 1$ (Proposition 6.2(1)). The terms $G^{*(n+1)}$ are built by iterated trapezoid accumulation along u and then v, one insertion per step, on $u, v \in [0, 12]$. The partial sums are compared with $\frac{1}{2} J_0(\sqrt{uv})$.

Feynman's checkerboard (Proposition 6.2(3)). The lattice amplitude is propagated with 1 for keeping direction and $i\varepsilon\mu$ for a corner. The first leg moves right, and the amplitude arriving left-moving at $t = 8$ is compared with $i\varepsilon\mu\, J_0$ on $|x| \le 7.5$.

Descent to $3 + 1$ (Proposition 6.4). The $1 + 1$ sum with $n \le 60$, minus its $n = 0$ term, is differentiated on the (u, v) grid ($h = 0.01$), and $-\partial_r/(2\pi r)$ is applied with $\partial_r = \partial_v - \partial_u$. The result is compared with $-\mu J_1(\mu\sqrt{w})/(4\pi\sqrt{w})$ at points with $r \ge 0.25$ and $t - r \ge 0.25$.

Composition, unitarity and the nonrelativistic limit (Proposition 7.1, Theorems 7.3 and 8.1).

- Pair composition: $\max \|M_k(t_1 + t_2) - M_k(t_2) M_k(t_1)\| = 1.1 \times 10^{-14}$ over 200 random modes, with $t_1 = 0.7$ and $t_2 = 1.9$.

α	root (+ branch)	$\operatorname{Im}\omega$, both branches	$(\pi/4)\lvert\alpha-2\rvert\mu^2/\lvert\omega\rvert$	growing modes, $k=0.1,1,5$
1.90	1.370022	-5.435×10^{-2}	5.03×10^{-2}	0, 0, 0
1.99	1.410296	-5.546×10^{-3}	5.51×10^{-3}	0, 0, 0
2.00	1.414214	0	0	–
2.01	1.418025	$+5.560\times10^{-3}$	5.60×10^{-3}	2, 2, 0
2.05	1.432250	$+2.790\times10^{-2}$	2.88×10^{-2}	2, 2, 2
2.10	1.447853	$+5.590\times10^{-2}$	5.95×10^{-2}	2, 2, 2

TABLE 3. Theorem 5.2. At $\alpha=2.01$ and $k=5$ the growth rate, 1.5×10^{-3}, is below the counting threshold. A time-stepped run with a real source and zero past gives $|J|=8.0\times10^{-2}\to1.4\times10^{-3}$ between $t=50$ and 200 at $\alpha=1.95$; bounded beats at $\alpha=2$; and $6.3\times10^{-1}\to5.9\times10^{1}$ at $\alpha=2.05$.

grid h	$n=2$	$n=5$	$n=10$	$n=20$	$n=40$	$n=60$
0.020	1.44×10^{2}	1.15×10^{3}	3.28×10^{1}	3.35×10^{-5}	3.35×10^{-5}	3.35×10^{-5}
0.010	1.44×10^{2}	1.15×10^{3}	3.28×10^{1}	8.38×10^{-6}	8.38×10^{-6}	8.38×10^{-6}

TABLE 4. Maximum error of the partial path sums. The large intermediate values are those of the J_0 power series at $z=12$. The residual falls by 4.0 when h halves, so it is discretization, not the series.

ε	$\max\lvert L_\varepsilon/(i\varepsilon\mu)-J_0\rvert$	ratio to previous
0.02	3.697×10^{-2}	–
0.01	1.810×10^{-2}	2.04
0.005	8.950×10^{-3}	2.02
0.0025	4.451×10^{-3}	2.01

TABLE 5. The checkerboard's corner-odd amplitude converges to $2c\,i\mu\,G_\mu=i\mu J_0$ at first order in ε.

t	max error	max|exact|	points
4	5.28×10^{-7}	3.090×10^{-2}	351
8	2.46×10^{-6}	2.330×10^{-3}	376

TABLE 6. The interior tail of the $3+1$ massive propagator, built from the summed histories.

- Packet $\psi\propto e^{-x^2/8+ix}$ evolved to $t=10$ by $e^{-i(\hat H-mc^2)t/\hbar}$ and by K_F, with $\hbar=m=1$ (Table 7). The predicted limit of the error times c^2 is $(t/8)\|k^4\hat\psi\|=3.672$.
- Trace readout: over $t\in[0,10]$ the relative spread of $\|J(t)\|^2$ is 2.5×10^{-16} on positive frequency and 1.6×10^{-1} on a mixed solution. The relative spread of the Klein–Gordon charge of the mixed solution is 4.4×10^{-16}.

10. Remarks on scope

Remark 10.1 (What is derived and what is not). In summary:

- *Derived from the axioms:* Lorentz invariance of the memory exactly at $\alpha=2$; the Klein–Gordon equation and the mass as the kernel's own rate; conservation exactly at $\alpha=2$,

c	$\|\mathrm{KG} - \mathrm{Schrödinger}\|$	error $\times c^2$	norm drift
2	6.058×10^{-1}	2.423	0
4	2.076×10^{-1}	3.321	0
8	5.608×10^{-2}	3.589	2.2×10^{-16}
16	1.426×10^{-2}	3.651	2.2×10^{-16}
32	3.581×10^{-3}	3.667	2.2×10^{-16}

TABLE 7. Positive-frequency Klein–Gordon evolution, rest phase removed, against Feynman's free kernel.

with decay below and runaway above; absolute convergence of the history sum and its resummation; the checkerboard in $1+1$; the corner tail in $3+1$; the pair composition.

- *Derived under the one-sign restriction:* unitarity in $\mathcal{H}$, conservation of the closure $\int JJ^\dagger$ and of its trace readout, and the limit to Feynman's kernel.
- *Reading:* the balance reading at $\alpha = 2$ (Axiom 3); the edge's update rule is not used.
- *Open:*
  - the sign of frequency is an input, inherited from the source's past, as at the edge;
  - interaction between carriers at $\alpha = 2$, where the reciprocal (return-leg) structure that the edge lacks is available [2, Rem. 7.1];
  - a first-order (spin) form in $3+1$ dimensions;
  - the relation between the $\alpha = 2$ mass and the edge's rest rotation, and mass ratios between carriers;
  - a frame-independent position readout.

Remark 10.2 (Histories are memory). Each term of the Feynman sum is the past re-entering the present n times along the light cone, weighted by the order-two lag τ per leg and by $-\mu^2$ per re-entry. The sum over histories is the memory series of the CAL at $\alpha = 2$. Its Markov form is the pair $(J, \partial_t J)$, and its Galilean limit is Feynman's kernel.

11. Conclusion

At closure order $\alpha = 2$, a memory spread over the light cone is Lorentz invariant, and it is so only at that order. Its kernel is the retarded Green's function of the wave operator: Kirchhoff's shell in $3+1$ dimensions, and one order-one accumulation per null leg in $1+1$.

With the transport chain the identity and no factor on the memory, a carrier whose present balances its light-cone past obeys the Klein–Gordon equation. Its mass is the kernel's own rate, and only ratios between carriers are physical. The law is conservative exactly at $\alpha = 2$, decays below it, and runs away above it.

The balance equation is a sum over piecewise-null histories, with $-\mu^2$ at each corner. In $1+1$ dimensions it is Feynman's checkerboard term by term, and in $3+1$ dimensions the corners alone fill the cone. The past enters only through the pair $(J, \partial_t J)$.

Each sign of frequency is unitary. The sign is inherited from the source's past, and on it the carrier's trace readout is conserved. In the slow limit the positive-frequency path sum is Feynman's free kernel. The dynamics is real throughout, and the i appears only with the inherited sign of frequency.

Appendix A. Notation

α	closure order
ν_τ	light-cone profile at lag τ, radius $c\tau$ (Definition 2.1)
K_α, m_α	order-α light-cone memory and its symbol
R, φ, h_α	polar form of $(s, c\lvert k\rvert)$, and angular factor
$\Box = \partial_t^2 - c^2\nabla^2$	wave operator
G, G_μ	retarded Green's functions of $\Box$ and $\Box + \mu^2$
τ_c	the kernel's lag scale, the only scale
$\mu = 1/(\sqrt{2}\,\tau_c)$, $mc^2 = \hbar\mu$	rest rate and mass
$u = ct - x$, $v = ct + x$	null coordinates in $1+1$
Φ_{2n+1}	checkerboard amplitude with $2n+1$ corners
$\mathcal{D}$	descent operator (Proposition 4.5)
$w = c^2t^2 - r^2$	interval inside the cone
$\hat{H} = \sqrt{m^2c^4 - \hbar^2c^2\nabla^2}$	positive-frequency generator
$Q = \int JJ^\dagger$, q	closure of the carrier; Klein–Gordon charge
K_F	Feynman's free kernel

Appendix B. Numerical methods

Symbol. The integral of Definition 2.3 is computed on a logarithmic grid $\tau = e^u$, $u \in [-14, 8]$, with 4×10^5 trapezoid intervals.

Dispersion. The real part of the retarded symbol of T, at $\operatorname{Re} s = 10^{-12}$, is bisected 200 times on $(|k|, 50)$.

Path sums. $G^{*(n+1)}$ is computed on the (u, v) grid by cumulative trapezoid integration in u and then in v, times $1/4$ ($c = 1$). J_0, J_1 and I-type checks use the power series with 160 terms.

Checkerboard. Two amplitude arrays (arriving right-moving and left-moving) are updated per time step: $R_{j+1} \leftarrow R_j + i\varepsilon\mu L_j$ and $L_{j-1} \leftarrow L_j + i\varepsilon\mu R_j$. Only lattice sites of the right parity are compared.

Descent. Central differences on the (u, v) grid, along the diagonal $u + v = 2t$.

Packets. A periodic grid of 4096 points on a box of length 400, with a unitary FFT, and exact spectral evolution.

Near $\alpha = 2$.

- Roots: Newton iteration on the continued closed-form symbol.
- Growing modes: the winding number on a rectangle 2×10^5 points per side, with its lower edge at $\operatorname{Im}\omega = 0.002$, which avoids the light-cone branch points.
- Time stepping: the trapezoid product rule on the lag integral, with step 0.02.

References

[1] S. Leizerman, *The Causal Accumulation Law: The Four Corners of α, Algebraic Closure, and the Standard Model*, LIOR Research Initiative (2026).

[2] S. Leizerman, *Schrödinger evolution from the bare Causal Accumulation Law at $\alpha \to 1^-$: feedback on the accumulated past, unitarity in the limit, and carrier entanglement as time dilation*, LIOR Research Initiative (2026).

[3] S. Leizerman, *An Axiomatic Derivation of the Born Rule at the δ-Kernel Limit and Closure Order $\alpha = 2$ of the Causal Accumulation Law*, LIOR Research Initiative (2026).

[4] R. P. Feynman, Space-time approach to non-relativistic quantum mechanics, Rev. Mod. Phys. **20**, 367–387 (1948).

[5] R. P. Feynman, A. R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw-Hill (1965), Problem 2-6.

[6] T. Jacobson, L. S. Schulman, Quantum stochastics: the passage from a relativistic to a non-relativistic path integral, J. Phys. A **17**, 375–383 (1984).

[7] B. Gaveau, T. Jacobson, M. Kac, L. S. Schulman, Relativistic extension of the analogy between quantum mechanics and Brownian motion, Phys. Rev. Lett. **53**, 419–422 (1984).
[8] S. Johnston, Particle propagators on discrete spacetime, Class. Quantum Grav. **25**, 202001 (2008).
[9] R. Courant, D. Hilbert, *Methods of Mathematical Physics, Vol. II*, Interscience (1962).
[10] P. A. M. Dirac, Classical theory of radiating electrons, Proc. R. Soc. Lond. A **167**, 148–169 (1938).
[11] T. D. Newton, E. P. Wigner, Localized states for elementary systems, Rev. Mod. Phys. **21**, 400–406 (1949).
[12] M. H. Stone, On one-parameter unitary groups in Hilbert space, Ann. of Math. **33**, 643–648 (1932).
[13] M. Reed, B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, Academic Press (1980).

LIOR Research Initiative
Email address: samleizerman@outlook.com