

Unit-1

One Words

- 1) Each performance of random experiment is called a trial.
- 2) The collection of all possible outcomes of a random exp is called the sample space.
- 3) sample space is denoted by S .
- 4) The elements of sample space are called sample points.
- 5) The outcome of "head turning up" is denoted by H .
- 6) The outcome of "tail turning up" is denoted by T .
- 7) Any subset A of sample space S is called an event.
- 8) The event S is called a pure event.
- 9) The event ϕ is called an impossible event.

ucted highways
ways, project
Certs remain

Roads on a roll
Construction of roads

ecture: Finish line yet to be reached

BUDGET EXPECTATIONS



(4)

- 2) $A = \{1, 3, 5\}$ is the event of getting an odd number.
- 1) $B = \{2, 4, 6\}$ is the event of getting an even number.
- 2) $\bar{A}, \bar{B}, \bar{C}$ are called complement events.
- 3) f/N is called the relative frequency of the event A .
- 4) $C_1: P(A) \geq 0$ for all $A \subseteq S$.
- 5) $C_2: P(S) = 1$
- 6) $C_3: \text{If } \{A_n\}$ is any finite (or) infinite seq of disjoint events then $P(\cup A_i) = \sum P(A_i)$.
- 7) P is called the Probability set function.
- 8) The number $P(A)$ is called the Probability of the event A .
- 9) $P(A) = \frac{\text{No. of cases favourable to } A}{\text{Total no. of cases}}$
- 10) If $A_i \cap A_j = \emptyset \neq i, j$ with $i \neq j$ then seq. of subsets is said to be mutually disjoint.
- 11) If $\bigcup_{n=1}^{\infty} A_n = S$, then seq. of events is said to be exhaustive.

5

- 22) If seq of events $A_1, A_2 \dots A_n$ is mutually disjoint & exhaustive, then $P(\bigcup_{n=1}^{\infty} A_n) = \sum_{n=1}^{\infty} P(A_n) = 1$.
- 23) $P(A/B) = \frac{P(A \cap B)}{P(B)}$
- 24) $P(A/B) \geq 0$
- 25) $P(A/A) = 1$
- 26) $P(A_1 \cup A_2 \cup \dots / B) = P(A_1/B) + P(A_2/B) + \dots$ provide $A_1, A_2 \dots$ are mutually disjoint events.
- 27) $P(A \cap B) = P(B) P(A/B)$ is called multiplication theorem for probabilities.
- 28) A is independent of B if $P(A/B) = P(A)$.
- 29) If A & B are 2 independent events, then $P(A \cap B) = P(A) P(B)$.
- 30) A set of events $A_1, A_2 \dots A_n$ are said to be pairwise independent if $P(A_i \cap A_j) = P(A_i) P(A_j)$.
- 31) The events $A_1, A_2 \dots A_n$ are independent if $P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) P(A_2) \dots P(A_n)$.
- 32) pairwise independent of n events \neq independence of n events.
- 3) If A & B are independent events then $A \cap \bar{B}$ are also independent events.

Unit-2

- 1) A fn $x: S \rightarrow \mathbb{R}$ which assigns to each elmt $\omega \in S$ one & only one real no's is called a random variable.
- 2) The space of random variable x is defined to be set of real no's $\xi = \{x(\omega) / \omega \in S\}$.
- 3) P' defines a probability set fn on ξ .
- 4) P' is called the induced probability of random variable x .
- 5) The fn $F: \mathbb{R} \rightarrow \mathbb{R}$ defined by $F(x) = P(X \leq x)$, where $-\infty < x < \infty$ is called distribution fn of random variable x .
- 6) If $F(x)$ is distribution fn of random vble x & if $a < b$ then $P(a < x \leq b) = F(b) - F(a)$.
- 7) The events $a < x \leq b$ & $x \leq a$ are disjoint.
- 8) $P(X = x_i)$ is also written as $P(x_i)$ or simply P_i .
- 9) If random vble takes atmost countable no. of values x_1, x_2, \dots, x_n it is called a discrete random vble.

$$\sum P_i = 1$$

11) Any subset A of \mathcal{S} , $P(A) = \sum_{x_i \in A} P(x_i)$.

$$12) P(x = x_i) = f(x_i) = P_i.$$

$$13) F(x) = \sum_{x_i < x} P_i = \sum_{x_i \leq x} f(x_i).$$

14) f is called probability density fn of discrete random vble.

15) A random vble x is said to be continuous random vble if it can take any value in an interval which may be finite or infinite.

$$16) P(A) = \int_a^b f(x) \cdot dx.$$

17) $P(A)$ is also written as $P(A) = P(a < x < b)$.

$$18) A = \{a\}.$$

$$19) P(A) = \int_a^a f(x) dx = 0$$

20) x is continuous random vble, probability of every set consisting of single pt is zero.

$$21) F(x) = \int_{-\infty}^x f(t) \cdot dt.$$

22) $F(x)$ is called distribution fn of continuous random variable x .

1) $f'(x) = f(x)$ at each pt where $f(x)$ is continuous.

2) $F(0) = 1$

3) $F(-\infty) = 0$

4) $P(a \leq x \leq b) = F(b) - F(a)$

5) $F(x)$ is an increasing fn of x .

6) $P_i = P(x = x_i), i = 1, 2, \dots$

7) Mathematical expectation of x , denoted by $E(x)$.

8) $E(x)$ is denoted by $\sum P_i x_i$.

9) It is easy to verify the following results if x & y are random vbls.

32) $E(c) = c$ where c is constant.

33) $E(cx) = c E(x)$.

34) $E(ax + b) = a E(x) + b$.

35) $E(x + y) = \underline{E(x) + E(y)}$

36) $E(xy) = E(x)E(y)$ if x & y are independent random vbls.

37) $E(\psi(x)) = \sum P_i \psi(x_i)$.

6

Statistics-II

Bayes's theorem:-

Let $\{A_i\}$ be a sequence of mutually exclusive & exhaustive events in a sample space S s.t. $P(A_i) > 0$ for all i . Let B be any event with $P(B) > 0$ then,

$$P(A_i/B) = \frac{P(A_i) P(B/A_i)}{\sum P(A_i) P(B/A_i)}$$

Pf: By Bayes's rule we have,

$$P(A_i/B) = \frac{P(A_i) P(B/A_i)}{P(B)} \quad \text{--- (1)}$$

we claim that,

$$P(B) = \sum_i P(A_i) P(B/A_i)$$

Since the events $\{A_i\}$ are mutually exclusive & exhaustive, we have,

$$\bigcup_i A_i = S \text{ \& } A_i\text{'s are disjoint.}$$

$$\therefore B = B \cap S$$

$$= B \cap \left(\bigcup_i A_i \right) \quad [\because A_i\text{'s are exhaustive}]$$

$$B = \bigcup_i (B \cap A_i)$$

$$P(B) = P\left(\bigcup_i (B \cap A_i)\right)$$

$$P(B) = \sum_i P(B \cap A_i) \quad [A_i\text{'s are mutually disjoint}]$$

$$= \sum_i P(A_i) P(B/A_i) \quad [\because P(B/A_i) = \frac{P(B \cap A_i)}{P(A_i)}]$$

Then, $P(B) = \sum_i P(A_i) P(B/A_i) \quad \text{--- (2)}$

sub (2) in (1)

$$P(A_i/B) = \frac{P(A_i) P(B/A_i)}{\sum_i P(A_i) P(B/A_i)} //$$

2) Boole's inequality:-

If A & B are any 2 events in a sample space then P.T. i) $P(A \cap B) \geq 1 - P(\bar{A}) - P(\bar{B})$ ii) (Generalised Boole's inequality). For the events $A_1, A_2, \dots, A_n, \dots$ is a sample space $P(\bigcap_{i=1}^{\infty} A_i) \geq 1 - \sum_{i=1}^{\infty} P(\bar{A}_i)$.

Proof: i) For any 2 events A & B we have,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\begin{aligned} P(A \cap B) &= [1 - P(\bar{A})] + [1 - P(\bar{B})] - P(A \cup B) \\ &= 1 - P(\bar{A}) - P(\bar{B}) + [1 - P(A \cup B)] \\ &\geq 1 - P(\bar{A}) - P(\bar{B}) \quad (\because 0 \leq P(A \cup B) \leq 1) \end{aligned}$$

ii) Let $B_1 = \bigcap_{i=2}^{\infty} A_i$ so that $\bigcap_{i=1}^{\infty} A_i = A_1 \cap B_1$

$$\text{Now, } P\left(\bigcap_{i=1}^{\infty} A_i\right) = P(A_1 \cap B_1) \geq 1 - P(\bar{A}_1) - P(\bar{B}_1)$$

(by i))

$$= 1 - P(\bar{A}_1) - P\left(\overline{\bigcap_{i=2}^{\infty} A_i}\right)$$

$$\begin{aligned}
 &= 1 - P(\bar{A}_1) - P\left(\bigcup_{i=2}^{\infty} \bar{A}_i\right) \geq 1 - P(\bar{A}_1) - \sum_{i=2}^{\infty} P(\bar{A}_i) \\
 &= 1 - \sum_{i=1}^{\infty} P(\bar{A}_i) \\
 &= P\left(\bigcap_{i=1}^{\infty} A_i\right) \geq 1 - \sum_{i=1}^{\infty} P(\bar{A}_i) //
 \end{aligned}$$

Unit - 1

One words

6

Infrastructure: India

Though the govt has constructed highways and developed inland waterways, project funding and road safety concerns remain

MAHENDRA DAS

New Delhi, India

9,800-

tion of

financ

the main

any of the

"Between

2018, 6,700

were consi

cal) year, v

more than

built about

to March 15

linum wor

Transport

istry officia

But thea

may be to

pared to th

scious stra

when they

complete

tion of lan

ance," sha

The con

ment-fun

changed

awarded

struction

to imp

roads, t

moved fr

ive criter

ment-bas

Buildin

express

routes v

cost will

scious st

ways, lar

tioned

outlets

gates

gates

gates