

# Supervised Learning (COMP0078) – Coursework 1

Mark Herbster

Due : 14 November 2018.

## Submission

You may work in groups of up to two.<sup>1</sup> You should produce a report about your results. You will not only be assessed on the **correctness/quality** of your answers but also on **clarity of presentation**. Additionally make sure that your code is *well commented*. Please submit on moodle i) your report as well as a ii) zip file with your source code. Finally, please ensure that if you are working in a group both of your names are on the coversheet. Regarding the use of libraries, you should implement regression using matrix algebra directly. Otherwise libraries are okay.

Questions please e-mail [sl-support@cs.ucl.ac.uk](mailto:sl-support@cs.ucl.ac.uk) .

## 1 PART I [50%]

### 1.1 Linear Regression

“Pluralitas non est ponenda sine neccesitate” – William of Ockham (ca. 1285-1349)

**Linear regression overview:** Given a set of data:

$$\{(\mathbf{x}_1, y_1), (\mathbf{x}_2, y_2), \dots, (\mathbf{x}_m, y_m)\} \quad (1)$$

where  $\mathbf{x} = (x_1, \dots, x_n)$  is a vector in  $\mathbb{R}^n$  and  $y$  is a real number. Linear regression finds a vector  $\mathbf{w} \in \mathbb{R}^n$  such that the sum of squared errors

$$\text{SSE} = \sum_{t=1}^m (y_t - \mathbf{w} \cdot \mathbf{x}_t)^2 \quad (2)$$

is minimized. This is expressible in matrix form by defining  $X$  to be the  $m \times n$  matrix

$$X = \begin{pmatrix} x_{1,1} & x_{1,2} & \dots & x_{1,n} \\ x_{2,1} & x_{2,2} & \dots & x_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m,1} & x_{m,2} & \dots & x_{m,n} \end{pmatrix}, \quad (3)$$

and defining  $\mathbf{y}$  to be the column vector  $\mathbf{y} = (y_1, \dots, y_m)$ . The vector  $\mathbf{w}$  then minimizes

$$(\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}).$$

In linear regression with basis functions we fit the data sequence with a linear combination of basis functions  $(\phi_1, \phi_2, \dots, \phi_k)$  where  $\phi_i : \mathbb{R}^n \rightarrow \mathbb{R}$  which defines a feature map from  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^k$

$$\phi(\mathbf{x}) = (\phi_1(\mathbf{x}), \dots, \phi_k(\mathbf{x})), \quad \mathbf{x} \in \mathbb{R}^n.$$

We use the basis functions to transform the data as follows

$$\{((\phi_1(\mathbf{x}_1), \dots, \phi_k(\mathbf{x}_1)), y_1), \dots, ((\phi_1(\mathbf{x}_m), \dots, \phi_k(\mathbf{x}_m)), y_m)\}, \quad (4)$$

---

<sup>1</sup>Affiliates students should work alone.

and then applying linear regression above to this transformed dataset. In matrix notation we may denote this transformed dataset as


$$\Phi := \begin{pmatrix} \phi(\mathbf{x}_1) \\ \vdots \\ \phi(\mathbf{x}_m) \end{pmatrix} = \begin{pmatrix} \phi_1(\mathbf{x}_1) & \dots & \phi_k(\mathbf{x}_1) \\ \vdots & \ddots & \vdots \\ \phi_1(\mathbf{x}_m) & \dots & \phi_k(\mathbf{x}_m) \end{pmatrix}, \quad (m \times k) \quad (5)$$

Linear regression on the transformed dataset thus finds a  $k$ -dimensional vector  $\mathbf{w} = (w_1, \dots, w_k)$  such that

$$(\Phi \mathbf{w} - y)^T (\Phi \mathbf{w} - y) = \sum_{t=1}^m (y_t - \sum_{i=1}^k w_i \phi_i(\mathbf{x}_t))^2 \quad (6)$$

is minimized.

A common basis ( $n = 1$ ) used is the polynomial basis  $\{\phi_1(x) = 1, \phi_2(x) = x, \phi_3(x) = x^2, \phi_4(x) = x^3, \dots, \phi_k(x) = x^{k-1}\}$  of dimension  $k$  (order  $k - 1$ ) in the figure below we give a simple fit of four points produced by a linear ( $k = 2$ ) and cubic ( $k = 4$ ) polynomial.



**Figure 1: Data set  $\{(1, 3), (2, 2), (3, 0), (4, 5)\}$  fitted with basis  $\{1, x\}$  and basis  $\{1, x, x^2, x^3\}$**


1. **[5 pts]:** For each of the polynomial bases of dimension  $k = 1, 2, 3, 4$  fit the data set of Figure 1  $\{(1, 3), (2, 2), (3, 0), (4, 5)\}$ .
  - (a) Produce a plot similar to Figure 1, superimposing the four different curves corresponding to each fit over the four data points.
  - (b) Give the equations corresponding to the curves fitted for  $k = 1, 2, 3$ . The equation corresponding to  $k = 4$  is  $-5 + 15.17x - 8.5x^2 + 1.33x^3$ .
  - (c) For each fitted curve  $k = 1, 2, 3, 4$  give the mean square error where  $\text{MSE} = \frac{\text{SSE}}{m}$ .
2. **[10 pts]:** In this part we will illustrate the phenomena of *overfitting*.
  - (a) Define

$$g_\sigma(x) := \sin^2(2\pi x) + \epsilon. \quad (7)$$

where  $\epsilon$  is a random variable distributed normally with mean 0 and variance  $\sigma^2$  thus  $g_\sigma(x)$  is a *random* function such that  $\sin^2(2\pi x)$  is computed and then the normal noise is added on each “call” of the function. We then sample “ $x_i$ ” uniformly at random from the interval  $[0, 1]$  30 times creating  $(x_1, \dots, x_{30})$  and apply  $g_{0.07}$  to each  $x$  creating the data set

$$S_{0.07, 30} = \{(x_1, g_{0.07}(x_1)), \dots, (x_{30}, g_{0.07}(x_{30}))\}. \quad (8)$$

- i. Plot the function  $\sin^2(2\pi x)$  in the range  $0 \leq x \leq 1$  with the points of the above data set superimposed. The plot should resemble



- ii. Fit the data set with a polynomial bases of dimension  $k = 2, 5, 10, 14, 18$  plot each of these 5 curves superimposed over a plot of data points.<sup>2</sup>
- (b) Let the training error  $te_k(S)$  denote the MSE of the fitting of the data set  $S$  with polynomial basis of dimension  $k$ . Plot the log of the training error versus the polynomial dimension  $k = 1, \dots, 18$  (this should be a decreasing function).
- (c) Generate a test set  $T$  of a thousand points,

$$T_{0.07,1000} = \{(x_1, g_{0.07}(x_1)), \dots, (x_{1000}, g_{0.07}(x_{1000}))\}. \quad (9)$$

Define the test error  $tse_k(S, T)$  to be the MSE of the test set  $T$  on the polynomial of dimension  $k$  fitted from training set  $S$ . Plot the log of the test error versus the polynomial dimension  $k = 1, \dots, 18$ . Unlike the training error this is not a decreasing function. This is the phenomena of *overfitting*. Although the training error decreases with growing  $k$  the test error eventually increases since rather than fitting the function, in a loose sense, we begin to fit to the noise.

- (d) For any given set of random numbers we will get slightly different training curves and test curves. It is instructive to see these curves smoothed out. For this part repeat items (b) and (c) but instead of plotting the results of a single “run” plot the average results of a 100 runs (note: plot the  $\log(\text{avg})$  rather than the  $\text{avg}(\log)$ ).

3. [5 pts]: Now use basis (for  $k = 1, \dots, 18$ )

$$\sin(1\pi x), \sin(2\pi x), \sin(3\pi x), \dots, \sin(k\pi x).$$

Repeat the experiments in 2 (b-d) with the above basis.

## 1.2 Boston housing and kernels

In this section we will use kernel methods to extend linear regression. Boston housing is a classic dataset where you are given 13 values and a goal is to predict the 14th which is the median house price. It is described in more detail at <http://www.cs.toronto.edu/~dave/data/boston/bostonDetail.html>. There are 506 entries which we will split into train and test.

The boston housing data set as “.mat” file is located at

<http://www.cs.ucl.ac.uk/staff/M.Herbster/boston>

otherwise please go to URL above to retrieve in as a text file.

4. [10 pts]: “Baseline versus full linear regression.”  
Rather than use all of our attributes for prediction it is often useful to see how well a baseline method works for a problem. In this exercise, we will compare the following:

<sup>2</sup>Depending on your implementation you may have numerical errors for large values of  $k$  this is ‘ok’ for the purposes of this exercise.

- (a) Predicting with the mean  $y$ -value on the training set.
- (b) Predicting with a single attribute and a bias term.
- (c) Predicting with all the attributes

The training set should be 2/3, and the test set should be 1/3, of your data in (a)-(c). In the following average your results over 20 runs (each run based on a different (2/3,1/3) random split).

- a. Naive Regression. Create a vector of ones that is the same length as the training set using the function `ones`. Do the same for the test set. By using these vectors we will be fitting the data with a constant function. Perform linear regression on the training set. Calculate the MSE on the training and test sets and note down the results.
- b. Give a simple interpretation of the constant function in ‘a.’ above.
- c. Linear Regression with single attributes. For each of the thirteen attributes, perform a linear regression using only the single attribute but incorporating a bias term so that the inputs are augmented with an additional 1 entry,  $(\mathbf{x}_i, 1)$ , so that we learn a weight vector  $\mathbf{w} \in \mathbb{R}^2$ .
- d. Linear Regression using all attributes. Now we would like to perform linear regression using all of the data attributes at once.

Perform linear regression on the training set using this regressor, and incorporate a bias term as above. Calculate the MSE on the training and test sets and note down the results. You should find that this method outperforms any of the individual regressors.

### 1.3 Kernelised ridge regression

For nonlinear regression, the dual version will prove important. Obviously, linear regression is not capable of achieving a good predictive performance on a nonlinear data set. Here, the dual formulation will prove extremely useful, in combination with the *kernel trick*.

For our exercises we will use a slight variation on the optimisation (as compared to the notes) that defines ridge regression for a training set with  $\ell$  examples,

$$\mathbf{w}^* = \operatorname{argmin}_{\mathbf{w} \in \mathbb{R}^n} \frac{1}{\ell} \sum_{i=1}^{\ell} (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 + \gamma \mathbf{w}^\top \mathbf{w}. \quad (10)$$

i.e, we have replace the sum of the square errors with the mean square error (MSE).<sup>3</sup> For a given kernel function  $K$  define the kernel matrix  $\mathbf{K}$  for a training set of size  $\ell$  elementwise via

$$K_{i,j} := K(\mathbf{x}_i, \mathbf{x}_j)$$

The dual optimisation formulation after kernelization is

$$\boldsymbol{\alpha}^* = \operatorname{argmin}_{\boldsymbol{\alpha} \in \mathbb{R}^\ell} \frac{1}{\ell} \sum_{i=1}^{\ell} \left( \sum_{j=1}^{\ell} \alpha_j K_{i,j} - y_i \right)^2 + \gamma \boldsymbol{\alpha}^\top \mathbf{K} \boldsymbol{\alpha}. \quad (11)$$

Now if we use  $\mathbf{y} := (y_1, \dots, y_\ell)^\top$  to denote a vector that contain the  $y$ -values of training set we may solve in the dual as follows

$$\boldsymbol{\alpha}^* = (\mathbf{K} + \gamma \ell \mathbf{I}_\ell)^{-1} \mathbf{y} \quad (12)$$

where  $\mathbf{I}_\ell$  denotes the  $\ell \times \ell$  identity matrix. The evaluation of the regression function on a test point can be reformulated as:

$$y_{\text{test}} = \sum_{i=1}^{\ell} \alpha_i^* K(\mathbf{x}_i, \mathbf{x}_{\text{test}}) \quad (13)$$

---

<sup>3</sup>The motivation is that we wish the regularisation parameter  $\gamma$  to ‘scale’ somewhat independently of training set size.

where the  $K$  is the kernel function.

5. [20 pts] “Kernel Ridge Regression”

In this exercise we will perform kernel ridge regression (KRR) on the data set using the Gaussian kernel,

$$K(\mathbf{x}_i, \mathbf{x}_j) = \exp\left(-\frac{\|\mathbf{x}_i - \mathbf{x}_j\|^2}{2\sigma^2}\right). \quad (14)$$

For this exercise, you will hold out 2/3 of data for training and report the test results on the remaining 1/3.

- Create a vector of  $\gamma$  values  $[2^{-40}, \dots, 2^{-26}]$  and a vector of  $\sigma$  values  $[2^7, 2^{7.5}, \dots, 2^{12.5}, 2^{13}]$  (recall that  $\sigma$  is the variance parameter in the Gaussian kernel see equation (14)). Perform kernel ridge regression on the training set using five-fold cross-validation to choose among all pairing of the values of  $\gamma$  and  $\sigma$ . Choose the indices of the  $\gamma$  and  $\sigma$  values that perform the best to compute the predictor that you will use to report the test and training error.
- Plot the “cross-validation error” (mean over folds of validation error) as a function of  $\gamma$  and  $\sigma$ .
- Calculate the MSE on the training and test sets for the best  $\gamma$  and  $\sigma$ .
- Repeat “exercise 4a,c,d” and “exercise 5c” over 20 random (2/3, 1/3) splits of your data record the train/test error and the standard deviations of the train/test errors and summarise these results in the following type of table.

Method	MSE train	MSE test
Naive Regression	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 1)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 2)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 3)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 4)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 5)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 6)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 7)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 8)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 9)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 10)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 11)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 12)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (attribute 13)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Linear Regression (all attributes)	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$
Kernel Ridge Regression	?? ?? $\pm \sigma$	?? ?? $\pm \sigma$

## 2 PART II [50%]

### 2.1 Questions

**Notation:** We overload  $[\cdot]$  as follows  $[n] := \{1, 2, \dots, n\}$  if  $n$  is a positive integer and  $[\text{pred}] = 1$  if  $\text{pred}$  is a logical predicate which is true and  $[\text{pred}] = 0$  otherwise.

6. [10 pts] *Bayes estimator.* In both of the following subquestions you will need to find the Bayes estimator with respect to the the probability mass function  $p(x, y)$  over  $(X, Y)$  where  $X$  and  $Y$  are finite thus  $\sum_{x \in X} \sum_{y \in Y} p(x, y) = 1$ .

- (a) For this subquestion  $Y = [k]$  and let  $\mathbf{c} \in [0, \infty)^k$  be a vector of  $k$  costs. Define  $L_{\mathbf{c}} : [k] \times [k] \rightarrow [0, \infty)$  as,

$$L_{\mathbf{c}}(y, \hat{y}) := [y \neq \hat{y}]c_y$$

as the the imbalanced classification loss function, i.e., if we don't predict the correct outcome  $y$  we suffer  $c_y$  loss. Derive the Bayes estimator.

(b) For this subquestion  $Y \subset \mathfrak{R}$ . Let  $L(y, \hat{y}) := |y - \hat{y}|$ . Derive the Bayes estimator.

7. [10 pts] *Kernel modification* Consider the function  $K_c(\mathbf{x}, \mathbf{z}) := c + \sum_{i=1}^n x_i z_i$  where  $\mathbf{x}, \mathbf{z} \in \mathfrak{R}^n$ .


(a) For what values of  $c \in \mathfrak{R}$  is  $K_c$  a positive semidefinite kernel? Give an argument supporting your claim. ("The closer your argument is to a proof the more likely it is to receive full credit.")

(b) Suppose we use  $K_c$  as a kernel function with linear regression (least squares). Explain how  $c$  influences the solution.


8. [15 pts] Suppose we perform linear regression with a Gaussian kernel  $K_\beta(\mathbf{x}, \mathbf{t}) = \exp(-\beta \|\mathbf{x} - \mathbf{t}\|^2)$  to train a classifier for two-class data (i.e.,  $y \in \{-1, 1\}$ ). This classifier depends on the parameter  $\beta$  selected for the kernel. How should one choose  $\beta$  so that the learned linear classifier simulates a 1-NEAREST NEIGHBOR CLASSIFIER? Give an argument supporting your reasoning. ("The closer your argument is to a proof the more likely it is to receive full credit'.')

9. [15pts]: *Hint: "Think about vector spaces 'broadly'"*.


We consider the problem of generalized Whack-A-Mole! We'll play this game on a board with  $n \times n$  holes, from which individual moles pop-up. Using a mallet you can then whack a mole and make it go hide underground. As you might know, however, whacking a mole will often have the effect of making other moles rise and come out from the ground. In this version of the game, every time you hit a hole with the mallet you'll cause the hole and the immediate four adjacent holes to change: any moles currently there will hide, while new moles pop-up from previously empty holes. Imagine, for example, that you find yourself in the following game configuration, and you prepare yourself to hit the hole at position 10 with the mallet.



You will cause the moles in positions 9, 10, 11, and 14 to hide, but you'll also make a mole appear at position 6. Thus ending up in the next configuration.



**Example:** Consider the  $4 \times 4$  board with moles at positions 1,2,4,8,9,10,11, and 14. This is the same configuration of moles as given in the initial illustration above. After hitting holes at positions 10 then 2, you'll end up in the following configuration.



**Task:** Design an algorithm for an  $n \times n$  board which, given an initial board configuration, finds a sequence of holes that you can hit in order to empty the board if such a sequence exists. The algorithm must be polynomial in  $n$  and you must provide an argument that it is correct. No credit will be given for algorithms that are not polynomial in  $n$ . There is no need to implement your algorithm.