

class10: Structural Bioinformatics (Pt. 1)

Jiachen Fan (A17662703)

1: Introduction to the RCSB Protein Data Bank (PDB)

Download stats from '<https://www.rcsb.org/stats/summary>'

At the time of writing there are 183,201 protein structure. In UniProt there are 251600,768 protein sequences. Only a small amount of proteins structure are available.

```
round(183201/251600768*100, 2)
```

```
[1] 0.07
```

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
stats <- read.csv('Data Export Summary.csv', row.names = 1)
head(stats)
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158,844	11,759	12,296	197	73	32
Protein/Oligosaccharide	9,260	2,054	34	8	1	0
Protein/NA	8,307	3,667	284	7	0	0
Nucleic acid (only)	2,730	113	1,467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183,201					
Protein/Oligosaccharide	11,357					
Protein/NA	12,265					
Nucleic acid (only)	4,327					
Other	205					
Oligosaccharide (only)	22					

We want to generate a function and apply it to our files. We can use the 'gsub()' function to remove commas.

```
string <- c('10', '100', 1, '1,000')
x <- string
as.numeric(gsub(",", "", x))
```

```
[1] 10 100 1 1000
```

```
rm.comma <- function(x){
  as.numeric(gsub(",", "", x))
}
```

```
pdbstats <- apply(stats, 2, rm.comma) # '2' means apply to column
pdbstats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other	Total
[1,]	158844	11759	12296	197	73	32	183201
[2,]	9260	2054	34	8	1	0	11357
[3,]	8307	3667	284	7	0	0	12265
[4,]	2730	113	1467	13	3	1	4327
[5,]	164	9	32	0	0	0	205
[6,]	11	0	6	1	0	4	22

We want our rownames back.

```
rownames(pdbstats) <- rownames(stats)
pdbstats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158844	11759	12296	197	73	32
Protein/Oligosaccharide	9260	2054	34	8	1	0
Protein/NA	8307	3667	284	7	0	0
Nucleic acid (only)	2730	113	1467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183201					
Protein/Oligosaccharide	11357					

Protein/NA	12265
Nucleic acid (only)	4327
Other	205
Oligosaccharide (only)	22

```
totals <- apply(pdbstats, 2, sum)
round(totals/totals["Total"]*100,2)
```

X.ray	EM	NMR	Multiple.methods
84.83	8.33	6.68	0.11
Neutron	Other	Total	
0.04	0.02	100.00	

It is 84.83% for X.ray and 8.33% for EM.

Q2-3 Lets skip these....

Q2: What proportion of structures in the PDB are protein?

```
round(pdbstats['Total'][1]/sum(pdbstats['Total'])*100, 2)
```


[1] NA

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?


22824 structures.

Using Mol* to examine HIV-Pr

Here is a rubbish pic of HIV-Pr that is not very useful yet.



Add a nicer pic colored by secondary structure with catalytic active site ASP 25 shown in each chain



> Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

Because it is 'ball&stick' type, the water molecules are represented by 'O'

Q5: There is a critical "conserved" water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have

HOH 332.

Using the bio3d package

```
library(bio3d)

pdb <- read.pdb('1hsg')
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD  
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE  
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP  
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

Q7: How many amino acid residues are there in this pdb object?

198 amino acid residues.

Q8: Name one of the two non-protein residues?

HOH

Q9: How many protein chains are in this structure?

2

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

```
head(pdb$atom$resid)
```

```
[1] "PRO" "PRO" "PRO" "PRO" "PRO" "PRO"
```

```
pdb$atom$resid[pdb$calpha]
```

```
[1] "PRO" "GLN" "ILE" "THR" "LEU" "TRP" "GLN" "ARG" "PRO" "LEU" "VAL" "THR"
[13] "ILE" "LYS" "ILE" "GLY" "GLY" "GLN" "LEU" "LYS" "GLU" "ALA" "LEU" "LEU"
[25] "ASP" "THR" "GLY" "ALA" "ASP" "ASP" "THR" "VAL" "LEU" "GLU" "GLU" "MET"
[37] "SER" "LEU" "PRO" "GLY" "ARG" "TRP" "LYS" "PRO" "LYS" "MET" "ILE" "GLY"
[49] "GLY" "ILE" "GLY" "GLY" "PHE" "ILE" "LYS" "VAL" "ARG" "GLN" "TYR" "ASP"
[61] "GLN" "ILE" "LEU" "ILE" "GLU" "ILE" "CYS" "GLY" "HIS" "LYS" "ALA" "ILE"
[73] "GLY" "THR" "VAL" "LEU" "VAL" "GLY" "PRO" "THR" "PRO" "VAL" "ASN" "ILE"
[85] "ILE" "GLY" "ARG" "ASN" "LEU" "LEU" "THR" "GLN" "ILE" "GLY" "CYS" "THR"
[97] "LEU" "ASN" "PHE" "PRO" "GLN" "ILE" "THR" "LEU" "TRP" "GLN" "ARG" "PRO"
[109] "LEU" "VAL" "THR" "ILE" "LYS" "ILE" "GLY" "GLY" "GLN" "LEU" "LYS" "GLU"
[121] "ALA" "LEU" "LEU" "ASP" "THR" "GLY" "ALA" "ASP" "ASP" "THR" "VAL" "LEU"
[133] "GLU" "GLU" "MET" "SER" "LEU" "PRO" "GLY" "ARG" "TRP" "LYS" "PRO" "LYS"
[145] "MET" "ILE" "GLY" "GLY" "ILE" "GLY" "GLY" "PHE" "ILE" "LYS" "VAL" "ARG"
[157] "GLN" "TYR" "ASP" "GLN" "ILE" "LEU" "ILE" "GLU" "ILE" "CYS" "GLY" "HIS"
[169] "LYS" "ALA" "ILE" "GLY" "THR" "VAL" "LEU" "VAL" "GLY" "PRO" "THR" "PRO"
[181] "VAL" "ASN" "ILE" "ILE" "GLY" "ARG" "ASN" "LEU" "LEU" "THR" "GLN" "ILE"
[193] "GLY" "CYS" "THR" "LEU" "ASN" "PHE"
```

Predicting functional motions of a single structure

Run a Normal Mode Analysis (NMA)

```
adk <- read.pdb('6s36')
```

Note: Accessing on-line PDB file


PDB has ALT records, taking A only, rm.alt=TRUE

```
m <- nma(adk)
```

Building Hessian... Done in 0.04 seconds.

Diagonalizing Hessian... Done in 0.36 seconds.

```
plot(m)
```



```
mktrj(m, file="m.pdb")
```