



K-Means connection



K-Means

1. Randomly initialize parameters $\theta = \{\mu_1, \dots, \mu_C\}$

K-Means

1. Randomly initialize parameters $\theta = \{\mu_1, \dots, \mu_C\}$
2. Until convergence repeat:
 - a) For each point compute closest centroid

$$c_i = \arg \min_c \|x_i - \mu_c\|^2$$

K-Means

1. Randomly initialize parameters $\theta = \{\mu_1, \dots, \mu_C\}$
2. Until convergence repeat:
 - a) For each point compute closest centroid

$$c_i = \arg \min_c \|x_i - \mu_c\|^2$$

- b) Update centroids

$$\mu_c = \frac{\sum_{i:c_i=c} x_i}{\#\{i : c_i = c\}}$$

K-Means from GMM perspective

From GMM to K-means:

- Fix covariances to be identical $\Sigma_c = I$

K-Means from GMM perspective

From GMM to K-means:

- Fix covariances to be identical $\Sigma_c = I$
- Fix weights to be uniform $\pi_c = \frac{1}{\# \text{ of Gaussians}}$

K-Means from GMM perspective

From GMM to K-means:

- Fix covariances to be identical $\Sigma_c = I$
- Fix weights to be uniform $\pi_c = \frac{1}{\# \text{ of Gaussians}}$

$$p(x_i \mid t_i = c, \theta) = \frac{1}{Z} \exp(-0.5 \|x_i - \mu_c\|^2)$$

K-Means from EM perspective

E-step

$$q^{k+1} = \arg \min_q \mathcal{KL} [q(T) \parallel p(T \mid X, \theta^k)]$$

K-Means from EM perspective

E-step

$$q^{k+1} = \arg \min_{q \in Q} \mathcal{KL} [q(T) \parallel p(T \mid X, \theta^k)]$$


Where Q is the set of delta-functions

K-Means from EM perspective

E-step

$$q^{k+1} = \arg \min_{q \in \mathcal{Q}} \mathcal{KL} [q(T) \parallel p(T \mid X, \theta^k)]$$

Where \mathcal{Q} is the set of delta-functions




K-Means from EM perspective

E-step

$$q^{k+1} = \arg \min_{q \in \mathcal{Q}} \mathcal{KL} [q(T) \parallel p(T \mid X, \theta^k)]$$

Where \mathcal{Q} is the set of delta-functions



K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \max_c p(t_i = c \mid x_i, \theta)$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \max_c p(t_i = c \mid x_i, \theta)$$

$$p(t_i \mid x_i, \theta) = \frac{1}{Z} p(x_i \mid t_i, \theta) p(t_i \mid \theta)$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \max_c p(t_i = c \mid x_i, \theta)$$

$$\begin{aligned} p(t_i \mid x_i, \theta) &= \frac{1}{Z} p(x_i \mid t_i, \theta) p(t_i \mid \theta) \\ &= \frac{1}{Z} \exp(-0.5 \|x_i - \mu_c\|^2) \pi_c \end{aligned}$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \max_c p(t_i = c \mid x_i, \theta) = \arg \min_c \|x_i - \mu_c\|^2$$

$$\begin{aligned} p(t_i \mid x_i, \theta) &= \frac{1}{Z} p(x_i \mid t_i, \theta) p(t_i \mid \theta) \\ &= \frac{1}{Z} \exp(-0.5\|x_i - \mu_c\|^2) \pi_c \end{aligned}$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \min_c \|x_i - \mu_c\|^2$$

K-Means from EM perspective

E-step

$$q^{k+1}(t_i) = \begin{cases} 1 & \text{if } t_i = c_i \\ 0 & \text{otherwise} \end{cases}$$

$$c_i = \arg \min_c \|x_i - \mu_c\|^2$$

Exactly like in K-Means!