

0.0.1 Renormalization-scale dependence

In this section we consider renormalization-scale dependence of our results. In principle, such scale μ should not appear in the cross sections, as it does not correspond to any fundamental constant or kinematical scale in the problem. The completely resummed perturbative expansion of an observable is indeed formally independent on μ . In practise, truncated perturbative expansions exhibit a residual scale dependence, because of neglected higher orders.

We start with deriving the strong coupling $\alpha_s(Q^2)$ as a function of $\alpha_s(\mu^2)$ and μ^2/Q^2 , to do so we expand the explicit ?? in powers of $\alpha_s(\mu^2)$ and obtain:

$$\begin{aligned} \alpha_s(Q^2) = & \alpha_s(\mu^2) + \alpha_s^2(\mu^2)b_0 \ln\left(\frac{\mu^2}{Q^2}\right) + \alpha_s^3(\mu^2)\left[b_1 \ln\left(\frac{\mu^2}{Q^2}\right) + b_0^2 \ln^2\left(\frac{\mu^2}{Q^2}\right)\right] \\ & + \alpha_s^4(\mu^2)\left[b_2 \ln\left(\frac{\mu^2}{Q^2}\right) + \frac{5}{2}b_0b_1 \ln^2\left(\frac{\mu^2}{Q^2}\right) + b_0^3 \ln^3\left(\frac{\mu^2}{Q^2}\right)\right] \\ & + \alpha_s^5(\mu^2)\left[b_3 \ln\left(\frac{\mu^2}{Q^2}\right) + \left(\frac{3}{2}b_1^2 + 3b_0b_2\right) \ln^2\left(\frac{\mu^2}{Q^2}\right) \right. \\ & \left. + \frac{13}{3}b_0^2b_1 \ln^3\left(\frac{\mu^2}{Q^2}\right) + b_0^4 \ln^4\left(\frac{\mu^2}{Q^2}\right)\right] + \mathcal{O}(\alpha_s^6(\mu^2)) \end{aligned} \quad (1)$$

for brevity we'll write:

$$\alpha_s(Q^2) = \alpha_s(\mu^2) + c_1\alpha_s^2(\mu^2) + c_2\alpha_s^3(\mu^2) + c_3\alpha_s^4(\mu^2) + c_4\alpha_s^5(\mu^2) + \mathcal{O}(\alpha_s^6(\mu^2)) \quad (2)$$

where c_i are the coefficients of the expansion above.

$$\begin{aligned} c_1 &= b_0 \ln\left(\frac{\mu^2}{Q^2}\right) \\ c_2 &= b_1 \ln\left(\frac{\mu^2}{Q^2}\right) + b_0^2 \ln^2\left(\frac{\mu^2}{Q^2}\right) \\ c_3 &= b_2 \ln\left(\frac{\mu^2}{Q^2}\right) + \frac{5}{2}b_0b_1 \ln^2\left(\frac{\mu^2}{Q^2}\right) + b_0^3 \ln^3\left(\frac{\mu^2}{Q^2}\right) \\ c_4 &= b_3 \ln\left(\frac{\mu^2}{Q^2}\right) + \left(\frac{3}{2}b_1^2 + 3b_0b_2\right) \ln^2\left(\frac{\mu^2}{Q^2}\right) + \frac{13}{3}b_0^2b_1 \ln^3\left(\frac{\mu^2}{Q^2}\right) + b_0^4 \ln^4\left(\frac{\mu^2}{Q^2}\right) \end{aligned} \quad (3)$$

now we substitute eq. (1) into $\lambda = b_0\alpha_s(Q^2)L$ and obtain:

$$\lambda(Q^2) = \lambda(\mu^2)\left(1 + c_1\alpha_s(\mu^2) + c_2\alpha_s^2(\mu^2) + c_3\alpha_s^3(\mu^2) + c_4\alpha_s^4(\mu^2) + \mathcal{O}(\alpha_s^5(\mu^2))\right) \quad (4)$$

and formally expand in powers of $\alpha_s(\mu^2)$ all the relevant functions, now $\lambda = \lambda(\mu^2)$:

$$\begin{aligned}
Lf_1(Q^2) &= Lf_1(\lambda) + \frac{c_1}{b_0}\lambda^2 f_1'(\lambda) + \frac{\lambda}{b_0} \left(c_2 \lambda f_1'(\lambda) + \frac{1}{2} c_1^2 \lambda^2 f_1''(\lambda) \right) \alpha_s \\
&\quad + \frac{\lambda}{b_0} \left(c_3 \lambda f_1'(\lambda) + c_1 c_2 \lambda^2 f_1''(\lambda) + \frac{1}{6} c_1^3 f_1^{(3)}(\lambda) \right) \alpha_s^2 + \frac{\lambda^2}{b_0} \left(c_4 f_1'(\lambda) \right. \\
&\quad \left. + \frac{1}{2} c_2^2 \lambda f_1''(\lambda) + c_1 c_3 \lambda f_1''(\lambda) + \frac{1}{2} c_1^2 c_2 \lambda^2 f_1^{(3)}(\lambda) + \frac{1}{24} c_1^4 \lambda^3 f_1^{(4)}(\lambda) \right) \alpha_s^3 \\
f_2(Q^2) &= f_2(\lambda) + c_1 \lambda f_2'(\lambda) \alpha_s + \left(c_2 \lambda f_2'(\lambda) + \frac{1}{2} c_1^2 \lambda^2 f_2''(\lambda) \right) \alpha_s^2 \\
&\quad + \left(c_3 \lambda f_2'(\lambda) + c_1 c_2 \lambda^2 f_2''(\lambda) + \frac{1}{6} c_1^3 f_2^{(3)}(\lambda) \right) \alpha_s^3 \\
\alpha_s f_3(Q^2) &= \alpha_s f_3(\lambda) + c_1 \lambda f_3'(\lambda) \alpha_s^2 + \left(c_2 \lambda f_3'(\lambda) + \frac{1}{2} c_1^2 \lambda^2 f_3''(\lambda) \right) \alpha_s^3 \\
\alpha_s^2 f_4(Q^2) &= \alpha_s^2 f_4(\lambda) + c_1 \lambda f_4'(\lambda) \alpha_s^3 \\
\alpha_s^3 f_5(Q^2) &= \alpha_s^3 f_5(\lambda)
\end{aligned} \tag{5}$$

As usual, the terms from the expansion proportional to α_s corrects f_3 , the terms proportional to α_s^2 corrects f_4 and so on.

So the additional terms in the functions f_i for $i = 0, 1, 3$, to partially compensate for the scale change $Q^2 \rightarrow \mu^2$, read:

$$\begin{aligned}
\delta f_1\left(\lambda, \frac{\mu^2}{Q^2}\right) &= 0 \\
\delta f_2\left(\lambda, \frac{\mu^2}{Q^2}\right) &= \lambda^2 f_1'(\lambda) \log \frac{\mu^2}{Q^2} \\
\delta f_3\left(\lambda, \frac{\mu^2}{Q^2}\right) &= \frac{1}{2} b_0 \lambda^3 f_1''(\lambda) \log^2 \frac{\mu^2}{Q^2} + \lambda^2 \left(b_0 f_1'(\lambda) \log^2 \frac{\mu^2}{Q^2} + \frac{b_1}{b_0} f_1'(\lambda) \log \frac{\mu^2}{Q^2} \right) \\
&\quad + b_0 \lambda f_2'(\lambda) \log \frac{\mu^2}{Q^2} \\
\delta f_4\left(\lambda, \frac{\mu^2}{Q^2}\right) &= \frac{1}{6} b_0^2 \lambda^4 f_1^{(3)}(\lambda) \log^3 \frac{\mu^2}{Q^2} + \lambda^3 \left(b_0^2 f_1''(\lambda) \log^3 \frac{\mu^2}{Q^2} + b_1 f_1''(\lambda) \log^2 \frac{\mu^2}{Q^2} \right) \\
&\quad + \lambda^2 \left(b_0^2 f_1'(\lambda) \log^3 \frac{\mu^2}{Q^2} + \frac{5}{2} b_1 f_1'(\lambda) \log^2 \frac{\mu^2}{Q^2} + \frac{b_2}{b_0} f_1'(\lambda) \log \frac{\mu^2}{Q^2} + \frac{1}{2} b_0^2 f_2''(\lambda) \log^2 \frac{\mu^2}{Q^2} \right) \\
&\quad + \lambda \left(b_0^2 f_2'(\lambda) \log^2 \frac{\mu^2}{Q^2} + b_0 f_3'(\lambda) \log \frac{\mu^2}{Q^2} + b_1 f_2'(\lambda) \log \frac{\mu^2}{Q^2} \right) \\
\delta f_5\left(\lambda, \frac{\mu^2}{Q^2}\right) &= \frac{1}{24} b_0^3 \lambda^5 f_1^{(4)}(\lambda) \log^4 \frac{\mu^2}{Q^2} + \lambda^4 \left(\frac{1}{2} b_0^3 f_1^{(3)}(\lambda) \log^4 \frac{\mu^2}{Q^2} \right. \\
&\quad + \frac{1}{2} b_0 b_1 f_1^{(3)}(\lambda) \log^3 \frac{\mu^2}{Q^2} \left. \right) + \lambda^3 \left(\frac{3}{2} b_0^3 f_1''(\lambda) \log^4 \frac{\mu^2}{Q^2} + \frac{7}{2} b_0 b_1 f_1''(\lambda) \log^3 \frac{\mu^2}{Q^2} \right. \\
&\quad + \frac{b_1^2}{2b_0} f_1''(\lambda) \log^2 \frac{\mu^2}{Q^2} + b_2 f_1''(\lambda) \log^2 \frac{\mu^2}{Q^2} + \frac{1}{6} b_0^3 f_2^{(3)}(\lambda) \log^3 \frac{\mu^2}{Q^2} \left. \right) \\
&\quad + \lambda^2 \left(b_0^3 f_1'(\lambda) \log^4 \frac{\mu^2}{Q^2} + \frac{13}{3} b_0 b_1 f_1'(\lambda) \log^3 \frac{\mu^2}{Q^2} + \frac{3b_1^2}{2b_0} f_1'(\lambda) \log^2 \frac{\mu^2}{Q^2} \right. \\
&\quad + 3b_2 f_1'(\lambda) \log^2 \frac{\mu^2}{Q^2} + \frac{b_3}{b_0} f_1'(\lambda) \log \frac{\mu^2}{Q^2} + b_0^3 f_2''(\lambda) \log^3 \frac{\mu^2}{Q^2} + b_0 b_1 f_2''(\lambda) \log^2 \frac{\mu^2}{Q^2} \\
&\quad + \frac{1}{2} b_0^2 f_3''(\lambda) \log^2 \frac{\mu^2}{Q^2} \left. \right) + \lambda \left(b_0^3 f_2'(\lambda) \log^3 \frac{\mu^2}{Q^2} + b_0^2 f_3'(\lambda) \log^2 \frac{\mu^2}{Q^2} \right. \\
&\quad + \frac{5}{2} b_1 b_0 f_2'(\lambda) \log^2 \frac{\mu^2}{Q^2} + b_0 f_4'(\lambda) \log \frac{\mu^2}{Q^2} + b_2 f_2'(\lambda) \log \frac{\mu^2}{Q^2} + b_1 f_3'(\lambda) \log \frac{\mu^2}{Q^2} \left. \right)
\end{aligned}$$

(6)