



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Please write carefully and clearly in complete sentences. Your explanations are your only representative when your work is being graded.

1. Solve the following problems:

(a) $y^{(4)} + 2y'' + y = 8 \sin t - 16 \cos t$;

(b) $y''' - 3y'' + 3y' - y = 6e^t$;

(c) $y' + y^2 \sin x = 0$.

Solutions.

- (a) The characteristic equation is $r^4 + 2r^2 + 1 = 0$. The roots are $i, i, -i, -i$, and the general solution of the homogeneous equation is

$$y_c(t) = c_1 \cos t + c_2 \sin t + c_3 t \cos t + c_4 t \sin t.$$

Our initial assumption for a particular solution is $Y(t) = A \sin t + B \cos t$, but we must multiply this choice by t^2 to make this choice by t^2 to make it different from all solutions of the homogeneous equation. Thus our assumption is $Y(t) = At^2 \sin t + Bt^2 \cos t$. Plugging this into the original equation, we obtain $A = -1$, $B = 2$. Hence, the general solution of the nonhomogeneous equation is

$$y(t) = y_c(t) + Y(t) = c_1 \cos t + c_2 \sin t + c_3 t \cos t + c_4 t \sin t - t^2 \sin t + 2t^2 \cos t,$$

where c_1, c_2, c_3, c_4 are arbitrary constants.

- (b) The characteristic equation is $r^3 - 3r^2 + 3r - 1 = (r - 1)^3$. So, the general solution of the homogeneous equation is

$$y_c(t) = c_1 e^t + c_2 t e^t + c_3 t^2 e^t,$$

where c_1, c_2, c_3 are arbitrary constants. Since $e^t, t e^t, t^2 e^t$ are all solutions of the homogeneous equation, we need to find a particular solution of the form $Y(t) = At^3 e^t$ of the nonhomogeneous equation, where A is an

undetermined coefficient. Plugging this into the original equation, we obtain $A = 1$. In conclusion, the general solution of the nonhomogeneous equation is

$$y(t) = c_1 e^t + c_2 t e^t + c_3 t^2 e^t + t^3 e^t,$$

where c_1, c_2, c_3 are arbitrary constants.

(c) If $y \neq 0$, then

$$-\frac{1}{y^2} dy = \sin x dx \Rightarrow d(y^{-1}) = -d(\cos x) \Rightarrow y^{-1} + \cos x = c,$$

where c is an arbitrary constant. Also, $y = 0$ is a solution.

2. Use the method of undetermined coefficients to find a particular solution of

$$\mathbf{x}' = \begin{bmatrix} -2 & 1 \\ 1 & -2 \end{bmatrix} \mathbf{x} + \begin{bmatrix} 2e^{-t} \\ 3t \end{bmatrix}.$$

Solution. To use the method of undetermined coefficients, we write $\mathbf{g}(t)$ in the form

$$\mathbf{g}(t) = \begin{bmatrix} 2 \\ 0 \end{bmatrix} e^{-t} + \begin{bmatrix} 0 \\ 3 \end{bmatrix} t.$$

Note that -1 is an eigenvalue of the coefficient matrix, and therefore we must include both $\mathbf{a}te^{-t}$ and $\mathbf{b}e^{-t}$ in the form of a solution. Therefore the solution of the system can be assumed to be of the form

$$\mathbf{x} = \mathbf{v}(t) = \mathbf{a}te^{-t} + \mathbf{b}e^{-t} + \mathbf{c}t + \mathbf{d}$$

where $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are vectors to be determined. Plugging the above $\mathbf{x}(t)$ into the original system, we obtain

$$\begin{aligned} \mathbf{x}' &= -\mathbf{a}te^{-t} + (\mathbf{a} - \mathbf{b})e^{-t} + \mathbf{c} \\ &= \mathbf{A}\mathbf{a}te^{-t} + \left(\mathbf{A}\mathbf{b} + \begin{bmatrix} 2 \\ 0 \end{bmatrix} \right) e^{-t} + \left(\mathbf{A}\mathbf{c} + \begin{bmatrix} 2 \\ 0 \end{bmatrix} \right) t + \mathbf{A}\mathbf{d}. \end{aligned}$$

Equating all the corresponding coefficients, we obtain

$$\mathbf{a} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b} = k \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \mathbf{c} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{d} = \begin{bmatrix} -\frac{4}{3} \\ -\frac{5}{3} \end{bmatrix},$$

where k is an arbitrary constant. We can simply take $k = 0$, and thus a particular solution of the previous system is given as follows:

$$\mathbf{v}(t) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} te^{-t} - \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^{-t} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} t - \frac{1}{3} \begin{bmatrix} 4 \\ 5 \end{bmatrix}.$$

3. Determine the type and stability of the critical point $(4, 3)$ of the locally linear system

$$\begin{aligned} \frac{dx}{dt} &= 33 - 10x - 3y + x^2, \\ \frac{dy}{dt} &= -18 + 6x + 2y - xy. \end{aligned}$$

Solution. With $F(x, y) = 33 - 10x - 3y + x^2$, $G(x, y) = -18 + 6x + 2y - xy$ and $x_0 = 4, y_0 = 3$ we have

$$J(x, y) = \begin{bmatrix} -10 + 2x & -3 \\ 6 - y & 2 - x \end{bmatrix}, \quad J(4, 3) = \begin{bmatrix} -2 & -3 \\ 3 & -2 \end{bmatrix}.$$

The associated linear system

$$\begin{aligned} \frac{du}{dt} &= -2u - 3v, \\ \frac{dv}{dt} &= 3u - 2v \end{aligned}$$

has characteristic equation $(\lambda + 2)^2 + 9 = 0$, with complex conjugate roots $\lambda = -2 \pm 3i$. Hence, $(0, 0)$ is an asymptotically stable spiral of the linear system. It follows that $(4, 3)$ is an asymptotically stable spiral point of the original locally linear system.

4. Construct a suitable Liapunov function to determine the stability of the origin for the system

$$\begin{aligned} \frac{dx}{dt} &= x^3 - 2y^3, \\ \frac{dy}{dt} &= xy^2 + x^2y + \frac{1}{2}y^3. \end{aligned}$$

Solution. We can take $V(x, y) = x^2 + 2y^2$ and $\dot{V} = 2(x^2 + y^2)^2$. Both of them are positive definite, therefore the origin is unstable.

5. Consider the system

$$\mathbf{x}' = \begin{bmatrix} 2 & 3 \\ -1 & -2 \end{bmatrix} \mathbf{x}$$

Find a fundamental matrix.

Solution. A fundamental matrix is given as follows:

$$\Psi(t) = \begin{bmatrix} -3e^t & e^{-t} \\ e^t & -e^{-t} \end{bmatrix}.$$

6. Transform $u'' - 2u' - 3u = 0$ into a system of first order equations.

Solution.

$$\begin{cases} x_1' = x_2 \\ x_2' = 2x_2 + 3x_1. \end{cases}$$

7. If two functions $f(t)$ and $g(t)$ are linearly independent on the interval $(-1, 1)$, then the Wronskian $W[f, g](t) \neq 0$ for all $(-1, 1)$. True or False?

Solution. False.

8. Write down the definition of the matrix exponential e^{At} .

Solution.

$$e^{At} = \mathbf{I} + \sum_{k=1}^{\infty} \frac{\mathbf{A}^k t^k}{k!}.$$

9. All solutions of $y' + 2y = \sin t$ converges to a particular solution as $t \rightarrow \infty$. True or False.

Solution. True.

10. Can be a saddle point asymptotically stable in a particular direction? Yes or No?

Solution. Yes.

11. Suppose that $f(x, y)$ and $f_y(x, y)$ are continuous on the xy -plane. Is it possible that $y_1(x) = \sin x$ and $y_2(x) = \cos x$ are solutions of $y'(x) = f(x, y)$? Yes or no?

Solution. No.

12. Use Laplace transform to solve the following initial value problem

$$\begin{cases} y^{(4)} + 2y'' + y = 4te^t, \\ y(0) = y'(0) = y''(0) = y'''(0) = 0. \end{cases}$$

Solution.

$$Y(s) = \frac{4}{(s-1)^2(s^2+1)^2} = \frac{A}{(s-1)^2} + \frac{B}{s-1} + \frac{Cs+D}{(s^2+1)^2} + \frac{Es+F}{s^2+1}.$$

Solve for $B = -2$, $C = 2$, $D = 0$, $E = 2$, and $F = 1$. Therefore, the solution of the given initial value problem is

$$y(t) = (t-2)e^t + (t+1)\sin t + 2\cos t.$$